

中華昆蟲 9: 87-136 (1989)
Chinese J. Entomol. 9: 87-136 (1989)

A LIST OF CERAMBYCID-BEETLES FROM TAIWAN (II)⁽¹⁾

Subfamily Lamiinae

Shingo Nakamura⁽²⁾, Hiroshi Makihara⁽³⁾, Akiko Saito⁽⁴⁾,
Yau-I Chu⁽⁵⁾ and Masuo Hayashi⁽⁶⁾

(Accepted for publication February 20, 1988)

Till present, 311 species of Cerambycid beetles which belong to subfamily Lamiinae are recorded from Taiwan. They are listed in the following catalogue:

Subfamily Lamiinae 粗天牛亞科

Tribe Mesosini 星斑天牛族

Genus *Mesoereis* MATSUSHITA, 1933 大星斑天牛屬

Mesoereis MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 338 (Type-species: *M. koshunensis* MATSUSHITA, 1933).

Mesoereis koshunensis MATSUSHITA, 1933 恒春大星斑天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 339, pl. 5, f. 6 (Koshun).

Distr.: Taiwan.

M. obscurus MATSUSHITA, 1933 黃翅大星斑天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 339, pl. 5, f. 5 (Hori).

Distr.: Taiwan.

Genus *Falsomesosella* PIC, 1925 假星斑天牛屬

Falsomesosella PIC, 1925, Mel. Exot.-Ent., 44: 27 (Type-species: *F. albofasciata* PIC, 1925).

(1) During the course of the present work, the authors debited for the co-operation on the following colleague to whom our sincerely thanks are due, Drs. T. Niisato, N. Obayashi, T. Shimomura, and Miss M. J. Lin. Also this work is financially supported by National Science Council, R. O. C. under the project No. of NSC 75-0409-B002-47.

(2) 179 Setoyama, Moto-machi, Shobara, Hiroshima, Japan.

(3) Sect. Forest Entomology, Forestry and Forest Products Research Institute, Japan.

(4) Div. of Entomology, National Institute of Agro-Environmental Sciences, Japan.

(5) Dept. of Plant Pathology and Entomology, National Taiwan University, Taipei, Taiwan, R. O. C.

(6) Osaka Jonan Women's Junior College, Japan.

Falsomesosella horishana GRESSITT, 1938 埔里姬星斑天牛

GRESSITT, 1938, Philip. J. Sci., 65: 161, pl. 1, f. 9 (Hori).

Distr.: Taiwan.

F. subalba GRESSITT, 1938 白姬星斑天牛

GRESSITT, 1938, Philip. J. Sci., 65: 162, pl. 1, f. 10 (Kuraru).

Distr.: Taiwan.

Genus *Mesosa* LATREILLE, 1829 星斑天牛屬

Mesosa LATREILLE, 1829, in Cuvier, Regne Anim. Ins. 2: 124 (Type-species: *Lamia cruculionoides* LINNE, 1761).

Subgenus *Mesosa* LATREILLE, 1829, s. str.

Mesosa (*Mesosa*) *perplexa* PASCOE, 1858 褐翅星斑天牛

PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 243 (N. China).

alternans SCHWARZER, 1925, Ent. Blaett., 21: 62 (*Saimia*) (Kankau, Kosempo, Pilam).

?*formosana* PIC, 1925, Mel. Exot.-Ent., 44: 30 (?*Mimocoptops*) (Taiwan).

Distr.: Taiwan; Japan, E. China.

Subgenus *Aphelocnemia* STEPHENS, 1831

Aphelocnemia STEPHENS, 1831, Brit. Ent. Man. Errata, p. 414 (Type-species: *Lamia nebulosa* FABRICIUS, 1781).

Mesosa AURIVILLIUS, 1922, Coleop. Cat., (73): 137 (nec LATREILLE, 1829).

Helixoea PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 124 (Type-species: *Agelasta rupta* PASCOE, 1862).

Haplocnemia GEMMINGER et HAROLD, 1873, Cat. Col., 10: 3038.

M. (*Aphelocnemia*) *latifasciata* (WHITE, 1858) 寬帶星斑天牛

WHITE, 1858, Proc. Zool. Soc. London, 26: 401 (*Cacia*) (Shanghai).

luteopubens PIC, 1917, Mel. Exot.-Ent., 23: 7 (Tonkin).

Distr.: Taiwan; E. China, Tonkin.

Subgenus *Perimesosa* BREUNING, 1939

Mesosa subgen. *Perimesosa* BREUNING, 1939, Novit. Ent., 9, suppl. 3, fasc. (52): 409 (Type-species: *Mesosa hirsuta* BATES, 1884).

M. (*Perimesosa*) *atrostigma* GRESSITT, 1942 黑點星斑天牛

GRESSITT, 1942, Notes d'Ent. Chinoise, 9: 84, pl. 1, f. 4 (China).

Distr.: Taiwan; E. China.

M. (*P.*) *binigrovittata* BREUNING, 1942 雙點星斑天牛

BREUNING, 1942, Novit. Ent., 11-12: 143 (Hokuzanko).

Distr.: Taiwan.

M. (*P.*) *chassoti* BREUNING, 1970 夏叟氏星斑天牛

BREUNING 1970, Bull. Soc. Ent. Mulhouse, p. 89 (Taiwan).

Distr.: Taiwan.

Subgenus *Saimia* PASCOE, 1866

Saimia PASCOE, 1866, Proc. Zool. Soc. London, p. 234 (Type-species: *Saimia albidor-salis* PASCOE, 1864).

Pachyosa FAIRMAIRE, 1897, Bull. Soc. Ent. Fr., p. 71 (Type-species: *Pachyosa cervinopicta* FAIRMAIRE, 1897).

M. (*Saimia*) *kojimai* (HAYASHI, 1974) 小島氏星斑天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 37 (*Coptops*) (Liukuei).

Distr.: Taiwan.

M. (*S.*) *kuntzeni* MATSUSHITA, 1933 臺灣白條星斑天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ. 34: 342 (Hozan).

Distr.: Taiwan.

Genus *Cacia* NEWMAN, 1842 鬚角星斑天牛屬

Cacia NEWMAN, 1842, Entomol. 1: 290 (Type-species: *C. spinigera* NEWMAN, 1842).

Subgenus *Ipocregyes* PASCOE, 1866

Ipocregyes PASCOE, 1866, Trans. Ent. Soc. London, (3)3: 113 (Type-species: *I. newmanni* PASCOE, 1866).

Cacia (*Ipocregyes*) *arisana* (KANO, 1933) 阿里山刷角星斑天牛

KANO, 1933, Kontyû, 6: 281 (*Ipocregyes arisanus*) (Arisan).

scopiferus MITONO, 1934, Trans. Nat. Hist. Soc. Formosa, 24: 492 (*Cenodocus*) (Hassenzan).

Distr.: Taiwan.

C. (*I.*) *formosana* (SCHWARZER, 1925) 蓬萊星斑天牛

SCHWARZER, 1925, Ent. Blaett., 21: 60 (*Mesosa*) (Hozan).

Distr.: Taiwan; E. China.

C. (*I.*) *watantakkuni* (Kano, 1933) 鬚角星斑天牛

KANO, 1933, Kontyû, 6: 280 (Nankotaizan).

Distr.: Taiwan.

Subgenus *Corethrophora* BLANCHARD, 1853

Corethrophora BLANCHARD, 1853, Voy. Pole Sud. Zool., 4: 301 (Type-species: *C. semiluctuosa* BLANCHARD, 1853).

C. (*Corethrophora*) *shirupiti* KANO, 1939 蘭嶼纓角星斑天牛

KANO, 1939, Annot. Zool. Jap., 18: 31, pl. 2, f. 4 (Lanyu I.)

Distr.: Taiwan (Lanyu I.).

Genus *Agelasta* NEWMAN, 1842 棋斑天牛屬

Agelasta NEWMAN, 1942, Entomol., 1: 269 (Type-species: *A. transversa* NEWMAN, 1842).

Subgenus *Dissosira* PASCOE, 1865

Dissosira PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 124, note (Type-species: *Agelasta catenata* PASCOE, 1862)

Agelasta (*Dissosira*) *cana* BREUNING, 1939 大坂棋斑天牛

BREUNING, 1939, Novit. Ent., 9, suppl. 3: 520 (Taihanroku).

Distr.: Taiwan.

A. (*D.*) *tonkinea* PIC, 1925

PIC, 1925, Bull. Soc. Ent. France, p. 188 (Tonkin).

formosana SCHWARZER, 1925, Ent. Blaett., 21: 61 (Fuhosho et Kosempo).

Distr.: Taiwan; W. China, Hainan I., Assam, Laos, Tonkin.

subsp. *tonkinea* PIC, 1925, s. str. 吹粉棋斑天牛

PIC, 1925, Bull. Soc. Ent. France, p. 188 (Tonkin).

Distr.: Taiwan; Assam, Laos, Tonkin.

Subgenus *Parasaimia* BREUNING, 1935

Parasaimia BREUNING, 1935, Fol. Zool. Hydrobiol., 8: 267 (Type-species: *Mimocoptops laosensis* PIC, 1925).

A. (*Parasaimia*) *hirticornis* (GRESSITT, 1936) 毛角星斑天牛

GRESSITT, 1936, Philip. J. Sci., 61: 102, pl. 1, f. 13 (*Saimia*?) (Hassenzan).

Distr.: Taiwan.

Genus *Mutatocoptops* PIC, 1923 小笠原星天牛屬

Mutatocoptops PIC, 1923, Mel. Exot.-Ent., 44: 29 (Type-species: *M. alboapicalis* PIC, 1925).

Saimia subgen. *Paracoptops* HELLER, 1926, Tijdschr. Ent., 69: 36 (Type-species:

Saimia (*Paracoptops*) *tuberosa* HELLER, 1926 = *Saimia bituberosa* PASCOE, 1866).

Mutatocoptops anancyloides (SCHWARZER, 1925) 灰紋星斑天牛

SCHWARZER, 1925, Ent. Blaett., 21: 61 (*Coptops*) (Hozan).

Distr.: Taiwan; Laos.

Tribe Homonoeini 淡色燦天牛族

Genus *Micromulciber* AURIVILLIUS, 1913 四條天牛屬

Micromulciber AURIVILLIUS, 1913, Ark. Zool., 8, 22: 25 (Types-species: *Mulciber biguttatus* PASCOE, 1867)

Kamikiria MATSUSHITA, 1933, J. Fac. Agric. Hokkaido Imp. Univ., 34(2): 348 (Xylorhizini) (Type-species: *Kamikiria plagiata* MATSUSHITA, 1933 = *Micromulciber 4-signatus* SCHWARZER, 1925).

Micromulciber quadrisignatus SCHWARZER, 1925 白星僞叉尾天牛

SCHWARZER, 1925, Ent. Blaett., 21: 64 (Sokutsu).

plagiata MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 349, pl. 4, f. 5 (Koshun, Kuraru).

Distr.: Taiwan; Japan (Ryukyu Is.).

Genus *Bumetopia* PASCOE, 1858 淡色燦天牛屬

Bumetopia PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 252 (Type-species: *B. oscitans* PASCOE, 1858).

Bumetopia oscitans PASCOE, 1858 蘭嶼燦天牛

PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 252, pl. 26, f. 7 (Hong Kong).

Distr.: Taiwan; Japan, S. China.

B. lanshuana HAYASHI, 1974 蘭嶼淡色燦天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 40 (Lanyu Is.).

Distr.: Taiwan (Lanyu I.).

B. lutoana HAYASHI, 1974 綠島淡色燦天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 41 (Lutau I.).

Distr.: Taiwan (Lutau I.).

B. stolata (MATSUSHITA, 1931) 蘭嶼超扁條天牛MATSUSHITA, 1931, Trans. Sapporo Nat. Hist. Soc., 12: 45, f. 2 (*Tmesisternus*)
(Lanyu I.).*tobacoximae* KANO, 1931, Bull. Biogeogr. Soc. Japan, 2: 185, pl. 16, f. 2 (*Homonoea*)
(Lanyu I.).

Distr.: Taiwan (Lanyu I., Lutau I.).

Tribe Apomecynini 白星銹天牛族

Genus *Apomecyna* LATREILLE, 1829 白星銹天牛屬*Apomecyna* LATREILLE, 1829, in Cuvier, Regne Anim., ed. 2, 5: 126 (Type-species:
Lamia alboguttata MEGERLE, 1802).*Apomecyna naevia* BATES, 1873 黃斑細銹天牛

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 317 (Japan).

Distr.: Taiwan(?); Japan.

A. historio (FABRICIUS, 1792) 胸紋銹天牛FABRICIUS, 1792, Ent. Syst., 1(2): 288 (*Lamia*) (India).*alboguttata* MEGERIE, 1802, Cat. Ins. Append. Nov., 473: 10 (*Saperda*).*quadrifasciata* THOMSON, 1868, Physis., 2: 59.*maculaticollis* PIC, 1918, Mel. Exot.-Ent., 28: 6.Distr.: Taiwan; Japan (Ryukyu Is.), Korea, Hainan I., Philippines, Indonesia,
India, Laos.*A. saltator* (FABRICIUS, 1781) 香瓜銹天牛FABRICIUS, 1781, Mant. Ins., 1: 141 (*Lamia*).*neglecta* PASCOE, 1865, Mant. Ins., 1: 152.*pertigera* THOMSON, 1868, Physis., 2: 160.*multinotata* PIC, 1918, Mel. Exot.-Ent., 28: 5.*cantator excavaticeps* GRESSITT, 1942, Philip. J. Sci., 72: 161, pl. 4, f. 9-10 (Hainan
I.).*excavaticeps*: GRESSITT, 1942, Lingn. Nat. Hist. Surv. Mus. Spec. Publ., 8: 25.

Distr.: SE Asia.

Genus *Ropica* PASCOE, 1858 雙星銹天牛屬*Ropica* PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 247 (Type-species: *R. piperata*
PASCOE, 1858).*Ropica dorsalis* SCHWARZER, 1925 雙星銹天牛SCHWARZER, 1925, Ent. Blaett., 21: 145 (*R. formosana* var. *dorsalis*) (Taihorin,
Sokutsu et Kankau).

Distr.: Taiwan; Hainan I., S. China, Hong Kong.

R. formosana BATES, 1866 臺灣雙星銹天牛

BATES, 1866, Proc. Ent. Soc. London, p. 351 (Taiwan).

Distr.: Taiwan.

R. fuscolaterimaculata HAYASHI, 1974 蘭嶼緣紋銹天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 42 (Lanyu I.).

Distr.: Taiwan. (Lanyu I.).

R. honesta PASCOE, 1865 蘭嶼雙星銹天牛

PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 190 (New Guinea, Moluccas).

formosana BATES, var. *dorsalis* SCHWARZER, 1925, Ent. Bl. Biol. Syst. Kafer, 21(4): 145 (Taihorin, Kankau).

dorsalis: GRESSITT, 1951, Longicornia, 2: 491.

rufescens PIC, 1926, Melang Exot.-Ent., 46: 5 (Tonkin).

burketi GRESSITT, 1937, Lingn. J. Sci., 16: 609 (SE China).

langana PIC, 1945, Echange, 61(499): 3 (Tonkin).

Distr.: Taiwan (Lanyu I.); Japan, New Guinea, SE Asia.

R. variabilis SCHWARZER, 1925 雙紋銹天牛

SCHWARZER, 1925, Ent. Blaett., 21: 68 (*R. formosana* var.).

Distr.: Taiwan.

Genus *Iproca* GRESSITT, 1940 尖翅銹天牛屬

Iproca GRESSITT, 1940, Philip. J. Sci., 72: 165 (Type-species: *I. acuminata* GRESSITT, 1940).

Iproca flavolineata HAYASHI, 1971 黃條尖翅銹天牛

HAYASHI, 1971, Ent. Rev. Japan, 23(1): 10 (Shihmen Reservoir).

Distr.: Taiwan.

Genus *Microlera* BATES, 1873 菱天牛屬

Microlera BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 380 (Type-species: *M. ptinoides* BATES, 1873).

Microlera kanoi HAYASHI, 1971 鹿野氏菱天牛

HAYASHI, 1971, Ent. Rev. Japan, 23(1): 9 (Jujiro).

Distr.: Taiwan.

Genus *Asaperda* BATES, 1873 華夏黑斑天牛屬

Asaperda BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 385 (Type-species: *A. rufipes* BATES, 1873).

Asaperda meridiana MATSUSHITA, 1931 卓社黑斑天牛

MATSUSHITA, 1931, Trans. Sapporo Nat. Hist. Soc., 12: 45 (Taihorin, Hozan et Hori).

takushaensis KANO, 1933, Kontyû, 6: 283 (Takusha).

Distr.: Taiwan; SE China.

Genus *Pseudokamikiria* TIPPMAN, 1942 假叉尾天牛屬

Pseudokamikiria TIPPMAN, 1942, Keleop. Rund., 33: 124.

Pseudokamikiria gressitti (TIPPMAN, 1942) 克雷氏假叉尾天牛

TIPPMAN, 1942, Koleop. Rund., 33: 124 (*Kamikiria*) (Musha).

Distr.: Taiwan.

Genus *Doliops* WATERHOUSE, 1841 假硬象天牛屬

Doliops WATERHOUSE, 1841, Proc. Ent. Soc. London, p. 278, (Type-species: *D. curculionoides* WATERHOUSE, 1841).

Doliops similis MIWA et MITONO, 1933 假硬象天牛

MIWA et MITONO, 1933, Trans. Nat. Hist. Soc. Formosa, 23: 373 (n. n. for *pachyrhynchoides* MIWA et MITONO, 1932).

pachyrrhynchoides MIWA et MITONO (nec HELLER), 1932, Trans. Nat. Hist. Soc. Formosa, 22: 306, f. (*Aprophata*), (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

Genus *Mimoplocia* BREUNING, 1949 黃紋細銹天牛屬

Mimoplocia BREUNING, 1949, Bull. Inst. Soc. Nat. Belg., 25, 38: 19 (Type-species: *Plocia notata* NEWMAN, 1842).

Mimoplocia notata (NEWMAN, 1842) 蘭嶼黃紋細銹天牛

NEWMAN, 1842, Entomologist, 1: 292 (*Plocia*) (Philippines).

albicornis HEYROVSKY, 1935, Cas. Cesk. Spol. Ent., 32: 19 (*Epaphra*).

Distr.: Taiwan (Lanyu I.); Philippines.

Genus *Atimura* PASCOE, 1863 瘤條銹天牛屬

Atimura PASCOE, 1863, Trans. Ent. Soc. London, (3)1: 548 (Type-species: *A. terminata* PASCOE, 1863).

Atimura formosana MATSUSHITA, 1933 蓬萊筒型矮天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 371 (Shijuwei).

Distr.: Taiwan.

A. fulva SCHWARZER, 1925 黃翅矮天牛

SCHWARZER, 1925, Ent. Blaett., 21: 145 (Kankau).

Distr.: Taiwan.

Genus *Xylariopsis* BATES, 1834 白領天牛屬

Xylariopsis BATES, 1884, J. Linn. Soc. London, 18: 247 (Type-species: *X. mimica* BATES, 1884).

Xylariopsis esakii MITONO, 1943 江崎氏白領天牛

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 33: 582 (Hassenzan: Reimei).

Distr.: Taiwan.

Genus *Sybra* PASCOE, 1865 豔紋矮天牛屬

Sybra PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 141, 198 (Type-species: *Ropica*

stigmatica PASCOE, 1859).

Sybra pascoei LAMEERE, 1893

LAMEERE, 1893, Annls. Soc. Ent. Fr., 62: 285 (Indo-China (Annan)).

Distr.: Taiwan; Japan (Ryukyu Is.), Hainan I., Vietnam.

subsp. *taiwanella* GRESSITT, 1951 巴斯氏矮天牛

GRESSITT, 1951, Longicornia, 2; 499 (Rokki).

Distr.: Taiwan.

S. ordinata BATES, 1873

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 318 (Japan).

Distr.: Taiwan; Japan, Hainan I., SE China.

subsp. *flavostriata* HAYASHI, 1968 黃條麗紋矮天牛

HAYASHI, 1968, Ent. Rev. Japan, 21(1): 16 (Japan (Ryukyu Is.)).

alternatus: MIWA, 1933, Trans. Nat. Hist. Soc. Formosa, 23(124): 13 (Japan (Ryukyu Is.)).

Distr.: Taiwan; Japan (Ryukyu Is. (Yaeyama I.)).

S. baculina BATES, 1866 後紋矮天牛

BATES, 1866, Proc. Zool. Soc. London, p. 352 (Taiwan).

Distr.: Taiwan; Japan (Ryukyu Is.).

S. maculiclunis MATSUSHITA, 1931 凤山矮天牛

MATSUSHITA, 1931, Mitt. Zool. Mus. Berl., 17: 404 (Hozan, Taihorin, Alikang).

Distr.: Taiwan.

S. kotoensis MATSUSHITA, 1934 蘭嶼矮天牛

MATSUSHITA, 1934, Trans. Nat. Hist. Soc. Formosa, 23(135): 541 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

S. miscanthivola MAKIHARA, 1979 盧葦矮天牛

MAKIHARA, 1979, Ent. Rev. Japan, 30(1/2): 25 (Japan (Yonaguni I.)).

Distr.: Taiwan; Japan (Yonaguni I.).

S. narai HAYASHI, 1976 南山矮天牛

HAYASHI, 1976, Bull. Osaka Jonan Women's Jr. Coll., 11: 14 (Nanshanchi).

Distr.: Taiwan.

S. punctatostriata BATES, 1866 棉天牛

BATES, 1866, Proc. Zool. Soc. London, p. 351 (Taiwan).

gossypi MATSUMURA, 1912, Spec. Bull. Agr. Exp. Sta. Formosa, 5: 219, pl. 11, f. 1-4.

Distr.: Taiwan; Hainan I., SE China.

S. bioculata PIC, 1925 眼紋矮天牛

PIC, 1925, Bull. Soc. Ent. France, p. 302 (China).

Distr.: Taiwan; E China.

subsp. *quadrinotata* SCHWARZER, 1925 角紋矮天牛

SCHWARZER, 1925, Ent. Blaett., 21: 145 (Kankau).

Distr.: Taiwan.

S. botelensis BREUNING et OHBAYASHI, 1966 耶美矮天牛

BREUNING et OHBAYASHI, 1966, Bull. Jap. Ent. Acad., 2(3): 13 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

S. albomaculata BREUNING, 1939

BREUNING, 1939, Festschr. E. Strand, 5: 264 (N. China).

Distr.: Taiwan; N China.

subsp. formosana BREUNING, 1964 白紋矮天牛

BREUNING, 1964, Ent. Abh. Mus. Tierk. Dresden, 30: 305.

Distr.: Taiwan.

Genus *Sybrodiboma* BREUNING, 1853 臺灣矮天牛屬

Sybrodiboma BREUNING, 1853, Bull. Inst. R. Sci. Nat. Belg., 29(8): 16 (Type-species: *Sybra subfasciata* BATES, 1884).

Sybrodiboma taiwanensis HAYASHI, 1974 臺灣矮天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 51 (Lishan).

Distr.: Taiwan.

Genus *Palausybra* GRESSITT, 1956 欠翅矮天牛屬

Palausybra GRESSITT, 1956, Insects Micronesia, 17(3): 153 (Type-species: *P. vestigialis* GRESSITT, 1956).

Palausybra chibi HAYASHI, 1974 特微天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 41 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

Genus *Paraneosybra* HAYASHI, 1978 黃條矮天牛屬

Paraneosybra HAYASHI, 1978, Ent. Rev. Japan, 32(1/2): 107 (Type-species: *P. fulvofasciata* HAYASHI, (1978)).

Paraneosybra fulvofasciata HAYASHI, 1978 南山黃條矮天牛

HAYASHI, 1978, Ent. Rev. Japan, 32(1/2): 108 (Nanshanchi).

Distr.: Taiwan.

Genus *Brachysybra* BREUNING 1940 短圓矮天牛屬

Brachysybra BREUNING, 1940, Folia Zool. Hydrob., 10: 127 (Type-species: *B. elliptica* BREUNING, 1940).

Brachysybra elliptica BREUNING, 1940 長圓矮天牛

BREUNING, 1940, Folia Zool. Hydrob., 10: 428 (Philippines).

Distr.: Taiwan (Lanyu I.); Philippines.

Genus *Neosybra* BREUNING, 1939 姫艷紋矮天牛屬

Neosybra BREUNING, 1939, Festchr. E. Strand, 5: 276 (Type-species: *N. ropicoides* BREUNING, 1939).

Cornalis: MATSUSHITA, (nec THOMSON, 1864) 1933, J. Fac. Agric. Hokkaido Imp. Univ., 34(2): 370.

Neosybra costata (MATSUSHITA, 1933) 隆條矮天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 370 (*Cornalis*) (Toroen).

Distr.: Taiwan.

N. sinuicosta GRESSITT, 1951 波條矮天牛

GRESSITT, 1951, Longicornia, 2: 502, pl. 20, f. 2-4 (Shinhassenzan).

Distr.: Taiwan.

Genus *Rondonia* BREUNING, 1962 雙紋矮天牛屬

Ronconia BREUNING, 1962, Bull. Soc. R. Sci. Nat. Laos, 3: 7 (Type-species: *R. ropicoides* BREUNING, 1962).

Rondonia bisignata HAYASHI, 1976 雙紋矮天牛

HAYASHI, 1976, Bull. Osaka Jonan Women's Jr. Coll., 11: 16 (Kenting Park).

Distr.: Taiwan.

Tribe Hippopsini 瘦天牛族

Genus *Aulaconotus* THOMSON, 1864 滿翅天牛屬

Aulaconotus THOMSON, 1864, Syst. Ceramb., p. 99 (Type-species: *A. pachypezoides* THOMSON, 1864).

A. gracilicornis MAKIHARA et SAITO, 1985 繖角溝翅天牛

MAKIHARA et SAITO, 1985, Mis. Rep. Hiwa Mus. Nat. Hist., 23: 14 (Kenting Park).

pachypezoides: TAKASU, 1981 (nec THOMSON), Sayabane, 7: 17 (Wulai) (In Japanese).

Distr.: Taiwan.

Genus *Pothyne* THOMSON, 1864 白條瘦天牛屬

Pothyne THOMSON, 1864, Syst. Ceramb., p. 97 (Type-species: *P. variegata* THOMSON, 1864).

Neopothyne MATSUSHITA, 1931, Trans. Sapporo Nat. Hist. Soc., 12: 46 (Type-species: *N. variegata* MATSUSHITA, 1931 = *Pothyne variegata* THOMSON, 1864).

Pothyne albolineata MATSUSHITA, 1933 白條瘦天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 383 (Taihorin).

Distr.: Taiwan.

P. formosana SCHWARZER, 1925

SCHWARZER, 1925, Ent. Blaett., 21: 146 (Kosempo, Kankau et Sokutsu).

Distr.: Taiwan; Japan (Ryukyu Is.).

subsp. *formosana* SCHWARZER, 1925, s. str. 臺灣沉香瘦天牛

SCHWARZER, 1925, Ent. Blaett., 21: 146 (Kosempo, Kankau et Sokutsu).

Distr.: S. Taiwan.

subsp. *nanshanchina* TAKAKUWA et KUSAMA, 1979 南山沉香瘦天牛

TAKAKUWA et KUSAMA, 1979, Elytra, 7(1):10 (Nanshanchi, Lienhwachi, Hui-Sun Exptl. Forest, Palin, Yangmingshan).

Distr.: Central & North Taiwan.

P. lanhsuensis HAYASHI, 1978 蘭嶼瘦天牛

HAYASHI, 1978, Ent. Rev. Japan, 32(1/2): 108 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

P. semiaulaconotus HAYASHI, 1974 顆胸瘦天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 43 (Taiwan).

Distr.: Taiwan.

P. variegata THOMSON, 1864 灰紋瘦天牛

THOMSON, 1864, Syst. Ceramb., p. 97 (Malaysia).

Distr.: Taiwan; Japan (Ryukyu Is.), Siam, Burma, Assam.

P. virginialis TAKAKUWA et KUSAMA, 1979 墾丁瘦天牛

TAKAKUWA et KUWAMA, 1979, Elytra, 7(1): 4 (Kenting Park).

Distr.: Taiwan.

P. silacea PASCOE, 1871 琉球白條瘦天牛

PASCOE, 1871, Ann. Mag. Nat. Hist., (4)8: 278 (Japan).

imasakai HAYASHI, 1976, Bull. Osaka Jonan Women's Jr. Coll., 11: 18 (Japan (Yonaguni I.)).

Distr.: Taiwan; Japan, China, Hainan I.

P. annulata BREUNING, 1942 縱條瘦天牛

BREUNING, 1942, Folia Zool. Hydrobiol., 6: 167 (Japan).

silacea MATSUSHITA (nec PASCOE), 1933, J. Fac. Agric. Hokkaido Imp. Univ., 34; 384.

silacea KOJIMA et HAYASHI (nec PASCOE), 1969, Insects' Life in Japan, 1: 106, pl. 32, f. 4; KUSAMA (nec PASCOE), 1973, List Ecology & Dist. Jap. Ceramb., p. 94.

Distr.: Taiwan(?); Japan, Hainan I.

Genus Illinoia DISCOON 1964 L. M. M.

P. piceus GRESSITT, 1951 黑身瘦天牛

GRESSITT, 1951, Longicornia, 2: 543 (China).

Distr.: Taiwan; China.

P. leptissimus GRESSITT, 1936 阿里山瘦天牛

GRESSITT, 1936, Philip. J. Sci., 61: 103, pl. 1, f. 14 (Arisan, Taiheizan).

Distr.: Taiwan; SE. China.

Genus *Metopoplectus* GRESSITT, 1937 紅長角天牛屬

Metopoplectus GRESSITT, 1937, Philip. J. Sci., 61: 104 (Type-species: *M. taiwanensis* GRESSITT, 1937).

Metopoplectus taiwanensis GRESSITT, 1936 姬棗紅長角天牛

GRESSITT, 1936, Philip. J. Sci., 61: 105, pl. 1, f. 15 (Hori).

Distr.: Taiwan.

Genus *Cleptometopus* THOMSON, 1864 斜帶長角天牛屬

Cleptometopus THOMSON, 1864, Syst. Ceramb., pp. 95, 389 (Type-species: *C. terrestris* THOMSON, 1864).

Apophrena PASCOE, 1866, Trans. Ent. Soc. London, (3)3: 323, 324 (Type-species: *A. filifera* PASCOE, 1866).

Smermus LACORDAIRE, 1869, Hist. Nat. Ins., Gen. Col., 9: 692, 696 (Type-species: *S. mniszechi* LACORDAIRE, 1869).

Acroama JORDAN, 1894, Novit. Zool., 1: 501 (Type-species: *A. armata* JORDAN, 1894).

Anapophrena PIC, 1925, Melang Exot.-Ent., 43: 28 (Type-species: *A. lutenotata* PIC, 1925).

Mimocleptometopus PIC, 1934, Mat. Long., 11(2): 36 (Type-species: *M. undulatus* PIC, 1934).

Metopoplectus GRESSITT, 1936, Philip. J. Sci., 61: 104 (Type-species: *M. taiwanensis* GRESSITT, 1936).

Itohigea MATSUSHITA, 1938, Insecta Mats., 12(2/3): 102 (Xylorhizini) (Type-species: *I. bimaculata* MATSUSHITA, 1938).

Cleptometopus orientalis MITONO, 1934 棗紅長角天牛

MITONO, 1934, Trans. Nat. Hist. Soc. Formosa, 24: 492 (Hori, Karenko, Kahodai).

Distr.: Taiwan.

Tribe Pteropliini 鎏天牛族

Genus *Abryna* NEWMAN, 1842 鎏麗天牛屬

Abryna NEWMAN, 1842, Entomologist, 1: 289 (Type-species: *A. coenosa* NEWMAN, 1842).

Abryna coenosa NEWMAN, 1842

NEWMAN, 1842, Entomologist, 1: 289 (Philippines).

?*subuniformis* PIC, 1925, Melang Exot.-Ent., 43: 30.

Distr.: Taiwan; Japan (Ryukyu Is.), S. China, Philippines.

subsp. obscura SCHWARZER, 1925 鎏麗天牛
 SCHWARZER, 1925, Ent. Blaett., 21: 65 (Taihorin).
superba PIC, 1932, Echange, 48: 24 (Taiwan).
 Distr.: Taiwan.

Genus *Niphona* MULSANT, 1839 灰叉尾天牛屬

Niphona MULSANT, 1839, Col. France Long., 1: 369 (Type-species: *N. picticornis* MULSANT, 1839).
Camptocnema THOMSON, 1864, Syst. Ceramb., pp. 54, 367 (Type-species: *N. lateralis* WHITE, 1858).
Ocheutes THOMSON, 1864, Syst. Ceramb., pp. 54, 368 (Type-species: *N. scopulifera* DEJ.).
Aelara THOMSON, 1864, Syst. Ceramb., pp. 55, 368 (Type-species: *N. regis-fernandi* PAIVA, 1860).

Niphona furcata (BATES, 1873) 灰身叉尾天牛

BATES, 1873, Ann. Mag. Nat. Hist., (4)7: 314 (*Aelara*) (Japan).
 Distr.: Taiwan; Japan, China.

N. parallela (WHITE, 1858) 印度叉尾天牛

WHITE, 1858, Ann. Mag. Nat. Hist., (3)2: 267 (*Aelara*) (N. India).
minor LAMEERE, 1893, Ann. Soc. Ent. France, 62: 284 (*Aelara*) (Cambodia).
 Distr.: Taiwan; China, Burma, N. India, Cambodia, Hainan.

N. yanoi MATSUSHITA, 1934

MATSUSHITA, 1934, Trans. Nat. Hist. Soc. Formosa, 24: 240 (Kuraru).
 Distr.: Taiwan; Japan (Ryukyu Is.), Hainan I.

subsp. yanoi MATSUSHITA 1934, s. str. 矢野氏叉尾天牛

MATSUSHITA, 1934. Trans. Nat. Hist. Soc. Formosa, 24: 240 (Kuraru).
 Distr.: Taiwan; Japan.

Genus *Nipholophia* GRESSITT, 1951 擬叉尾天牛屬

Nipholophia GRESSITT, 1951, Longicornia, 2: 458 (Type-species: *N. chujoi*, GRESSITT, 1951).

Nipholophia chujoi GRESSITT, 1951 中條氏擬叉尾天牛

GRESSITT, 1951, Longicornia, 2: 458 (Musha).

Distr.: Taiwan.

Genus *Falsoterinaea* MATSUSHITA, 1938 平山天牛屬

Falsoterinaea MATSUSHITA, 1938, Kontyû, 12: 95 (n. n. for *Hirayamaia* MATSUSHITA, 1937).

Falsoterinaea fuscorufa (MATSUSHITA, 1937) 褐紅平山天牛

MATSUSHITA, 1937, Kontyû, 11: 103 (*Hirayamaia*) (Musha).
 Distr.: Taiwan.

Genus *Pterolophia* NEWMAN, 1842 鎏天牛屬

Pterolophia NEWMAN, 1842, Entomologist, 1: 370 (Type-species: *Mesosa?* *bigibbera* NEWMAN, 1842).

Subgenus *Pterolophia* NEWMAN, 1842, s. str.

Albana MULSANT, 1846, Col. Fr. Long. Suppl., p. 13 (Type-species: *A. m-griseum* MULSANT, 1846).

Prioneta BLANCHARD, 1853, in d'Urville, Voy. Pole Sud., Zool., 4: 292, pl. 17, f. 11 (Type-species: *P. albosignata* BLANCHARD, 1853).

Praonetha PASCOE, 1862, J. Ent., 1: 348 (n. n. pro. *Prioneta* BLANCHARD, 1853).

Theticus THOMSON, 1858, Arc. Nat., 2: 190 (Type-species: *T. biarcuatus* THOMSON, 1858).

Alyattes THOMSON, 1864, Syst. Ceramb., p. 48 (Type-species: *A. guennensis* THOMSON, 1864).

Prionetopsis THOMSON, 1864, Syst. Ceramb., p. 49 (Type-species: *P. batteata* THOMSON, 1864=*Lamia inaequalis* FABRICIUS, 1801).

Cormia PASCOE, 1864, J. Ent., 2: 281 (Type-species: *C. ingrata* PASCOE, 1864).

Anaches PASCOE, 1865, Trans. Ent. Soc. London, 3(3): 160 (Type-species: *Sthenias dorsalis* PASCOE, 1858).

Eurycotyle BLESSIG, 1873, Trudy Russk. Ent. Obschch., 9: 210 (Type-species: *E. maacki* BLESSIG, 1873).

Acroptycha QUEDENFELDT, 1888, Berl. Ent. Z., 32: 209 (Type-species: *A. spinifera* QUEDENFELDT, 1888).

Incamelomorpha PIC, 1926, Melang. Exot-Ent., 46: 4 (Type-species: *Pterolophia (Incamelomorpha) depensis* PIC, 1926) (subg.).

Paralophira AURIVILLIUS, 1924, Zool. Meded., 8: 30 (Type-species: *P. quadrinodosa* AURIVILLIUS, 1924).

Pterolophia (*P.*) *angusta* (BATES, 1873) 圓紋鎔天牛

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 317 (*Praonetha*) (Japan).

Distr.: Taiwan(?); Japan, NE. China.

P. (*P.*) *bigibbera* (NEWMAN, 1842) 雙瘤鎔天牛

NEWMAN, 1842, Entomologist, 1: 323 (*Mesosa?*) (Philippines).

camura NEWMAN, 1842, Entomologist, 1: 371 (Philippines).

binodosa BATES, 1866, Proc. Zool. Soc. London, p. 350 (*Praonetha*) (Taiwan).

saipana OHBAYASHI, 1941, Tenthredo, 3(3): 225 (Saipan Is.).

Distr.: Taiwan; Japan (Ogasawara Is.), Philippines, Mariana.

P. (*P.*) *kaleea* (BATES, 1866)

BATES, 1866, Proc. Zool. Soc. London, p. 351 (*Praonetha*) (Taiwan).

subleliopodina BREUNING et OHBAYASHI, 1964, Bull. Japan. Ent. Acad., 1(4): 16 (Japan (Ryukyu Is.)).

subexternefasciata MITONO, 1941, Cat. Col. Japan., (8): 172 (nec PIC, 1926).

matsushitai KUSAMA, 1965, New Ins. Collect., (Ge), p. 470 (Japan (Ryukyu Is.)) (nec BREUNING, 1937).

Distr.: Taiwan; Japan (Ryukyu Is.), China.

subsp. *kaleea* (BATES, 1866) s. str. 姬後紋锈天牛

BATES, 1866, Proc. Zool. Soc. London, p. 351 (*Praonetha*) (Taiwan).

Distr.: Taiwan; Japan (Ryukyu Is.).

subsp. *inflexa* GRESSITT, 1940 華南姬後紋锈天牛

GRESSITT, 1940, Lingnan Soc. J., 19: 11, pl. 1, f. 3 (China).

Distr.: Taiwan; S. China.

P. (P.) matsushitai BREUNING, 1938 松下氏锈天牛

BREUNING, 1938, Festchr. E. Strand, 4: 289 (Kuraru).

Distr.: Taiwan.

P. (P.) obscura SCHWARZER, 1925

SCHWARZER, 1925, Ent. Blaett., 21: 66 (Taiwan).

Dirst.: Taiwan.

subsp. *obscura* SCHWARZER, 1925 藤花锈天牛

SCHWARZER, 1925, Ent. Blaett., 21: 66 (Taiwan).

Distr.: Taiwan.

subsp. *postmaculatoides* BREUNING, 1968 曇紋锈天牛

BREUNING, 1968, Bull. Soc. Ent. Mulhouse, p. 57 (n. n.)

obscura postmaculata BREUNING et OHBAYASHI, 1966, Bull. Jap. Ent. Acad., 2(3): 13.

Distr.: Taiwan.

P. (P.) reduplicata GRESSITT, 1951 八仙锈天牛

GRESSITT, 1951, Longicornia, 2: 471, pl. 18, f. 9 (Hassenzan)

Distr.: Taiwan.

P. (P.) zonata (BATES, 1873) 白尾锈天牛

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 315 (*Praonetha*) (Japan).

Distr.: Taiwan(?); Japan.

P. (P.) laterialba (SCHWARZER, 1925) 白緣锈天牛

SCHWARZER, 1925, Ent. Blaett., 21: 68 (*Albana*) (Anping).

Distr.: Taiwan.

P. (P.) variantennalis BREUNING, 1970 大埔林锈天牛

BREUNING, 1970, Koleop. Rdsch., 48: 10 (Taihorinsho).

Distr.: Taiwan.

Subgenus *Hylobrotus* LACORDAIRE, 1869

Hylobrotus LACORDAIRE, 1869, Hist. Nat. Ins., Gen. Col., 9: 522, 538 (Type-species:

H. ploemi LACORDAIRE, 1872).

P. (H.) annulata (CHEVROLAT, 1845) 輪紋锈天牛

CHEVROLAT, 1845, Rev. Zool., 8: 99 (*Coptops*) (Macao).

bowringii PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 170 (*Praonetha*) (Hong Kong).

scutellata SCHWARZER, 1925, Ent. Blaett., 21: 66 (Kosempo, Alikang, Kankau).

annulicornis PIC, 1925, Bull. Soc. Ent. Fr., p. 138 (China).

lacosa PIC, 1926, Melang. Exot.-Ent., 46: 3 (Tonkin).

lunigera AURIVILLIUS subsp. *formosana* HAYASHI, 1960, Ent. Rev. Japan, 11(1):

28 (Japan (Kita-Ogari-jima Is.)); SAMUELSON, 1965, Pacific Insects, 7(1): 108 (part: C. Ryukyu). (nec SCHWARZER, 1925).

Distr.: Taiwan; Japan, S. China, Hong Kong, Macao, Hainan I., Korea, Burma.

P. (H.) latefascia SCHWARZER, 1925 橫帶銹天牛

SCHWARZER, 1925, Ent. Blaett., 21: 65 (Kosempo et Kankau).

fainanensis PIC, 1926, Melang Exot.-Ent., 45: 32 (Fainan).

yayeyamai BREUNING, 1961, Bull. Inst. R. Sci. Nat. Belg., 37(20): 9 (Japan (Ryukyu Is.)).

horii BREUNING et OHBAYASHI, 1967, Bull. Japan. Ent. Acad., 3(2): 3 (Japan (Ryukyu Is.)).

Distr.: Taiwan; Japan (Ryukyu Is.).

P. (H.) formosana SCHWARZER, 1925 虹紋銹天牛

SCHWARZER, 1925, Ent. Bl. Biol. Syst. Kafer, 21(2): 66 (*P. lunigera* var.) (Kosempo).

lateralis formosana KUSAMA, 1973, List Ecol. Dist. Japan. Ceramb., p. 99.

Distr.: Taiwan; Japan (Okinawa I.).

Subgenus *Ale* PASCOE, 1865

Ale PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 96, 131 (Type-species: *A. agraria* PASCOE, 1865).

Achriotypa PASCOE, 1875, Ann. Mag. Nat. Hist., (4)15: 71 (Type-species: *A. basalis* PASCOE, 1875).

P. (Ale) gibbosipennis PIC, 1926

PIC, 1926, Melang Exot.-Ent., 45: 32 (Taiwan).

cristipennis MATSUSHITA, 1931, Mitt. Zool. Mus. Berlin, 17: 403 (Hori).

Distr.: Taiwan; Japan (Ryukyu Is.).

subsp. *gibbosipennis* PIC, 1926, s. str. 瘤翅銹天牛

PIC, 1926, Melang Exot.-Ent., 45: 32 (Taiwan).

Distr.: Taiwan.

Subgenus *Ovalopraonetha* BREUNING, 1961

Ovalopraonetha BREUNING, 1961, Cat. Lam. Monde, 4: 259 (Type-species: *Pterolophia (O.) ovalis* BREUNING, 1938) (subg.).

P. (Ovalopraonetha) kanoi HAYASHI, 1966 鹿野氏銹天牛

HAYASHI, 1966, Bull. Osaka Jonan Women's Jr. Coll., 1: 7 (Rodofu).

Distr.: Taiwan.

Subgenus *Pseudale*, SCHWARZER, 1925

Pseudale SCHWARZER, 1925, Ent. Blaett., 21: 67 (Type-species: *P. fasciata* SCHWARZER, 1925).

P. (Pseudale) dorsotuberculare (HAYASHI, 1984) 糜背銹天牛

HAYASHI, 1984, Bull. Osaka Jonan Women's Jr. Coll., 17/18: 33, pl. 2, f. 8 (*Pseudale?*) (Sungkang).

P. (P.) fasciata SCHWARZER, 1925 昇條銹天牛

SCHWARZER, 1925, Ent. Blaett., 21: 67 (*Pseudale*) (Fuhosho).

camura OHBAYASHI (nec NEWMAN), 1963, Icon. Ins. Japon. Col., 2: 309, pl. 155, f. 10.

Distr.: Taiwan.

P. (P.) suginoi MAKIHARA, 1986 杉野氏銹天牛

MAKIHARA, 1986, Mis. Rep. Hiwa Mus. Nat. Hist., 24: 9 (Hungyeh Wenchuan).

Distr.: Taiwan.

Subgenus *Lychrosis* PASCOE, 1866

Lychrosis PASCOE, 1866, J. Linn. Soc. London, 9: 89 (Type-species: *Hathlia zebrinus* PASCOE, 1857).

P. (Lychrosis) zebrinus (PASCOE, 1857) 白斑銹天牛

PASCOE, 1857, Trans. Ent. Soc. London, (2)4: 252 (*Hathlia*) (N. India).

Distr.: Taiwan; N. India, Tonkin, Hainan.

Genus *Marmylaris* PASCOE, 1866 巴克氏銹天牛屬

Marmylaris PASCOE, 1866, J. Linn. Soc. London, 9: 88 (Type-species: *Hathlia buckleyi* PASCOE, 1857).

Marmylaris buckleyi (PASCOE, 1857) 巴克氏銹天牛

PASCOE, 1857, Trans. Ent. Soc. London, (2)4: 107 (*Hathlia*) (N. India).

Distr.: Taiwan(?); N. India, Himalaya.

Genus *Desisa* PASCOE, 1865 霜斑銹天牛屬

Desisa PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 163 (Type-species: *Praonetha subfasciata* PASCOE, 1862).

Mesopenthea SCHWARZER, 1925, Ent. Blaett., 21: 67 (Type-species: *M. variabilis* SCHWARZER, 1925).

Desisa kuraruana (MATSUSHITA, 1935) 霜斑銹天牛

MATSUSHITA, 1935, Trans. Nat. Hist. Soc. Formosa, 25: 312 (*Mesosella*) (Kuraru).

Distr.: Taiwan.

D. takasagona MATSUSHITA, 1933 蓬萊白條銹天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 364, pl. 5, f. 9 (Horisha).

Distr.: Taiwan.

D. variabilis (SCHWARZER, 1925) 斑紋銹天牛

SCHWARZER, 1925, Ent. Blaett., 21: 67 (*Mesopenthea*) (Sokutsu).

Distr.: Taiwan.

Genus *Egesina* PASCOE, 1864 新島氏矮天牛屬

Egesina PASCOE, 1864, Trans. Ent. Soc. London, (3)3: 29, 49 (Type-species: *E. rigida* PASCOE, 1864).

Cuphisia PASCOE, 1866, Proc. Zool. Soc. London, p. 229 (Type-species: *C. callosa* PASCOE, 1866).

Callienispia FISHER, 1925, Philip. J. Sci., 28: 210 (Type-species: *C. elegans* FISHER, 1925).

Platyzeargyra FISHER, 1925, Philip. J. Sci., 28: 213 (Type-species: *P. bakeri* FISHER, 1925).

Neoegesina FISHER, 1925, Philip. J. Sci., 28: 215 (Type-species: *N. ornata* FISHER, 1925).

Subgenus *Egesina* PASCOE, 1864

Egesina PASCOE, 1864, Trans. Ent. Soc. London, (3)3: 29, 49 (Type-species: *E. rigida* PASCOE, 1864).

Egesina (Egesina) albofasciata HAYASHI, 1974 白點矮天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 46 (Lushan or Hotso).

Distr.: Taiwan.

Subgenus *Niijimaia* MATSUSHITA, 1933

Niijimaia MATSUSHITA, 1933, J. Fac. Agric. Hokkaido Univ., 34(2): 386 (Type-species: *Niijimaia bifasciana* MATSUSHITA, 1933).

Egesina subgen. *Pseudenispia* BREUNING, 1938, Festschr. 60 Geburtst E. Strand., 4: 388 (Type-species: *Pseudenispia albomarmorata* BREUNING, 1938).

Egesina (Niijimaia) fascolaterimaculata HAYASHI, 1971 側紋新島天牛

HAYASHI, 1971, Ent. Rev. Japan, 23(1): 13 (Tienhsiang).

Distr.: Taiwan.

E. (N.) formosana (SCHWARZER, 1925) 蓬萊新島天牛

SCHWARZER, 1925, Ent. Blaett., 21: 68 (*Enispia*) (Kosempo).

Distr.: Taiwan.

Genus *Sthenias* CASTELNAU, 1840 長角銹天牛屬

Sthenias CASTELNAU, 1840, Hist. Nat. Col., 2: 466 (Type-species: *Lamia grisor* FABRICIUS, 1887).

Chalarus FAHRAEUS, 1872, Ofvers. Vet. Akad. Forhl., 29, 2: 45 (Type-species: *C. leucaspis* FAHRAEUS, 1872=*cylindrator* FABRICIUS, 1801).

Anomamomus FAHRAEUS, 1872, Col. Hefte, 10: 194 (n. n. for *Chalarus*).

Thysanodes NEWMAN, 1842, Entomologist, 1: 292.

Albosthenias BREUNING, 1961, Bull. Mus. Hist. Nat. Ser. 2, 32: 538 (Type-species: *A. albicollis* GAHAN, 1890).

Sthenias cylindricus GRESSITT, 1939 白紋長角銹天牛

GRESSITT, 1939, Notes d'Ent. Chinoise, 6: 114, pl. 3, f. 7 (China).

Distr.: Taiwan; E. China.

S. semicylindricus HAYASHI, 1974 大白紋長角銹天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 45 (Juisui).

Distr.: Taiwan.

Genus *Similosodus* McKEOWN, 1945 裸胸天牛屬

Similosodus McKEOWN, 1945, Rec. Austral. Mus., 21: 292 (n. n. for *Sodus* PASCOE, 1864).

Sodus PASCOE: (nec AMYOT), 1864, Trans. Ent. Soc. London, (3)3: 137 (Type-species: *S. verticalis* PASCOE, 1864).

Similosodus taiwanus HAYASHI, 1966 黃條裸胸天牛

HAYASHI, 1966, Bull. Osaka Jonan Women's Jr. Coll., 1: 8 (Paran).

Distr.: Taiwan.

Tribe Morimopsini 赤銅天牛族**Genus *Dolophrades* BATES, 1884 泥色粗角天牛屬**

Dolophrades BATES, 1884, J. Linn. Soc. London, 18: 237 (Type-species: *D. terrenus* BATES, 1884).

Corcadiohammus PIC, 1934, Mel. Exot.-Ent., 63: 8 (Type-species: *D. punctatus* PIC, 1934).

Dolophrades punctatus (PIC, 1934) 假瘤粗角天牛

PIC, 1934, Melang Exot.-Ent., 63: 8 (Taiwan).

Distr.: Taiwan.

Tribe Phrissomini 長角瘤叉尾天牛族**Genus *Mesechthistatus* BREUNING, 1950 瘤叉尾天牛屬**

Mesechthistatus BREUNING, 1950, Longicornia, 1: 222 (Type-species: *Echthistatus binodosus* WATERHOUSE, 1881).

Mesechthistatus yamahoi (MITONO, 1943) 山保氏瘤叉尾天牛

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 33: 578, f. 1 (*Echthistatus*) (Taito: Aroe).

Distr.: Taiwan.

Tribe Agnini 長角天牛族**Genus *Aethalodes* GAHAN, 1888 瘤條天牛屬**

Aethalodes GAHAN, 1888, Ann. Mag. Nat. Hist., (6)1: 270 (Type-species: *A. verrucosus* GAHAN, 1888)

Aethalodes verrucosus GAHAN, 1888

GAHAN, 1888, Ann. Mag. Nat. Hist., (6)1: 270, pl. 16, f. 1 (N. China).

armata NONFRIED, 1892, Ent. Nachr., 18: 93 (*Trachystola*) (China).

nodicollis FAIRMAIRE, 1900, Ann. Soc. Ent. France, 68: 640 (China).

Distr.: Taiwan; China, Tonkin.

subsp. *formosanus* KRIESCHE, 1924 蓬萊瘤條天牛

KRIESCHE, 1924, Deut. Ent. Zeit., p. 285 (*A. formosanus*) (Kosempo et Taihorin).

Distr.: Taiwan.

Genus *Aristobia* THOMSON, 1868 細角長天牛屬

Aristobia THOMSON, 1868, Physis, 2: 178 (Type-species: *Cerosterna hispida* SAUNDERS, 1863).

Aristobia hispida (SAUNDERS, 1863) 細角長天牛

SAUNDERS, 1863, Trans. Ent. Soc. London, (2)2: 112, pl. 4, f. 6 (N. China).

Distr.: Taiwan; China, Hainan I., Tonkin.

Genus *Psacothea* GAHAN, 1888 黃星天牛屬

Psacothea GAHAN, 1888, Ann. Mag. Nat. Hist., (6)2: 400 (Type-species: *Monohammus hilaris* PASCOE, 1857).

Psacothea hilaris (PASCOE, 1857)

PASCOE, 1857, Trans. Ent. Soc. Lond., (2)4: 193 (*Monohammus*) (N. China).

Distr.: Taiwan; Japan, Korea, China.

subsp. *hilaris* (PASCOE, 1857) s. str. 黃星長角天牛

PASCOE, 1857, Trans. Ent. Soc. Lond., (2)4: 193 (*Monohammus*) (N. China).

Distr.: Taiwan; Japan, Korea, China.

subsp. *botelensis* BREUNING et OHBAYASHI, 1966 蘭嶼黃星天牛

BREUNING et OHBAYASHI, 1966, Bull. Japan Ent. Acad., 2(6): 12

Distr.: Taiwan (Lanyu I.).

Genus *Epepeotes* PASCOE, 1866 縱紋長角天牛屬

Epepeotes PASCOE, 1866, Proc. Zool. Soc. London, p. 249 (Type-species: *Lamia lusca* FABRICIUS, 1866).

Epepeotes ambigenus CHEVROLAT, 1841

CHEVROLAT, 1841, Rev. Zool., 4: 228

ambiguus DILLON et DILLON, 1941, Read. Publ. Mus., Soc. Publ., 1: 43, pl. 3, f. 23.

Distr.: Taiwan; Philippines.

subsp. *formosanus* BREUNING, 1943 蘭嶼縱紋長角天牛

BREUNING, 1943, Novit. Ent., 13, suppl. 3: 228 (Lanyu I.).

Distr.: Taiwan (Lanyu I., Lutau I.).

Genus *Paraleprodera* BREUNING, 1935 紗紅長角天牛屬

Paraleprodera BREUNING, 1934, Fol. Zool. hydrobiol., 8: 253 (Type-species: *Lamia crucifera* FABRICIUS, 1792).

Paraleprodera diophthalma (PASCOE, 1856)

PASCOE, 1956, Trans. Ent. Soc. London, (2)4: 49 (*Morimus*) (N. China).

bioculata FAIRMAIRE, 1900, Ann. Soc. Ent. France, 68: 641 (*Leprodera*) (China).

Distr.: Taiwan; E. et W. China.

subsp. *formosana* (SCHWARZER, 1925) 雙爪長角天牛

SCHWARZER, 1925, Ent. Blaett., 21: 58 (*Leprodera bioculata* subsp.) (Kosempo).

Distr.: Taiwan.

P. itzingeri BREUNING, 1935 四星繖紅長角天牛

BREUNING, 1935, Fol. Zool. Hydrobiol., 8: 254 (Taiwan).

carolina: SCHWARZER (nec FAIRMAIRE), 1925, Ent. Blaett., 21: 58 (Taiwan).

Distr.: Taiwan.

Genus *Callophlophora* THOMSON, 1864 麗紋星天牛屬

Callophlophora THOMSON, 1864, Syst. Ceramb., p. 76 (Type-species: *Oplophora sollii* HOPE, 1840).

Oplophora HOPE (nec MILNE-EDWARDS), 1839, Proc. Linn. Soc. London, 1: 43 (Type-species: *Oplophora sollii* HOPE, 1840).

Callophophora albopicta (MATSHITA, 1933) 麗艷大星斑天牛
MATSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 333, pl. 4, f. 2 (*Cyriocrates*)
Taiwan).

elegans KANO, 1933, Kontyû, 6: 279 (*Melanauster*) (Hori).

Distr.: Taiwan.

Genus *Anoplophora* HOPE, 1839 星天牛屬

Anoplophora HOPE, 1839, Proc. Linn. Soc. London, 1: 43 (Type-species: *A. stanleyana* HOPE, 1839).

Melanauster THOMSON, 1868, Physis, 2: 181 (Type-species: *Cerambyx chinensis* FORSTER, 1771).

Micromelanauster PIC, 1931, Ent. Nachr. Bl., 5: 49 (Type-species: *Monohammus bowringi* WHITE, 1858).

Subgenus *Anoplophora* HOPE, 1839, s. str.

Anoplophora (A.) *beryllina* HOPE, 1840

HOPE, 1840, Proc. Linn. Soc. London, 1: 79 (*Monochamus*) (Assam).

melanostictus WHITE, 1858, Proc. Zool. Soc. London, 24: 407 (*Monochamus*) (N. India).

argentifer PIC, 1902, Mat. Long., 4, 1: 31 (*Melanauster*) (China).

Distr.: Taiwan(?); Assam, N. India, Burma, Tonkin, China.

subsp. *bowringii* (WHITE, 1858) 保玲氏四條天牛

WHITE, 1858, Proc. Zool. Soc. London, 26: 398, pl. 53, f. 1 (*Monochammus bowringii*) (Hong Kong).

Distr.: Taiwan(?); S. China.

A. (A.) *wusheana* CHANG, 1960 霧社星天牛

CHANG, 1960, Bull. Soc. Plant Pathology and Entomology, 11(3/4): 2, pl. 1, f. 2 (Wu-she).

Distr.: Taiwan.

A. *flavomaculata* (GRESSITT, 1934) 黃星天牛

GRESSITT, 1934, Philip. J. Sci., 55: 385 (*Melanauster*) (Karapin).

Distr.: Taiwan.

A. *lurida* (PASCOE, 1856) 黑星天牛

PASCOE, 1856, Trans. Ent. Soc. London, (2)4: 47 (*Monohammus*) (N. China).

Distr.: Taiwan; China.

A. *macularia* (THOMSON, 1865) 星天牛

THOMSON, 1865, Syst. Ceramb., p. 553 (*Callophophora*) (N. China).

Distr.: Taiwan; China.

A. *nobilis* (GANGLBAUER, 1889) 黃背星天牛

GANGLBAUER, 1889, Hor. Soc. Ent. Ross., 24: 82 (*Malanauster*) (China).

luteonotatus PIC, 1925, Mel. Exot.-Ent., 43: 21 (*Melanauster*) (China).

Distr.: Taiwan; China.

A. davidis (FAIRMAIRE, 1886) 大衛星天牛

FAIRMAIRE, 1886, Ann. Soc. Ent. France, (6)6: 355 (*Melanauster*) (China).

zonator GAHAN (nec THOMSON), 1894, Ann. Mus. Civ. Genova, 34: 39 (*Melanauster*) (Burma).

Distr.: Taiwan; China, Burma, Laos, Cambodia, Thailand.

A. versteegi (RITSEMA, 1881) 灰翅星天牛

RITSEMA, 1881, Notes Leyden Mus., 3: 155 (*Monohammus*) (Sumatra).

albescens PIC, 1920, Melang Exot.-Ent. 32: 2 (*Monochamus*) (Tonkin).

Distr.: Taiwan(?); Sumatra, Burma, Assam, NE. India, Tonkin, Laos, Hainan I., China, Vietnam.

Subgenus *Cyriocrates* THOMSON, 1868

Cyriocrates THOMSON, 1868, Physis, 2: 181 (Type-species: *Oplophora horsfieldii* HOPE, 1842).

A. (Cyriocrates) horsfieldi HOPE, 1842

HOPE, 1842, Proc. Ent. Soc. London, LXI (*Oplophora*) (Assam).

voluptuosa THOMSON, 1856, Rev. Mag. Zool., (2)8: 529 (*Cerosterna*).

Distr.: Taiwan; Tonkin, Assam, China.

subsp. *tonkinensis* (KRIESCHE, 1925). 荷菲氏星天牛

KRIESCHE, 1925, Deut. Ent. Zeit., p. 287 (*Cyriocrates*) (Tonkin).

Distr.: Taiwan; Tonkin.

Genus *Eupromus* PASCOE, 1858 紅星天牛屬

Eupromus PASCOE, 1858, Trans. Ent. Soc. London, Proc., p. 12 (Type-species: *Lamia rubra* DALMAN, 1817).

Eupromus ruber (DALMAN, 1817) 紅星天牛

DALMAN, 1817, in Schonh., Syn. Ins. 1, 3, app., p. 167 (*Lamia*) (E. Indies).

sieboldi GUERIN, 1844 (Icon. Regne Anim. Ins., p. 238 (*Oplophora (Callimation)*) (Japan)).

championi WHITE, 1858, Proc. Zool. Soc. London, 26: 398 (*Monohammus*) (China).

Distr.: Taiwan; Japan, China.

Genus *Eutaenia* THOMSON, 1857 却巴氏粗天牛屬

Eutaenia THOMSON, 1857, Arch. Ent., 1: 184 (Type-species: *Ceroplesia javeti* THOMSON, 1857).

Eutaeniopsis GAHAN, 1888, Ann. Mag. Nat. Hist., (6)1: 277 (Type-species: *Lamia trifasciella* WHITE, 1850).

Eutaenia formosana MATSUSHITA, 1941 高士佛粗天牛

MATSUSHITA, 1941, Ins. Mats., 15: 156 (Kuskus).

Distr.: Taiwan.

E. trifasciella (WHITE, 1850) 三條粗天牛

WHITE, 1850, Proc. Zool. Soc. London, p. 13, pl. 13, f. 1 (*Lamia (Celosterna)*) (Hong Kong).

javeti THOMSON, 1857, Arch. Ent., 1: 125 (*Ceroplesia*) (Penang).

trifasciata THOMSON, 1864, Syst. Ceramb., p. 72.

Distr.: Taiwan; N. India, Penang, Tonkin, S. China, Hong Kong.

Genus *Cereopsius* PASCOE, 1862 白條紋天牛屬

Cereopsius PASCOE, 1862, J. Ent., 1: 344 (Type-species: *C. exoletus* PASCOE, 1862).

Etymestia PASCOE, 1866, Trans. Ent. Soc. London, (3)3: 236 (Type-species: *Cereopsius helena* WHITE, 1858).

C. shamankarriyali KANO, 1939 呂宋粗天牛

KANO, 1939, Annot. Zool. Jap., 18: 30, pl. 2, f. 3 (Luzon).

Distr.: Taiwan (Lanyu I.); Philippines.

Genus *Blepephaeus* PASCOE, 1866 疊紋粗天牛屬

Blepephaeus PASCOE, 1866, Proc. Zool. Soc. London, p. 249 (Type-species: *Monohammus succinctor* CHEVROLAT, 1852).

B. succinctor (CHEVROLAT, 1852) 疊紋粗天牛

CHEVROLAT, 1852, Rev. Mag. Zool., (2)4: 417 (*Monohammus*) (China).

sublineatus WHITE, 1958, Proc. Zool. Soc. London, 26: 410 (*Monohammus*) (India).

obfuscatus WHITE, 1858, Proc. Zool. Soc. London, 26: 411 (*Monohammus*) (Hong Kong).

fleutiauxi LAMEERE, 1893, Ann. Soc. Ent. France, 62: 283 (*Coelosterna*) (Cochin-China).

humeralis PIC, 1925, Mel. Exot.-Ent., 43: 18 (China).

Distr.: Taiwan; China, Hainan, Tonkin-China, Assam, Tenasserim, India, Siam, Malacca.

Genus *Pebblephaeus* KUSAMA et TAKAKUWA, 1984 麗紋粗天牛屬

Pebblephaeus KUSAMA et TAKAKUWA, 1984, Longicorn-Beetles of Japan, p. 449 (Type-species: *Cereopsius decoloratus* SCHWARZER, 1925).

Pebblephaeus decoloratus (SCHWARZER, 1925)

SCHWARZER, 1925, Ent. Blaett., 21: 39 (*Cereopsius*) (Sokutsu et Kosempo).

Distr.: Taiwan; Japan (Ryukyu Is.).

subsp. *decoloratus* (SCHWARZER, 1925) s. str. 麗紋粗天牛

SCHWARZER, 1925, Ent. Blaett., 21: 59 (*Cereopsius*) (Sokutsu et Kosempo).

Distr.: Taiwan.

P. ziczac (MATSUSHITA, 1940) 曲紋粗天牛

MATSUSHITA, 1940, Ins. Mats., 14: 54 (*Etymestia*) (Musha).

Distr.: Taiwan.

Genus *Parapolytretus* BREUNING, 1944 繖紋粗天牛屬

Parapolytretus BREUNING, 1944, Novit. Ent., 14, suppl., 3: 370 (Type-species: *Cycos rugosus* MATSUSHITA, 1933).

Parapolytretus rugosus (MATSUSHITA, 1933) 鐵紋粗天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 335, pl. 4, f. 10 (*Cycos* (Otsubo)).

Distr.: Taiwan.

Genus *Nanohammus* BATES, 1884 褐白星天牛屬

Nanohammus BATES, 1884, J. Linn. Soc. London, 18: 243 (Type-species: *N. rufescens* BATES, 1884).

Microcycos PIC, 1934, Melang. Exot.-Ent., 63: 9 (Type-species: *Microcycos annulicornis* PIC, 1934).

Pararhodopis BREUNING, 1935, Fol. Zool. Hydrobiol., 7: 174 (Type-species: *Rhodopis aberrans* GAHAN, 1894).

Rarasanus MATSUSHITA, 1941, Ins. Mats., 15(4): 156 (Type-species: *Rarasanus subfasciatus* MATSUSHITA, 1941).

Nanohammus subfasciatus (MATSUSHITA, 1941) 拉拉山白星天牛

MATSUSHITA, 1941, Insecta Mats., 15(4): 157, f. 3 (*Rarasanus*) (Rarasan).

taiyal GRESSITT, 1951, Longicornia, 2, p. 628 (Sankakuho).

oshimanus BREUNING et OHBAYASHI, 1964, Bull. Japan. Ent. Acad., 1(4): 15 (Japan (Amami-Oshima)).

Distr.: Taiwan; Japan (Ryukyu Is.).

Genus *Paruraecha* BREUNING, 1935 四川長角天牛屬

Paruraecha BREUNING, 1935, Fol. Zool. Hydrobiol., 8: 64 (Type-species: *P. szetschuanica* BREUNING, 1935).

Subgenus *Arisania* GRESSITT, 1936

Arisania GRESSITT, 1936, Philip. J. Sci., 61: 106 (Type-species: *A. submarmorata* GRESSITT, 1936).

Paruraecha (Arisania) submarmorata (GRESSITT, 1936) 阿里山長角天牛

GRESSITT, 1936, Philip. J. Sci., 61: 107, pl. 1, f. 16 (*Arisania*) (Arisan).

Distr.: Taiwan.

Genus *Uraecha* THOMSON, 1964 叉尾天牛屬

Uraecha THOMSON, 1964, Syst. Ceramb., pp. 84, 383 (Type-species: *Uraecha bimaculata* THOMSON, 1864).

Uraecha angusta (PASCOE, 1856) 西藏細角長天牛

PASCOE, 1856, Trans. Ent. Soc. London, (2)4: 49 (*Monohammus?*) (N. China).

attenuata PIC, 1925, Melang. Exot.-Ent., 43: 23 (Tibet).

Distr.: Taiwan; China.

Genus *Xenohammus* SCHWARZER, 1931 雙紋細角長天牛屬

Xenohammus SCHWARZER, 1931, Senckenbergiana, 13: 204 (Type-species: *X. bimaculatus* SCHWARZER, 1931).

Xenohammus bimaculatus SCHWARZER, 1931 埔里雙紋細角長天牛

SCHWARZER, 1931, Senckenbergiana, 13: 204, f. 22.

filicornis GRESSITT, 1934, Philip. J. Sci., 55: 283 (*Monochammus*) (Hori).

Distr.: Taiwan; S. China, Hainan I. (?).

X. nebulosus SCHWARZER, 1931 雲紋細角長天牛

SCHWARZER, 1931, Senckenbergiana, 13: 205, f. 15 (Kosempo).

Distr.: Taiwan.

Genus *Monochamus* GUERIN-MENEVILLE, 1826 長角天牛屬

Monochamus GUERIN-MENEVILLE, 1826, Dict. Class. d'Hist. Nat., 11: 91 (Type-species:

Lamia sutor FABRICIUS, 1775= *Cerambyx sutor* LINNE, 1758).

Meges PASCOE, 1866, Trans. Ent. Soc. London, (3)3: 272 note (Type-species: *Monochamus gravidus* PASCOE, 1858).

Monochamus kasabi HEYROVSKY, 1955 卡薩氏長角天牛

HEYROVSKY, 1955, Ann. Mus. Nat. Hung., 6: 260 (Taiwan).

Distr.: Taiwan.

M. alternatus HOPE, 1842 松斑天牛

HOPE, 1842, Trans. Ent. Soc. London, Proc., p. 111 (China)

tessellata WHITE, 1858, Proc. Zool. Soc. London, p. 408 (Hong Kong).

Distr.: Taiwan; Japan, China.

M. bimaculatus GAHAN, 1888 雙紋長角天牛

GAHAN, 1888, Ann. Mag. Nat. Hist., (6)2: 260 (India).

Distr.: Taiwan(?); India, Burma, Siam, Malacca, Laos, Tonkin, S. China.

M. fascioguttatus GRESSITT, 1938 八仙長角天牛

GRESSITT, 1938, Philip. J. Sci., 65: 156 (Hassenzan).

sintikensis MATSUSHITA, 1939, Ins. Mats., 13: 58 (Shinchiku).

Distr.: Taiwan; S. China.

M. taiheizanensis MITONO, 1943 太平山長角天牛

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 33: 580 (Taiheizan).

Distr.: Taiwan.

M. convexicollis GRESSITT, 1942 狹胸長角天牛

GRESSITT, 1942, Notes d'Ent. Chinoies, 9: 83, pl. 1, f. 3 (Tiennu Shan).

Distr.: Taiwan; E. China.

Genus *Acalolepta* PASCOE, 1858 級毛天牛屬

Acalolepta PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 247 (Type-species: *A. pusio* PASCOE, 1858, monobasic).

Cyriola THOMSON, 1864, Syst. Ceramb., p. 16 (Type-species: *C. acanthocinoides* THOMSON, 1864).

Dihammus THOMSON, 1864, Syst. Ceramb., pp. 80, 381 (Type-species: *Monochamus longicornis* THOMSON, 1857).

Haplohammus BATES, 1884, J. Linn. Soc. London, 18: 239 (Type-species: *Monochamus luxuriosus* BATES, 1873).

Neanthes PASCOE, 1878, Ann. Mag. Nat. Hist., (5)2: 372 (Type-species: *Monohammus curialis* PASCOE, 1858).

Astynoscelis PIC, 1905, Mat. Long., 5, 1: 8 (Type-species: *A. longicornis* PIC, 1905).

Niphohammus MATSUSHITA, 1932, Ins. Mats., 6(4): 170 Type-species: *N. korolensis* MATSUSHITA, 1932).

Saitoa MATSUSHITA, 1937, Kontyû, 11: 104 (Type-species: *S. teneburosa* MATSUSHITA, 1937).

Acalolepta degener (BATES, 1873) 齊藤氏灰小天牛

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 310 (*Monohammus*) (Japan).

contemptus GAHAN, 1888, Ann. Mag. Nat. Hist., (6)2: 62 (*Haplohammus*) (China).

nanus GANGLBAUER, 1889, Trudy Russk. Ent. Obshch., 24: 81 (*Haplohammus*) (China).

longicornis PIC, 1905, Mat. Long., 5(1): 8 (*Astynoscelis*) (China).

savioi PIC, 1925, Melang Exot.-Ent., 45: 21: (*Orsidis*) (China).

teneburosa MATSUSHITA, 1937, Kontyû, 11(1/2): 104 (*Saitoa*) (Korea).

Distr.: Taiwan; China, Korea, Japan.

A. flocculata (GRESSITT, 1935)

GRESSITT, 1935, Philip. J. Sci., 57: 188 (*Monochamus*) (Hinokiyama).

Distr.: Taiwan; W. China.

subsp. *flocculata* (GRESSITT, 1935), s. str. 金星絨毛天牛

GRESSITT, 1935, Philip. J. Sci., 57: 188 (*Monochamus*) (Hinokiyama).

Distr.: Taiwan.

A. formosana (BREUNING, 1935) 蓬萊絨毛天牛

BREUNING, 1935, Fol. Zool. Hydrobiol., 7: 251 (*Dihammus*) (Kosempo).

Distr.: Taiwan.

A. grossescapa (BREUNING, 1942) 恒春絨毛天牛

BREUNING, 1942, Fol. Zool. Hydrobiol., 11: 117 (*Dihammus*) (Koshun).

Distr.: Taiwan.

A. permutans (PASCOE, 1857)

PASCOE, 1857, Trans. Ent. Soc. London, (2)4: 103 (*Monohammus*) (N. China).

Distr.: Taiwan; China, Tonkin.

subsp. *paucipunctata* GRESSITT, 1938 長角絨毛天牛

GRESSITT, 1938, Philip. J. Sci., 65: 157 (*Dihammus*) (Giran et Suisha).

Distr.: Taiwan.

A. rusticatrix (FABRICIUS, 1801)

FABRICIUS, 1801, Syst. El., 2: 294 (*Lamia*) (Sumatra).

Distr.: Taiwan; Sumatra, E. Indies, India, Tonkin, Philippines.

subsp. *rusticatrix* (FABRICIUS, 1801), s. str. 鎏絨毛天牛

FABRICIUS, 1801, Syst. El., 2: 294 (*Lamia*) (Sumatra).

Distr.: Taiwan; Sumatra, E. Indies, India, Tonkin, Philippines.

subsp. *formosensis* BREUNING et OHBAYASHI, 1966 臺灣鎔絨毛天牛

BREUNING et OHBAYASHI, 1966, Bull. Jap. Ent. Acad., 2(3): 13.

Distr.: Taiwan (Lanyu I.).

A. sublusca (THOMSON, 1857)

THOMSON, 1857, Arch. Ent., 1: 293 (*Monochamus*) (Singapore).

curialis PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 246 (*Haplohammus*) (N. China).

Distr.: Taiwan; Japan (Ryukyu Is.), China, Hainan I., Indo-China, Singapore.
subsp. *maculihumera* (MATSHUSITA, 1933) 肩紋絨毛天牛

MATSHUSITA, 1933, J. Fac. Agric. Hokkaido Imp. Univ., 34(2): 328 (*Dihammus*) (Horisha).

Distr.: Taiwan.

A. ikedai MITONO, 1943 池田氏絨毛天牛

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 33: 581 (*Dihammus*) (Sozan).

vitalisi GRESSITT (nec PIC), 1951, Longicornia, 2: 404 (*Dihammus*).

Distr.: Taiwan.

A. yokoyamai HAYASHI, 1971 橫山氏絨毛天牛

HAYASHI, 1971, Ent. Rev. Japan, 23(1): 15 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

Genus *Mimorsidis* BREUNING, 1938 紅翅天牛屬

Mimorsidis BREUNING, 1938, Novit. Ent., 8: 33 (Type-species: *M. lemóulti* BREUNING, 1938).

Mimorsidis taiwanensis HAYASHI, 1974 臺灣深條天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 47 (Kenting Park).

Distr.: Taiwan.

Tribe *Batocerini* 白條天牛族Genus *Batocera* CASTELNAU, 1840 白條天牛屬

Batocera CASTELNAU, 1840, Hist. Nat. Col., 2: 470 (Type-species: *Cerambyx rubus* LINNÉ, 1758).

Batocera davidis DEYROLLE, 1878 大衛大天牛

DEYROLLE, 1878, Ann. Soc. Ent. France, (5)8: 131 (E. China).

Dist.: Taiwan; S. China, Tonkin.

B. lineolata CHEVROLAT, 1852 白條天牛

CHEVROLAT, 1852, Revue Mag. Zool., (2)4: 417 (China).

chinensis THOMSON, 1857, Arc. Nat., 1: 170 (N. China).

catenata VOLLENHOVEN, 1871, Tijdschr. Ent., p. 215.

flachi SCHWARZER, 1914, Ent. Mitt., 3: 280 (China).

hauseri SCHWARZER, 1914, Ent. Mitt., 3: 280 (China).

Distr.: Taiwan; Japan, China, Korea, Quelpart I.

B. punctata SCHWARZER, 1925 臺灣大天牛

SCHWARZER, 1925, Ent. Blaett., 21: 60 (Taiwan).

Distr.: Taiwan.

B. rubus (LINNÉ, 1758)

LINNÉ, 1758, Syst. Nat., 10: 390 (*Cerambyx*) (India).

stigma VOET, 1778, Cat. Col., 2: 15, pl. 13, f. 54 (*Lamia*) (Tonkin, China).

octomaculata FABRICIUS, 1792, Ent. Syst., 1, 2: 290 (*Lamia*) (India).

Distr.: Taiwan; India, Assam, Peninsula of SE. Asia, Sunda Is., Tonkin, China, Hainan I., Korea.

subsp. *rubus* (LINNE, 1758) s. str. 無花果天牛

LINNE, 1758, Syst. Nat., 10: 390 (*Cerambyx*) (India).

Distr.: Taiwan; India, Assam, Peninsula of SE. Asia, Sunda Is., Tonkin, China, Hainan I., Korea.

subsp. *albofasciata* (DE GEER, 1775) 白條無花果天牛

DE GEER, 1775, Mem. Ins., 5: 106, pl. 13, f. 16 (*Cerambyx*) (E. Indies).

Distr.: Taiwan; E. Indies.

B. ushijimai N. OHBAYASHI, 1981 牛島氏大天牛

N. OHBAYASHI, 1981, Ent. Rev. Japan, 35(1/2): 47 (Musha).

Distr.: Taiwan.

Genus *Apriona* CHEVROLAT, 1852 桑天牛屬

Apriona CHEVROLAT, 1852, Rev. Mag. Zool., (2)4: 414 (Type-species: *Lamia germari* HOPE, 1831).

Apriona rugicollis CHEVROLAT, 1852 桑天牛

CHEVROLAT, 1852, Rev. Mag. Zool., (2)4: 418 (Shanghai).

plicicollis MOTSCHULSKY, 1853, Etudes Ent., 2: 48 (N. China).

germari: MATSUSHITA (nec HOPE), 1933, J. Fac. Agr. Hokkaido Univ., 34: 337.

Distr.: Taiwan; China, Assam, Korea, Hainan I., Indochina, Burma, India.

Tribe Aencylonotini 胡斑天牛族

Genus *Palimna* PASCOE, 1862 長角胡斑天牛屬

Palimna PASCOE, 1862, J. Ent., 1: 346 (Type-species: *Golsinda tessellata* PASCOE, 1857 = *Cerambyx annulatus* OLIVIER, 1792).

Cylanca THOMSON, 1864, Syst. Ceramb., pp. 58, 132, 484 (Type-species: *Golsinda tessellata* PASCOE, 1857).

Goniages PASCOE, 1865, Trans. Ent. Soc. London, (3)3: 135 (Type-species: *G. infausta* PASCOE, 1865).

Apalimna BATES, 1884, J. Linn. Soc. London (Zool.), 18: 241 (Type-species: *A. liturata* BATES, 1884).

Palimna formosana (KANO, 1933) 臺灣長角胡斑天牛

KANO, 1933, Kontyû, 6: 282 (*Apalimna*) (Musha: Kyuhabon).

Distr.: Taiwan.

P. palminoides (SCHWARZER, 1925)

SCHWARZER, 1925, Ent. Blaett., 21: 62 (*Apalimna*) (Fuhosho, Hozan, Kosempo et Taihorin).

Distr.: Taiwan; S. China, Hainan I., Laos, Bhutan, Sikkim, N. India.

subsp. palimnoides (SCHWARZER, 1925) s. str. 蜘形長角天牛

SCHWARZER, 1925, Ent. Blaett., 21: 62 (*Apalimna*) (Kosempo et Taihorin).

Distr.: Taiwan.; S. China, Laos, Bhutan, Sikkim, N. India.

Tribe Crossotini 紅斑天牛族

Genus *Moechotypa* THOMSON, 1864 豔天牛屬

Moechotypa THOMSON, 1864, Syst. Ceramb., pp. 55, 368 (Type-species: *M. arida* THOMSON, 1864 = *Niphona suffusa* PASCOE, 1862).

Scotinauges PASCOE, 1871, Ann. Mag. Nat. Hist., (4)8: 277 (Type-species: *Scotinauges diphysis* PASCOE, 1871).

Tylophorus BLESSIG, 1873, Trudy Russk. Ent. Obshch., 9: 213 (Type-species: *Thylophorus wulffiusi* BLESSIG, 1873).

Moechotypa delicatula (WHITE, 1858) 阿薩姆艷天牛

WHITE, 1858, Ann. Mag. Nat. Hist., (3)2: 268 (*Nyphona*) (Assam (Sylhet)).

suffusa var. *elongata* PIC, 1925, Melang Exot.-Ent., 44: 29 (India).

Distr.: Taiwan(?); S. China, Sumatra, Assam, India, Burma, Laos, Tonkin.

M. formosana (PIC, 1917) 埔里艷天牛

PIC, 1917, Mel. Exot.-Ent., 26: 7 (*Hecyrida*) (Taiwan).

Distr.: Taiwan.

Tribe Xylorhizini 縱條天牛族

Genus *Xylorhiza* CASTELNAU, 1840 縱條天牛屬

Xylorhiza CASTELNAU, 1840, Hist. Nat. Col., 2: 476 (Type-species: *X. venosa* CASTELNAU, 1840 = *X. adusta* WIEDEMANN, 1819).

Xylorhiza adusta (WIEDEMANN, 1819) 臺灣縱條天牛

WIEDEMANN, 1819, Zool. Mag., (1)3: 182 (*Lamia*) (Bengal).

venosa CASTELNAU, 1840, Hist. Nat. Col., 2: 476 (Peninsula of SE. Asia).

Distr.: Taiwan(?); Bengal, Burma, Peninsula of SE. Asia, S. China, Hainan I.

Genus *Microxylorhiza* HAYASHI, 1974 矮縱條天牛屬

Microxylorhiza HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 38 (Type-species: *M. matsudai* HAYASHI, 1974).

Microxylorhiza matsudai HAYASHI, 1974 松田氏矮縱條天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 39 (Sungkang).

Distr.: Taiwan.

Tribe Dorcaschematini 白天牛族

Genus *Olenecamptus* CHEVROLAT, 1835 白天牛屬

Olenecamptus CHEVROLAT, 1835, Mag. Zool., 5: 134 (Type-species: *O. serratus* CHEVROLAT, 1835).

Authades THOMSON, 1857, Archives Ent., 1: 191 (Type-species: *A. indianus* THOMSON, 1857).

Ibidimorphum MOTSCHULSKY, 1860, Schrenck's Reisen Amurl. Col., p. 153 (Type-species: *I. octopustulatum* MOTSCHULSKY, 1860).

Olenecamptus bilobus (FABRICIUS, 1801) 六星白天牛

FABRICIUS, 1801, Syst. Eleuth., 2: 324 (*Saperda*) ("Insulis oceanii australis" = Northern Australia).

Distr.: Taiwan; Japan (Ryukyu Is.), Philippines, Borneo, Java, Andaman I., Nias I., Seychelles I., Laos, Sri Lanka, Caroline I., Mariana I., Molucca, New Guinea, New Britain, I. N. Australia.

O. indianus (THOMSON, 1857) 印度白天牛

THOMSON, 1857, Archiv. Ent., 1: 192 (*Authades*) (India).

Distr.: Taiwan; Japan (Ryukyu Is.), China, Tonkin, Burma, Laos, Sri Lanka, India, Sikkim, Nepal.

O. cretaceus BATES, 1873

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 314 (Japan).

Distr.: Taiwan; Japan, E. China, Korea.

subsp. *marginatus* SCHWARZER, 1925 大狹胸白天牛

SCHWARZER, 1925, Ent. Blaett., 21: 63 (*O. marginatus*) (Kosempo).

Distr.: Taiwan.

O. formosanus PIC, 1914 蓬萊狹胸白天牛

PIC, 1914, Mat. Long., 9(1): 19 (Taiwan).

decemmaculatus PIC, 1916, Mat. Long., 10(1): 13 (Hanoi, Tonkin).

Distr.: Taiwan; Japan, N. Vietnam.

Genus *Microlenecamptus* PIC, 1925 姬白天牛屬

Microlenecamptus PIC, 1925, Melang Exot.-Ent., 43: 22 (*Cylindropomus* subg.) (Type-species: *C. (M.) albonotatus* PIC, 1925).

Microlenecamptus obsoletus (FAIRMAIRE, 1888) 蒼藍蛇紋天牛

FAIRMAIRE, 1888, Revue d'Ent., 7: 146 (*Olenecamptus*) (China).

albatus MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 353 (*Dorcaschema*) (Taihorin).

Distr.: Taiwan; China.

M. biocellatus (SCHWARZER, 1925) 蛇紋天牛

SCHWARZER, 1925, Ent. Blaett., 21: 63 (*Olenecamptus*) (Kankau).

virescens SAVIO, 1929, Notes d'Ent. Chinoise, 1, 2: 8, f. 6 (*Olenecamptus*) (China).

Distr.: Taiwan; E. China.

Tribe *Xenoleini* 霜斑長角天牛族

Genus *Xenolea* THOMSON, 1864 霜斑長角天牛屬

Xenolea THOMSON, 1864, Syst. Ceramb., p. 91 (Type-species: *X. collaris* THOMSON, 1864).

Aeschopalaea PASCOE, 1864, Trans. Ent. Soc. London, (3)3: 24 (Type-species: *A. agraria* PASCOE, 1864).

Xenolea asiatica (PIC, 1925) 亞洲長角天牛

PIC, 1925, Echange, 40(41): 16 (*Aeschopalaea*) (Tonkin).

Distr.: Taiwan; Japan; S. China; Hainan; Indo-China, India.

X. collaris THOMSON, 1864

THOMSON, 1864, Syst. Ceramb., p. 92 (*Brunnea*).

Distr.: Taiwan; Borneo.

subsp. *formosana* BREUNING, 1965 蓬萊長角天牛

BREUNING, 1965, Mitt. Zool. Mus. Berlin, 41: 177.

Distr.: Taiwan.

Tribe Rhodopinini 瘤角天牛族

Genus *Rhodopina* GRESSITT, 1951 瘤角天牛屬

Rhadopina GRESSITT, 1951, Longicornia, 2: 439 (n. n. pro. *Rhodopis* THOMSON, 1857).

Rhodopis THOMSON, 1857, Archives Ent., 1: 174 (Type-species: *Rhodopis pubera* THOMSON, 1857).

R. formosana (BREUNING, 1954) 蓬萊瘤角天牛

BREUNING, 1954, Bull. Inst. Sci. Nat. Belg., 30(28): 5 (*Rhodopis*) (Hori).

Distr.: Taiwan.

R. subuniformis GRESSITT, 1951 八仙瘤角天牛

GRESSITT, 1951, Longicornia, 2: 440, pl. 18, f. 3 (Shin-Hassenzan).

Distr.: Taiwan.

Genus *Euseboides* GAHAN, 1893 尾片天牛屬

Euseboides GAHAN, 1893, Ann. Mag. Nat. Hist., (6)11: 385 (Type-species: *E. plagiatus* GAHAN, 1893).

Euseboides matsudai GRESSITT, 1938

GRESSITT, 1938, Philip. J. Sci., 65: 163, pl. 1, f. (Hori)

Distr.: Taiwan; Hainan I.

subsp. *matsudai* GRESSITT, 1938, s. str. 松田氏三帶天牛

GRESSITT, 1938, Philip. J. Sci., 65: 163, pl. 1, f. 2 (Hori).

plagiatus KANO (nec GAHAN), 1928, Trans. Nat. Hist. Soc. Formosa, 18: 408 (Karenko et Koshun).

Distr.: Taiwan.

Genus *Diboma* THOMSON, 1864 琉球銹天牛屬

Diboma THOMSON, 1864, Syst. Ceramb., p. 46 (Type-species: *D. tranquilla* THOMSON, 1864).

Zotalemimon PIC, 1925, Melang Exot.-Ent., 43: 29 (Type-species: *Z. apicale* PIC, 1925= *Sybra posticata* GAHAN, 1895).

Donysia GRESSITT, 1940, Philip. J. Sci., 72(1/2): 179 (Type-species: *Sydonia costata* MATSUSHITA, 1933).

Sybrocentrura BREUNING, 1947, Ark. Zool., 39A(6): 57 (Type-species: *S. obscura* BREUNING, 1947 = *Sydonia ropicoides* GRESSITT, 1939).

Diboma costata (MATSUSHITA, 1933) 琉球锈天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 379, pl. 5, f. 1 (*Sydonia*) (Japan (Ryukyu Is.)).

loochooana BREUNING, 1940, Folia Zool. Hydrobiol., 10: 78 (Japan (Ryukyu Is.)).

Distr.: Taiwan; Japan, Hainan I., China.

Genus *Taiwanajinga* HAYASHI, 1978 臺灣瘤角天牛屬

Taiwanajinga HAYASHI, 1978, Ent. Rev. Japan, 32(1/2): 109 (Type-species: *T. albofasciata* HAYASHI, 1978).

Taiwanajinga albofasciata HAYASHI, 1978 蓮華白條锈天牛

HAYASHI, 1978, Ent. Rev. Japan, 32(1/2): 110 (Balengshe, Nansanchi, Lien-hwachi).

Distr.: Taiwan.

Genus *Doius* MATSUSHITA, 1933 土井氏锈天牛屬

Doius MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 380 (Type-species: *D. rufescens* MATSUSHITA, 1933).

Doius fukudai HAYASHI, 1969 福田氏锈天牛

HAYASHI, 1969, Ent. Rev. Japan, 21(2): 65 (Shokoo (Matsuoka) nr. Musha).

Distr.: Taiwan.

Genus *Mimectatina* AURIVILLIUS, 1928 深褐锈天牛屬

Mimectatina AURIVILLIUS, 1928, Ark. Zool., 19A(23): 27 (Type-species: *M. singularis* AURIVILLIUS, 1927).

Nipposybra BREUNING, 1939, Festschr. 60 Geburtst E. Strand, 5: 280 (Type-species: *N. fuscoplagiata* BREUNING, 1939).

Parasydonia BREUNING, 1949, Bull. Inst. R. Sci. Nat. Belg., 25(38): 24 (Type-species: *Sydonia divaricata* BATES, 1884).

Falsodoius BREUNING, 1953, Bull. Inst. R. Sci. Nat. Belg., 29(8): 16 (Type-species: *Doius meridianus* MATSUSHITA, 1933).

Mimectatina meridiana (MATSUSHITA, 1933) 深褐锈天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 380, pl. 4, f. 8 (*Doius*) (Hozan).

Distr.: Taiwan.

M. murakamii HAYASHI, 1978 村上氏锈天牛

HAYASHI, 1978, Ent. Rev. Japan, 32(1/2): 111 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

Genus *Graphidessa* BATES, 1884 網紋锈天牛屬

Graphidessa BATES, 1884, J. Linn. Soc. London (Zool.), 18: 248 (Type-species: *G. venata* BATES, 1884).

Graphidessa obliquefasciata KOMIYA et KUSAMA, 1973 斜紋銹天牛
 KOMIYA et KUSAMA, 1973, Rep. Fac. Soc. Shizuoka Univ., 8: 138, pl. 1 (Sungkang).
 Distr.: Taiwan.

G. variegata HAYASHI, 1974 梨山銹天牛
 HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 50 (Lishan in Tachiachi).
 Distr.: Taiwan.

Genus *Pseudanaesthetis* PIC, 1922 擬褐翅天牛屬

Pseudanaesthetic PIC, 1922, Melang Exot.-Ent., 37: 15 (Type-species: *P. langana* PIC).
Hirayamaia MATSUSHITA (nec MARUNO), 1937, Kontyû, 11: 103 (Type-species: *H. fuscocrofa* MATSUSHITA, 1937).

Falsoterinaea MATSUSHITA, 1938, Kontyû, 12: 95 (n. n. for *Hirayamaia* MATSUSHITA, 1937).

Pseudanaesthetis mizunumai HAYASHI, 1974 望洋平山天牛
 HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 48 (Mareppa).
 Distr.: Taiwan.

P. rufipennis (MATSUSHITA, 1933) 紅翅平山天牛
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 377, pl. 5, f. 3 (*Eupogonius*) (Taihoku).
fuscorufa MATSUSHITA, 1937, Kontyû, 11: 103 (?*Hirayamaia*) (Musha).
 Distr.: Taiwan.

Genus *Anaesthetobrium* PIC, 1923 黑胸褐翅天牛屬

Anaesthetobrium PIC, 1923, Melang Exot.-Ent., 40: 20 (Type-species: *A. luteipenne* PIC, 1923).

Paraphidora (*Maraphidola*) MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 377 (Type-species: *P. fuscoflava* MATSUSHITA, 1933).

Anaesthetobrium fuscoflavum (MATSUSHITA, 1933) 黑胸褐翅天牛
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 377, pl. 4, f. 4 (*Paraphidola*) (Taiwan).
 Distr.: Taiwan.

Genus *Cylindilla* BATES, 1844 穎頸天牛屬

Cylindilla BATES, 1884, J. Linn. Soc. London (Zool.), 18: 250 (Type-species: *C. griseascens* BATES, 1884).

Nicroestola GRESSITT, 1940, Philip. J. Sci., 72(1/2): 180 (Type-species: *M. bidentata* GRESSITT, 1940).

Cylindilla formosana (GRESSITT, 1951) 蓬萊稜頸天牛
 GRESSITT, 1951, Longicornia, 2: 513 (*Microestola*) (Arisan).
 Distr.: Taiwan.

Genus *Penthides* MATSUSHITA, 1933 黃粗毛天牛屬

Penthides MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34(2): 430 (Type-species: *P. flavus* MATSUSHITA, 1933).

Hirakura HAYASHI, 1957, Ent. Rev. Japan, 8(2): 47 (Type-species: *H. rufoflava* HAYASHI, 1957).

Penthides flavus MATSUSHITA, 1933

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34(2): 431, pl. 5, f. 8 (Baibara).

Distr.: Taiwan; Japan (Ryukyu Is.).

subsp. *flavus* MATSUSHITA, 1933, s. str. 黑角黃天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34(2): 432, pl. 5, f. 8 (Baibara).

Distr.: Taiwan.

Genus *Miccolamia* BATES, 1884 微北海天牛屬

Miccolamia BATES, 1884, J. Linn. Soc. London, 18: 253 (Type-species: *M. cleroides* BATES, 1884).

Miccolamia castaneoverrucosa HAYASHI, 1974 褐微瘤天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 52 (Sungkang).

Distr.: Taiwan.

M. albosetosa GRESSITT, 1951 白毛微瘤天牛

GRESSITT, 1951, Longicornia, 2: 533 (Shinsui).

Distr.: Taiwan.

Tribe *Hebestolini* 鹿角天牛族

Genus *Cervoglenea* GRESSITT, 1951 鹿角白星天牛屬

Cervoglenea GRESSITT, 1951, Longicornia, 2: 621 (Type-species: ?*Glenea lata* GRESSITT, 1935).

Cervoglenea lata (GRESSITT, 1935) 粗星天牛

GRESSITT, 1935, Philip. J. Sci., 57: 191 (?*Glenea*) (Kuraru).

Distr.: Taiwan.

Tribe *Acanthocinini* 白楊粗腿天牛族

Genus *Leiops* SERVILLE, 1835 星斑寬腿天牛屬

Leiops SERVILLE, 1835, Ann. Soc. Ent. France, 4: 86 (Type-species: *Cerambyx nebulosus* LINNÉ, 1758).

Leiops shibatai HAYASHI, 1974 柴田氏寬腿天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 58 (Sungkang).

Distr.: Taiwan.

Genus *Neacanista* GRESSITT, 1940 假長角寬腿天牛屬

Neacanista GRESSITT, 1940, Philip. J. Sci., 72: 182 (Type-species: *N. tuberculipennis* GRESSITT, 1940).

Neacanista shirakii (MITONO, 1943) 素木氏寬腿天牛

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 33: 584, f. 2 (*Acanthocinus*) (Hassenzan: Reimei).

Distr.: Taiwan.

Genus *Acanthocinus* GUERIN, 1826 白楊粗腿天牛屬

Acanthocinus GUERIN, 1826, Dist. Class. D'Hist. Nat., 9: 186 (Type-species: *Cerambyx aedilis* LINNÉ, 1767).

Astynomus DEJEAN, 1835, Cat. Col. (ed. 2), p. 337 (n. n.).

Aedilis SERVILLE, 1835, Ann. Soc. Ent. France, 4: 32 (Type-species: *A. montanus* SERVILLE, 1835 = *Cerambyx aedilis* LINNÉ, 1758).

Astynomus STEPHENS, 1839, Man. Brit. Col., p. 269 (Type-species: *Cerambyx aedilis* LINNÉ, 1758).

Acanthocinus gundaiensis KANO, 1933 郡大粗腿天牛

KANO, 1933, Kontyû, 6: 285 (Gundaisha nr. Mt. Morison).

Distr.: Taiwan.

Gens *Ostedes* PASCOE, 1859 挺胸粗腿天牛屬

Ostedes PASCOE, 1859, Trans. Ent. Soc. London, (2)4: 43 (Type-species: *O. pauperata* PASCOE, 1859).

Ostedes inermis SCHWARZER, 1925

SCHWARZER, 1925, Ent. Blaett., 21: 146 (Alikang, Hozan).

nubila MATSUSHITA, 1931, Mitt. Zool. Mus. Berlin, 17: 405 (Hozan, Taihorin).

Distr.: Taiwan; Japan (Ryukyu Is.), S. China, Hainan I.

subsp. *inermis* SCHWARZER, 1925 黑紋挺胸粗腿天牛

SCHWARZER, 1925, Ent. Blaett., 21: 146 (Alikang, Hozan).

Distr.: Taiwan.

O. subfasciata MATSUSHITA, 1933 挺胸粗腿天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 392, pl. 4, f. 9 (Baibara).

Distr.: Taiwan.

Genus *Clytosemia* BATES, 1884 叉紋黑天牛屬

Clytosemia BATES, 1884, J. Linn. Soc. London, 18: 253 (Type-species: *C. pulchra* BATES, 1884).

Clytosemia bicincta GRESSITT, 1938 雙點長條天牛

GRESSITT, 1938, Philip. J. Sci., 65: 164, pl. 1, f. 3 (Sakahen nr. Karenko).

Distr.: Taiwan.

Genus *Rondibilis* THOMSON, 1857 粗腿刺翅天牛屬

Rondibilis THOMSON, 1857, Archives Ent., 1: 306 (Type-species: *R. bispinosa* THOMSON, 1857).

Polimeta PASCOE, 1864, Trans. Ent. Soc. London, (3)3: 13 (Type-species: *Ostedes spinosula* PASCOE, 1860).

Rondibilis femorata GRESSITT, 1938 欠紋刺翅天牛

GRESSITT, 1938, Philip. J. Sci., 65: 165, pl. 1, f. 1 (Rokki).

Distr.: Taiwan.

R. horiensis KANO, 1933

KANO, 1933, Kontyû, 6: 285 (Hori).

Distr.: Taiwan; China.

subsp. *horienis* KANO, 1933 s. str. 埔里刺翅天牛

KANO, 1933, Kontyû, 6: 285 (Hori).

Distr.: Taiwan.

R. semielongatus HAYASHI, 1974 六龜刺翅天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 53 (Liukuei).

Distr.: Taiwan.

Genus *Euryctosemia* HAYASHI, 1963 寬叉紋黑天牛屬

Euryctosemia HAYASHI, 1963, Ent. Rev. Japan, 16(1): 15 (Type-species: *C. nomurai* HAYASHI, 1963).

Euryctosemia nomurai HAYASHI, 1963 野村氏刺翅天牛

HAYASHI, 1963, Ent. Rev. Japan, 16(1): 15, pl. 2, f. 10 (Japan (Yonaguni I.)).

Distr.: Taiwan (Lanyu I.); Japan (Yonaguni I.).

Genus *Sophronica* BLANCHARD, 1845 水蠟鍍天牛屬

Sophronica BLANCHARD, 1845, Hist. Nat. Ins., 2: 160 (Type-species: *S. carbonaria* PASCOE, 1864).

Dasyo PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 253 (Type-species: *D. lineata* PASCOE, 1858).

Elithiotes PASCOE, 1864, J. Ent., 2: 279 (Type-species: *E. hirsuta* PASCOE, 1864).

Lasiapheles BATES, 1873, Ann. Mag. Nat. Hist., (4)12: (Type-species: *L. obrioides* BATES, 1873).

Dasystola KOLBE, 1894, Stettin. Ent. Ztg., 55: 63 (Type-species: *D. hirta* KOLBE, 1894).

Phunginus PIC, 1922, Melang Exot.-Ent., 37: 15 (Type-species: *P. apicalis* PIC, 1922).

Mimanaesthetis PIC, 1926, Melang Exot.-Ent., 47: 16 (Type-species: *M. atripennis* PIC, 1926).

Eupogonioides FISHER, 1930, Indian Forest Rec., 14: 275 (Type-species: *E. gardneri* FISHER, 1930 = *Phunginus apicalis* PIC, 1922).

Sophronica obrioides (BATES, 1873) 水蠟鍍天牛

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 382 (Japan).

Distr.: Taiwan; Japan.

Genus *Anaespogonius* GRESSITT, 1938 寬胸瘤角天牛屬

Anaespogonius GRESSITT, 1938, Lingnan Sci. J., 17: 53 (Type-species: *A. fulvus* GRESSITT, 1938).

Anaespogonius ochraceofulvus HAYASHI 1974 寬胸瘤角天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 49 (Taichung).

Distr.: Taiwan.

Genus *Myrmexocentrus* BREUNING, 1957 蟻型天牛屬

Myrmexocentrus BREUNING, 1957, Bull. Inst. Roy. Sci. Nat. Belg., 33, 8: 20 (Type-species: *M. tibialis* BREUNING, 1957).

Myrmexocentrus quadrimaculatus HAYASHI, 1983 四紋蟻型天牛

HAYASHI, 1983, Bull. Osaka Jonan Women's Jr. Coll., 16: 43 (Kenting Park).

Distr.: Taiwan.

Genus *Eryssamena* BATES, 1884 刺翅天牛屬

Eryssamena BATES, 1884, J. Linn. Soc. London (Zool.), 18: 251 (Type-species: *E. saperdina* BATES, 1884).

Eryssamena taiwana HAYASHI, 1974 臺灣刺翅天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 54 (Fenchih).

Distr.: Taiwan.

E. shibatai HAYASHI, 1974 柴田氏細條天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 55 (Fenchih).

Distr.: Taiwan.

Genus *Neoeryssamena* HAYASHI, 1974 新刺翅天牛屬

Neoeryssamena HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 56 (Type-species: *N. mitonoana* HAYASHI, 1974).

Neoeryssamena mitonoana HAYASHI, 1974 水戶野氏細條天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 56 (Wushe).

Distr.: Taiwan.

Genus *Eoporis* PASCOE, 1864 灰粗腿天牛屬

Eoporis PASCOE, 1864, Trans. Ent. Soc. London, (3)3: 15 (Type-species: *E. elegans* PASCOE, 1864).

Subgenus *Eoporimimus* SCHWARZER, 1925

Eoporimimus SCHWARZER, 1925, Ent. Blaett., 21: 147 (Type-species: *E. bifasciana* SCHWARZER, 1925).

Eoporis (Eoporimimus) simillima HAYASHI, 1974 南山灰粗腿天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 57 (Nanshanchi).

Distr.: Taiwan.

E. (E.) mitonoi (SEKI, 1946) 水戶野氏灰粗腿天牛

SEKI, 1949, Ins. World, L: 573, f. (*Rondibilis*) (Mt. Hinoki).

Distr.: Taiwan.

E. (E.) bifasciana SCHWARZER, 1925 雙彩灰粗腿天牛

SCHWARZER, 1925, Ent. Blaett., 21: 147 (Sokutsu).

Distr.: Taiwan.

E. (E.) differens PIC, 1926 寮國灰粗腿天牛

PIC, 1926, Bull. Soc. Ent. Fr., p. 143.

Distr.: Taiwan; China, India, N. Vietnam, Laos.

Genus *Exocentrus* MULSANT, 1839 粗毛瑣天牛屬

Exocentrus MULSANT, 1839, Col. France Long., p. 152 (Type-species: *Cerambyx lusitanus* LINNÉ, 1767).

- Campomyne* PASCOE, 1864, Trans. Ent. Soc. London, (3)131: 27, 43 (Type-species: *C. calliooides* PASCOE, 1864).
- Oligopsis* THOMSON, 1864, Syst. Ceramb., p. 111 (Type-species: *O. exocentroides* THOMSON, 1864).
- Ispaterus* FAIRMAIRE, 1892, Revue Ent., 11: 122 (Type-species: *Ispaterus longipilis* FAIRMAIRE, 1892).
- Pseudocentrus* FAIRMAIRE, 1901, Revue Ent., 20: 230 (Type-species: *P. lineellus* FAIRMAIRE, 1901 = *Exocentrus reticulatus* FAIRMAIRE, 1896).
- E. alboguttatus* FISHER, 1925
FISHER, 1925, Philip. J. Sci., 28(2): 240.
Distr.: Taiwan; Hainan I., Bhutan, Andaman I., Sumatra, Laos, Tonkin, China.
subsp. *taiwanensis* KUSAMA et TAHIRA, 1978 白星瑣天牛
KUSAMA et TAHIRA, 1978, Elytra, 6(1): 22 (Liukuei).
Distr.: Taiwan.
- E. brevisetosus* GRESSITT, 1938 縮毛瑣天牛
GRESSITT, 1938, Philip. J. Sci., 65: 167, pl. 1, f. 4 (*E. testudineus* subsp.) (Bukai).
Distr.: Taiwan; W. China.
- E. formosofasciolatus* KUSAMA et TAHIRA, 1978 細帶瑣天牛
KUSAMA et TAHIRA, 1978, Elytra, 6(1): 17 (Nanshanchi).
Distr.: Taiwan.
- E. fulvobrunneus* KUSAMA et TAHIRA, 1978 桔草瑣天牛
KUSAMA et TAHIRA, 1978, Elytra, 6(1): 15 (Roshan).
Distr.: Taiwan.
- E. immaculatus* GRESSITT, 1951 欠紋瑣天牛
GRESSITT, 1951, Longicornia, 2: 525, 529 (Kuraru).
Distr.: Taiwan.
- E. kentingensis* KUSAMA et TAHIRA, 1978 墾丁瑣天牛
KUSAMA et TAHIRA, 1978, Elytra, 6(1): 12 (nr. Kenting Park).
Distr.: Taiwan.
- E. nanshanensis* KUSAMA et TAHIRA, 1978 南山瑣天牛
KUSAMA et TAHIRA, 1978, Elytra, 6(1): 14 (Nanshanchi).
Distr.: Taiwan.
- E. pseudovariepennis* KUSAMA et TAHIRA, 1978 廬山多紋瑣天牛
KUSAMA et TAHIRA, 1978, Elytra, 6(1): 18 (Roshan).
Distr.: Taiwan.
- E. rufithorax* GRESSITT, 1935 紅胸瑣天牛
GRESSITT, 1935, Trans. Nat. Hist. Soc. Formosa, 25: 236 (Pianan-Ambu).
Distr.: Taiwan.
- E. savioi* PIC, 1925 短翅瑣天牛
PIC, 1925, Bull. Soc. Ent. France, p. 138 (*E. curtipennis* var. *savioi*) (China).
Distr.: Taiwan; E. China.

E. seriatomaculatus SCHWARZER, 1925 後紋圓瑣天牛

SCHWARZER, 1925, Ent. Blaett., 21: 147 (Kankau).

Distr.: Taiwan.

E. taniguchii MAKIHARA, 1986 谷口氏瑣天牛

MAKIHARA, 1986, Mis. Rep. Hiwa Mus. Nat. Hist., 24: 13 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

E. variepennis (SCHWARZER, 1925) 斑翅瑣天牛

SCHWARZER, 1925, Ent. Blaett., 21: 147 (*Camptomyne?*) (Kankau).

Distr.: Taiwan.

E. venatoides KUSAMA et TAHIRA, 1978 梅峯瑣天牛

KUSAMA et TAHIRA, 1978, Elytra, 6(1): 10 (nr. Meifeng).

Distr.: Taiwan.

Genus *Sciades* PASCOE, 1864 刺胸天牛屬

Sciades PASCOE, 1864, Trans. Ent. Soc. London, (3)3: 27, 30 (Type-species: *Leiopus suffusus* PASCOE, 1859).

Subgenus *Estoliops* MATSUSHITA, 1943

Estoliops MATSUSHITA, 1943, Trans. Nat. Hist. Soc. Formosa, 33(242/243): 574 (Type-species: *E. fasciatus* MATSUSHITA, 1943).

Sciades (Estoliops) fasciatus (MATSUSHITA, 1943)

MATSUSHITA, 1943, Trans. Nat. Hist. Soc. Formosa, 33(242/243): 575, f. 3 (*Estoliops*) (Japan).

Distr.: Taiwan; Japan.

subsp. *taiwanensis* KUSAMA et ODA, 1978 黑帶刺胸天牛

KUSAMA et ODA, 1978, Rep. Fac. Soc. Shizuoka Univ., 13: 76, pl. 1, f. 4 (*Estoriops*) (Hui-sun Exp. Forest).

Distr.: Taiwan.

Subgenus *Miaenia* PASCOE, 1864

Miaenia PASCOE, 1864, Trans. Ent. Soc. London, (3)3: 27 (Type-species: *M. marmorea* PASCOE, 1864).

S. (Mienia) botelensis (GRESSITT, 1951) 紅頭瑣天牛

GRESSITT, 1951, Longicornia, 2: 531 (*Nonymoides*) (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

S. (Miaenia) subfasciata (SCHWARZER, 1925) 灰帶瑣天牛

SCHWARZER, 1925, Ent. Blaett., 21: 148 (*Miaenia*) (Sokutsu).

Distr.: Taiwan.

Subgenus *Granulimiaenia* BREUNING, 1977

BREUNING, 1977, Mitt. Zool. Mus. Berlin, 53(1): 241, 159 (Type-species: *Miaenia granulicollis* GRESSITT, 1938).

S. (Granulimiaenia) granulicollis (GRESSITT, 1938) 裸胸瑣天牛

GRESSITT, 1938, Philip. J. Sci., 65: 166, pl. 1, f. 5 (*Miaenia*) (Kuraru).

Distr.: Taiwan; S. China.

S. (G.) *changi* (KUSAMA et ODA, 1975) 張氏瑣天牛

KUSAMA et ODA, 1975, Rep. Fac. Soc. Shizuoka Univ., 13: 75 pl. 1, f. 3 (*Miaenia*)
(Hui-Sun Exp. Forest).

Distr.: Taiwan.

Tribe Astathini 藍艷天牛族

Genus *Cleonaria* THOMSON, 1864 藍艷黃天牛屬

Cleonaria THOMSON, 1864, Syst. Ceramb., p. 119 (Type-species: *C. bicolor* THOMSON, 1864).

Cleonaria taiwana HAYASHI, 1984 瑞穗黃天牛

HAYASHI, 1984, Bull. Osaka Jonan Women's Jr. Coll., 17/18: 36, pl. 2, f. 9 (Juisui).
Distr.: Taiwan.

Genus *Bacchisa* PASCOE, 1867 琉璃天牛屬

Bacchisa PASCOE, 1867, Trans. Ent. Soc. London, (3)3: 329, 342 (Type-species: *B. coronata* PASCOE, 1867).

Bacchisa atritarsis (PIC, 1912) 臺灣琉璃天牛

PIC, 1912, Mat. Long. 8, 2: 21 (*Chreonomma*) (China?).
Distr.: Taiwan; China.

B. fortunei (THOMSON, 1857)

THOMSON, 1857, Archives Ent., 1: 58, pl. 6, f. 2 (*Plaxomicrus*) (China).

Distr.: Taiwan; China, Korea, Japan.

subsp. *flavicornis* KANO, 1933 福晉氏琉璃天牛

KANO, 1933, Kontyû, 6: 291 (*Chreonomma fortunei* subsp. *flavicornis*) (Urai).

Distr.: Taiwan.

Genus *Anastathes* GAHAN, 1901 四紋黃天牛屬

Anastathes GAHAN, 1901, Trans. Ent. Soc. London, p. 60 (Type-species: *Astathes nigricornis* THOMSON, 1865).

Anastathes parva GRESSITT, 1935

GRESSITT, 1935, Philip. J. Sci., 57: 193 (Bukai et Hori).

Distr.: Taiwan; China, Hainan I.

subsp. *parva* GRESSITT, 1935, s. str. 四紋黃天牛

GRESSITT, 1935, Philip. J. Sci., 57: 193 (Bukai et Hori).

Distr.: Taiwan.

Genus *Astathes* NEWMAN, 1842 藍艷天牛屬

Astathes NEWMAN, 1842, Entomologist, 1: 299 (Type-species: *A. perplexa* NEWMAN, 1842).

Astathes episcopalis CHEVROLAT, 1852 大紫天牛

CHEVROLAT, 1852, Revue Zool., (2): 418 (China).

Distr.: Taiwan; China.

A. formosana BREUNING, 1956 蓬萊紫天牛
BREUNING, 1956, Longicornia, 3: 499 (Taihoku).

Distr.: Taiwan.

A. janthinipennis FAIRMAIRE, 1895
FAIRMAIRE, 1895, Ann. Soc. Ent. Belg., 39: 187.
Distr.: Taiwan; Tonkin, Hainan I., China.

subsp. *cyanoptera* GAHAN, 1900 紅翅紫天牛
GAHAN, 1900, Ann. Mag. Nat. Hist., (7)5: 353 (Hainan).
Distr.: Taiwan.

Tribe Saperdini 多星天牛族

Genus *Thyestilla* AURIVILLIUS, 1923 大麻天牛屬

Thyestilla AURIVILLIUS, 1923, Col. Cat., 74: 491 (n. n. pro. *Thyestes* THOMSON, 1864)
(Type-species: *T. pubescens* THOMSON, 1864).

Thyestilla gebleri (FALDERMANN, 1835) 大麻天牛
FALDERMANN, 1835, Mem. Acad. Sci. St. Petersb., 2: 434, pl. 5, f. 6 (*Saperda*) (N. China).
pubescens THOMSON, 1964, Syst. Ceramb., p. 116 (*Thyestes*) (Japan).
funebris GAHAN, 1888, Ann. Mag. Nat. Hist., (6)2: 67 (*Thyestes*) (China).
infernalis PIC, 1904, Echange, 10(231): 17 (*Phytoecia*) (Amur).
Lepesmei GILMOUR, 1950, Longicornia, 1: 554, f. 10 (China).
curtipennis PIC, 1952. Div. Ent., 11: 5 (*Phytoecia*).
gebleri ssp. *kadowakii* FUJIMURA, 1962, Kontyû, 30(3): 211, pl. 13, f. 26 (Japan (Okinawa Is.)).
Distr.: Taiwan(?); Japan, Korea, Siberia.

Genus *Glenida* GAHAN, 1888 多星妮天牛屬

Glenida GAHAN, 1888, Ann. Mag. Nat. Hist., (6)2: 65 (Type-species: *G. suffusa* GAHAN, 1888).

Glenida cyanipennis GAHAN, 1888 藍翅妮天牛
GAHAN, 1888, Ann. Mag. Nat. Hist., (6)2: 66 (China).
Distr.: Taiwan; S. China, Hainan I.

Genus *Mandibularia* PIC, 1925 巨顎天牛屬

Mandibularia PIC, 1925, Melang Exot.-Ent., 43: 24 (Type-species: *M. nigriceps* PIC, 1925).

Mandibularia quadricolor GRESSITT, 1951 多彩巨顎天牛
GRESSITT, 1951, Longicornia, 2: 561, pl. 21, f. 4 (SE. China).
Distr.: Taiwan; China.

Genus *Serixia* PASCOE, 1856 細角多星天牛屬

Serixia PASCOE, 1856, Trans. Ent. Soc. London, (2)4: 45 (Type-species: *S. apicalis* PASCOE, 1856).

Serixia albopleura GRESSITT, 1935 白側細角天牛

GRESSITT, 1935, Trans. Nat. Hist. Soc. Formosa, 25: 291 (Hassenzan).

Distr.: Taiwan.

S. botensis KANO, 1933 蘭嶼細角天牛

KANO, 1933, Kontyû, 6: 288 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

S. kisana (MATSUSHITA, 1937) 旗山細角天牛

MATSUSHITA, 1937, Kontyû, 11: 105 (*Neopraolea*) (Kisan).

?*longicornis* SCHWARZER (nec PASCOE), 1925, Ent. Blaett., 21: 149 (Kankau).

subsericea GRESSITT, 1938, Philip. J. Sci., 65: 170, pl. 1, f. 11 (Kuraru).

Distr.: Taiwan.

S. signaticornis SCHWARZER, 1925 環節細角天牛

SCHWARZER, 1925, Ent. Blaett., 21: 149 (Kankau).

Distr.: Taiwan.

S. testaceicollis KANO, 1933 褐胸細角天牛

KANO, 1933, Kontyû, 6: 287 (Urai).

Distr.: Taiwan.

S. hayashii, HAYASHI, 1974 林氏細角天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 60 (Nanshanchi).

Distr.: Taiwan.

S. juisuiensis HAYASHI, 1974 瑞穗細角天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 61 (Juisui).

Distr.: Taiwan.

Genus *Praolia* BATES, 1884 長角藍艷天牛屬

Praolia BATES, 1884, J. Linn. Soc. London (Zool.), 18: 261 (Type-species: *P. citrinipes* BATES, 1884).

Praolia atripennis (PIC, 1923) 黑翅細角天牛

PIC, 1923, Melang Exot.-Ent., 40: 20 (*Serixia*) (Taiwan).

griseipennis GRESSITT, 1938, Philip. J. Sci., 65: 169, pl. 1, f. 8, (*Serixia*) (Karenko: Sakahen).

Distr.: Taiwan.

P. citrinipes BATES, 1884 藍艷細角天牛

BATES, 1884, J. Linn. Soc. London (Zool.), 18: 261, pl. 2, f. 8 (Japan).

Distr.: Taiwan(?); Japan.

Genus *Neoserixia* SCHWARZER, 1925 長頸黃天牛屬

Neoserixia SCHWARZER, 1925, Ent. Blaett., 21: 148 (Type-species: *N. pulchra* SCHWARZER, 1925).

Subgenus *Neoserixia* SCHWARZER, 1925, s. str.

Neoserixia (*Neoserixia*) *longicollis* GRESSITT, 1935 長頸黃天牛

GRESSITT, 1935, Trans. Nat. Hist. Soc. Formosa, 25: 287 (Rokki: Chirifu).

Distr.: Taiwan.

N. (N.) infasciatus GRESSITT, 1935 姬長頸黃天牛
 GRESSITT, 1935, Trans. Nat. Hist. Soc. Formosa, 25: 289 (*N. longicollis* subsp.
infasciatus) (Hassenzan, Fukai).

Distr.: Taiwan.

N. (N.) pulchra SCHWARZER, 1925 港口黃天牛
 SCHWARZER, 1925, Ent. Blaett., 21: 148 (Kankau, Banshoryo).
 Distr.: Taiwan; S. China.
 subsp. *pulchra* SCHWARZER, 1925, s. str.
 SCHWARZER, 1925, Ent. Blaett., 21: 148 (Kankau, Banshoryo).
 Distr.: Taiwan.

N. (N.) schwarzeri GRESSITT, 1935 舒瓦氏黃天牛
 GRESSITT, 1935, Trans. Nat. Hist. Soc. Formosa, 25: 289 (Sakahen-Higashinoko).
 Distr.: Taiwan.

Subgenus *Neopraolia* MATSUSHITA, 1933

Neopraolia MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 406 (Type-species:
Neoserixia delicata MATSUSHITA, 1933).

Neoserixia (*Neopraolia*) *delicata* MATSUSHITA, 1933 凤山細角天牛
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 406 (*Neopraolia*) (Hozan).
 Distr.: Taiwan.

Genus *Menesia* MULSANT, 1856 黃紋天牛屬

Menesia MULSANT, 1856, Ann. Soc. Linn. Lyon, (2)3: 157 (Type-species: *M. perrisi*
 MULSANT, 1856 = *Saperda bipunctata* ZOOBKOFF, 1829).

Daphisia PASCOE, 1867, Trans. Ent. Soc. London, (3)3: 363, 418 (Type-species: *D. pulchella* PASCOE, 1867).

Menesia matsudai HAYASHI, 1985 松田氏黃紋天牛
 HAYASHI, 1985, Ent. Rev. Japan, 40(2): 135 (Lienhwachi).
 Distr.: Taiwan.

Genus *Paramenesia* BREUNING, 1952 似黃紋天牛屬

Paramenesia BREUNING, 1952, Ent. Arb. Mus. Georg. Frey, 3: 135 (Type-species: *P. theaphia* BATES, 1884).

Paramenesia kasugensis (SEKI et KOBAYASHI, 1935) 春日黃紋天牛
 SEKI et KOBAYASHI, 1935, Ent. Wld. Tokyo, 3(16): 225, f. (*Glenea*) (Japan).
 Distr.: Taiwan (new record: Baikei, 25 VI 1939).

Genus *Eutetrapha* BATES, 1884 華夏天牛屬

Eutetrapha BATES, 1884, J. Linn. Soc. London (Zool.), 18: 256 (Type-species: *E. variicornis* BATES, 1884 = *Saperda sedecimpunctata* MOTSCHULSKY, 1860).

Saperda ABDULLAN et ABDULLAN, 1966 (nec FABRICIUS, 1775), Proc. R. Ent. Soc. London, (B)35: 87.

Eutetrapha elegans HAYASHI, 1966 和社緣天牛
 HAYASHI, 1966, Bull. Osaka Jonan Women's Jr. Coll., 1: 9, f. 6 (Hosakei).
 Distr.: Taiwan.

Genus *Saperda* FABRICIUS, 1775 多星天牛屬

Saperda FABRICIUS, 1775, Syst. Ent., p. 184 (Type-species: *Cerambyx scalaris* LINNÉ, 1758).

Saperda kojimai MAKIHARA et NAKAMURA, 1985 小島氏黑肩天牛
MAKIHARA et NAKAMURA, 1985, Mis. Rep. Hiwa Mus. Nat. Hist., 23: 18 (Huanshan).
Distr.: Taiwan.

Tribe Gleneini 白星天牛族

Genus *Paraglenea* BATES, 1866 莎麻天牛屬

Paraglenea BATES, 1886, Proc. Zool. Soc. London, p. 352 (Type-species: *Glenea fortunei* SAUNDERS, 1853).

Paraglenea swinhoei BATES, 1866 黑紋蒼藍天牛
BATES, 1866, Proc. Zool. Soc. London, p. 352, f. 3 (Taiwan).
Distr.: Taiwan.

P. fortunei (SAUNDERS, 1853) 莎麻天牛
SAUNDERS, Trans. Ent. Soc. London, (2)2: 112, pl. 4, f. 1 (*Glenea*) (N. China).
chloromelas THOMSON, 1879, Revue Mag. Zool., (3)7: 21 (*Glenea*) (China).
Distr.: Taiwan; Japan, China, Tonkin.

Genus *Pareutetrapha* BREUNING, 1952 蒼藍天牛屬

Pareutetrapha BREUNING, 1952, Ent. Arb. Mus. Georg. Frey, 3: 135 (Type-species: *P. eximia* BATES, 1884).

Pareutetrapha magnifica (SCHWARZER, 1925)
SCHWARZER, 1925, Ent. Bl. Biol. Syst. Kafer, 21: 148 (*Saperda*) (Kosempo).
Distr.: Taiwan; Japan (Ryukyu Is.).
subsp. *magnifica* (SCHWARZER, 1925) s. str. 胸紋蒼藍天牛
SCHWARZER, 1925, Ent. Bl. Biol. Syst. Kafer, 21: 148 (*Saperda*) (Kosempo).
Distr.: Taiwan.

Genus *Glenea* NEWMAN, 1842 白星天牛屬

Glenea NEWMAN, 1842, Entomologist, 1: 301 (n. n. pro. *Sphenura* CASTELNAU, 1840).
Sphenura CASTELNAU, 1840, Hist. Nat. Ins., Gen. Col., 2: 489 (Type-species: *S. novembuttata* GUERIN, 1831).

Glenea anulicornia SCHWARZER, 1925 新島氏黃條天牛
SCHWARZER, 1925, Ent. Blaett., 21: 149 (Kankau).
nijimai MATSUSHITA, 1938, Kontyû, 12: 94 (*G. (Macroglenea)*) (Hori).
Distr.: Taiwan.

G. subabbreviata BREUNING, 1956 武界黃條天牛
BREUNING, 1956, Ent. Arb., 7: 791 (n. n. for *G. abbreviata* GRESSITT, 1951, *Longicornia*, 2: 573) (Bukai).
Distr.: Taiwan.

G. acutoides SCHWARZER, 1925 環紋黃條天牛

SCHWARZER, 1925, Ent. Blaett., 21: 151 (Kosempo).

obliqua GRESSITT, 1939, Notes d'Ent. Chinoise, 6: 119, pl. 2, f. 9-11 (China).

roubali HEYROVSKY, 1939, Casopis C. Spo. Ent., 36: 68 (China).

Distr.: Taiwan; China.

G. centroguttata FAIRMAIRE, 1897 一色氏黃紋天牛

FAIRMAIRE, 1897, Notes Leyden Mus., 19: 232 (Tibet).

issikii MITONO, 1934, Trans. Nat. Hist. Soc. Formosa, 24: 490, f. 1 (Shinchiku: Shiigao, Toryo, Doka, Kahodai & Piyasan).

Distr.: Taiwan; Japan, W. China.

G. chlorospila GAHAN, 1897 琉球藍星天牛

GAHAN, 1897, Ann. Mag. Nat. Hist., (6)19: 488 (Japan (Ryukyu Is.)).

Distr.: Taiwan; Japan.

G. chrysomaculata SCHWARZER, 1925 藍艷燦星天牛

SCHWARZER, 1925, Ent. Blaett., 21: 152 (Sokutsu, Alikang, Kosempo).

Distr.: Taiwan.

G. chujoi MITONO, 1937 桑黃紋天牛

MITONO, 1937, Mushi, 10: 149, f. (Shinchiku: Seto).

Distr.: Taiwan.

G. tonkinea AURIVILLIUS, 1926

AURIVILLIUS, 1926, Ark. Zool., 18(A9): 19, f. (Tonkin).

Distr.: Taiwan; Tonkin, Hainan I.

subsp. *diversenotata* SCHWARZER, 1925 海南白星天牛

SCHWARZER, 1925, Ent. Blaett., 21: 152 (Kosempo).

Distr.: Taiwan.

G. fainanensis PIC, 1916 曇斑條天牛

PIC, 1916, Mat. Long., 10, 1: 17 (Fainan).

horishana MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 412 (Hori).

pieliana subsp. *nigra* GRESSITT, 1940, Notes d'Ent. Chinoise, 7: 194, pl. 5, f. 5 (China).

Distr.: Taiwan.

G. iwasakii KANO, 133 岩崎氏金條天牛

KANO, 1933, Kontyû, 7: 139 (Japan (Ryukyu Is.)).

stolata MATSUMURA et MATSUSHITA, 1933, J. Fac. Agri. Hokkaido Univ., 34: 413, f. 7 (Japan (Ryukyu Is.)).

Distr.: Taiwan; Japan (Ryukyu Is.).

G. lacteomaculata SCHWARZER, 1925 白紋星天牛

SCHWARZER, 1925, Ent. Blaett., 21: 151 (Kosempo, Sokutsu).

Distr.: Taiwan.

G. sauteri SCHWARZER, 1925 曹太氏黃紋天牛

SCHWARZER, 1925, Ent. Blaett., 21: 150 (Fuhosho, Alikang, Kosempo).

Distr.: Taiwan; Japan (Ryukyu Is.).

G. luteicollis GRESSITT, 1935 黃胸欠紋天牛

GRESSITT, 1935, Philip. J. Sci., 57: 190 (Hassenzan).

Distr.: Taiwan.

G. mitonoana GRESSITT, 1951 藍星黃紋天牛

GRESSITT, 1951, Longicornia, 2: 576 (n. n.)

issikii var. *ornata* MITONO (nec GAHAN, 1889), 1934, Trans. Nat. Hist. Soc. Formosa, 24: 491, f. 2 (Karenko).

Distr.: Taiwan.

G. miwai MITONO, 1943

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 38: 585, f. 3 (Baibara, Seto, Chushinron).

Distr.: Taiwan; China.

subsp. *miwai* MITONO, 1943, s. str. 三輪氏錯紋天牛

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 38: 585, f. 3 (Baibara, Seto, Chushinron).

Distr.: Taiwan.

G. relicta PASCOE, 1868

PASCOE, 1868, Trans. Ent. Soc. London, (2)4: 258 (China).

Distr.: Taiwan; Japan, Saghalien, China, Korea, Assam, Hainan I., Tonkin.

subsp. *silhetica* PLAVILSTCHIKOV, 1927 褐肩白星天牛

PLAVILSTCHIKOV, 1927, Encyc. Ent. B. Col. 2, 2: 63 (n. n.).

assimilis GAHAN (nec JORDAN), 1897, Ann. Mag. Nat. Hist., (6)19: 494 (Assam)

Distr.: Taiwan; Assam, China, Hainan I.

subsp. *formosensis* BREUNING, 1960 臺灣褐肩白星天牛

BREUNING, 1960, Abh. Mus. Tierk. Dresden (Ent.), 25: 15.

Distr.: Taiwan.

G. versuta NEWMAN, 1842 蘭嶼白條天牛

NEWMAN, 1842, Entomologist, 1: 302 (Philippines).

Distr.: Taiwan (Lanyu Is.) (?); Philippines.

G. formosana SCHWARZER, 1925 蓬萊白條天牛

SCHWARZER, 1925, Ent. Blaett., 21: 150 (Kankau).

Distr.: Taiwan.

G. mounieri PIC, 1926

PIC, 1926, Melang Exot.-Ent., 47: 22 (Annam).

Distr.: Taiwan; Annam, Tonkin, Laos.

subsp. *horiensis* BREUNING, 1953 莫尼氏天牛

BREUNING, 1953, Bull. Inst. Sci. Nat. Belg., 29(8): 21 (Hori).

Distr.: Taiwan.

G. nanshanchiana HAYASHI, 1978 南山白條天牛

HAYASHI, 1978, Ent. Rev. Japan, 31(1/2): 89 (Nanshanchi).

Distr.: Taiwan.

Genus *Heteroglenea* GAHAN, 1897 長白條天牛屬

Heteroglenea GAHAN, 1897, Ann. Mag. Nat. Hist., (6)19: 490 (Type-species: *G. glechoma* PASCOE, 1867).

Heteroglenea glechoma (PASCOE, 1867) 長白條天牛

PASCOE, 1867, Trans. Ent. Soc. London, (3)3: 409 (*Matabello*).

Distr.: Taiwan (Lanyu I.) (?); Sumatra, Philippines, New Britain, Solomon Is.

Tribe *Phytoeciini* 菊花天牛族Genus *Nupserha* THOMSON, 1869 黑緣蘋果天牛屬

Nupserha THOMSON, 1860, Classif. Ceramb., p. 41 (Type-species: *Stibara cosmopolita* THOMSON, 1857).

Nupserha clypealis (FAIRMAIRE, 1895)

FAIRMAIRE, 1895, Ann. Soc. Ent. Belg., 39: 188 (*Oberea*) (Tonkin).

bisbinotaticollis PIC, 1917, Mat. Longic., 10(2): 13 (*Oberea*).

Distr.: Taiwan; N. Vietnam, China, Laos.

subsp. *formosana* BREUNING, 1960 胸紋蘋果天牛

BREUNING, 1960, Bull. Inst. Sci. Nat. Belg., 36(19): 19.

Distr.: Taiwan.

N. fricator (DALMAN, 1817) 黑尾細蘋果天牛

DALMAN, 1817, in Schonherr, Syn. Ins., 1, 3: 183 (*Saperda*) (Java).

Distr.: Taiwan(?); Burma, Hainan I., Malacca, Java, Borneo, Celebes.

N. kankauensis (SCHWARZER, 1925) 港口姬蘋果天牛

SCHWARZER, 1925, Ent. Blaett., 21: 153 (*Oberea marginella* var.) (Kankau).

Distr.: Taiwan; Hainan I., S. China.

Genus *Eumecocera* SOLSKY, 1871 僞蘋果天牛屬

Eumecocera SOLSKY, 1871, Hor. Soc. Ent. Ross., 7: 391 (Type-species: *Saperda impustulata* MOTSCHULSKY, 1860).

Eumecocera lineata (GRESSITT, 1951) 四條藍天牛

GRESSITT, 1951, Longicornia, 2: 609, pl. 21, f. 8 (*Stenostola*) (China).

Distr.: Taiwan; China.

Genus *Obereopsis* CHEVROLAT, 1885 巨眼蘋果天牛屬

CHEVROLAT, 1855, Rev. Zool. Ser. 2, 7: 289 (Type-species: *O. obscuritarsis* CHEVROLAT, 1855).

Paroberea KOLBE, 1893, Stett. Ent. Ztg., 54: 79.

Obereopsis bicoloripes (PIC, 1916) 黑尾黃翅蘋果天牛

PIC, 1916, Mat. Long., 10, 1: 16 (*Oberea*) (Taiwan).

Distr.: Taiwan.

O. lineaticeps (PIC, 1911) 背條蘋果天牛

PIC, 1911, Mat. Long., 8, 1: 21 (*Oberea*) (Taiwan).

Distr.: Taiwan; E. China.

O. signaticornis (MATSUSHITA, 1933) 環紋蘋果天牛

MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 423, pl. 4, f. 10 (*Oberea*)
(Kanko).

Distr.: Taiwan.

Genus *Dystomorphus* PIC, 1926 寬頭天牛屬

Dystomorphus PIC, 1926, Melang Exot.-Ent., 47: 11 (Type-species: *D. notatus* PIC, 1926).

Dystomorphus esakii HAYASHI, 1974 江崎氏寬頭天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 59 (Saramao-Matsumine).

Distr.: Taiwan.

Genus *Oberea* MULSANT, 1926 蘋果天牛屬

Oberea MULSANT, 1839, Col. France Long., p. 192 (Type-species: *Cerambyx oculatus* LINNÉ, 1758).

Isosceles NEWMAN, 1840, Entomologist, 1: 318.

Amaurostoma MÜLLER, 1906, Wien. Ent. Ztg., 25: 222 (Type-species: *Cerambyx erythrocephalus* SCHRANK, 1778).

Oberea binotaticollis PIC, 1915 短頸蘋果天牛

PIC, 1915, Melang Exot.-Ent., 13: 13 (Taiwan).

Distr.: Taiwan.

O. formosana PIC, 1911 蓬萊蘋果天牛

PIC, 1911, Mat. Long., 8, 1: 20 (Taiwan).

posticata: KANO (nec GAHAN), 1928, Trans. Nat. Hist. Soc. Formosa, 18: 126
(Chikuto).

Distr.: Taiwan; S. China, Hainan, Korea.

O. fuscipennis (CHEVROLAT, 1852) 褐翅細蘋果天牛

CHEVROLAT, 1852, Revue Mag. Zool., (2)4: 419 (*Isosceles*) (China).

fulveola BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 390 (Japan).

Distr.: Taiwan; Japan, Korea, China, Hainan I.

O. griseopennis SCHWARZER, 1925 樟灰蘋果天牛

SCHWARZER, 1925, Ent. Blaett., 21: 154 (Kankau, Kosempo et Sokutsu).

Distr.: Taiwan.

O. nigriceps (WHITE, 1844) 黑腳蘋果天牛

WHITE, 1844, Ann. Mag. Nat. Hist., 14: 425 (*Saperda* (*Isosceles*)) (Hong Kong).

Distr.: Taiwan(?); S. China, Hainan I.

O. nigriventris BATES, 1873 黑腹細蘋果天牛

BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 390 (Japan).

Distr.: Taiwan; Japan, China, Hainan I., Indo-China, Malaysia.

O. reductesignata PIC, 1916 黑尾蘋果天牛

PIC, 1916, Mat. Long., 10, 1: 17 (Taiwan).

nigriceps; KANO (nec WHITE), 1933, Kontyû, 6: 269 (Hori).

Distr.: Taiwan.

- O. taiwana* MATSUSHITA, 1933 臺灣黑腹蘋果天牛
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 423, pl. 5, f. 11 (Taihorin).
 Distr.: Taiwan.
- O. taihokensis* BREUNING, 1961 臺北蘋果天牛
 BREUNING, 1961, Frust. Ent., 5(4): 168 (Taihoku).
 Distr.: Taiwan.
- O. walkeri* GAHAN, 1894 華克氏蘋果天牛
 GAHAN, 1894, Trans. Ent. Soc. London, p. 489 (Hong Kong).
latipennis GRESSITT, 1939, Lingnan Sci. J., 18: 99, 104, pl. 3, f. 9 (*O. latipennis* forma) (China).
 Distr.: Taiwan; China.

Genus *Linda* THOMSON, 1864 長腳蘋果天牛屬

- Linda* THOMSON, 1964, Syst. Ceramb., pp. 122, 400 (Type-species: *Amphionycha femorata* CHEVROLAT, 1852).
Miocris FAIRMAIRE, 1902, Bull. Soc. Ent. France, p. 245 (Type-species: *M. nigroscutatus* FAIRMAIRE, 1902).
- Linda annulicornis* MATSUSHITA, 1933 環紋黃胸蘋果天牛
 MATSUSHITA, 1933, J. Fac. Agr. Hokkaido Univ., 34: 424, pl. 5, f. 14 (Kagi).
 Distr.: Taiwan; S. & SE. China.
- L. femorata* (CHEVROLAT, 1852) 粗腿蘋果天牛
 CHEVROLAT, 1852, Revue Zool., (2)4: 418 (*Amphionycha*) (China).
 Distr.: Taiwan; China.
- L. fraterna* (CHEVROLAT, 1852) 長角蘋果天牛
 CHEVROLAT, 1852, Revue Zool., (2)4: 419 (*Amphionycha*) (China).
 Distr.: Taiwan(?); China.
- L. signaticornis* SCHWARZER, 1925 紅胸黑翅蘋果天牛
 SCHWARZER, 1925, Ent. Blaett., 21: 154 (Kosempo, Kankau et Sokutsu).
 Distr.: Taiwan.
- L. subannulata* BREUNING 1966 摳環紋蘋果天牛
 BREUNING, 1966, Bull. Inst. Roy. Sci. Nat. Belg., 42(21): 19 (n. n. for *L. subannulicornis* BREUNING et OHBAYASHI, 1964).
 Distr.: Taiwan.

Genus *Epiglenea* BATES, 1884 四黃星天牛屬

- Epiglenea* BATES, 1884, J. Linn. Soc. London, 18: 259 (Type-species: *E. comes* BATES, 1884).
- Epiglenea comes* BATES, 1884
 BATES, 1884, J. Linn. Soc. London, 18: 259 (Japan).
 Distr.: Taiwan; Japan, China, Korea.
- subsp. formosana* (SCHWARZER, 1925) 蓬萊四星天牛
 SCHWARZER, 1925, Ent. Blaett., 21: 153 (*Daphisia formosana*) (Taihorin, Kosempo et Pilam).
 Distr.: Taiwan.

Genus *Phytoecia* MULSANT, 1839 菊花天牛屬

Phytoecia MULSANT, 1839, Col. France Long., p. 199 (Type-species: *Cerambyx cylindricus* LINNÉ, 1758).

Phytoecia rufiventris GAUTIER, 1870 菊花天牛

GAUTIER, 1870, Petites. Nouv. Ent., 1: 104 (E. Siberia).

punctigera BLESSIG, 1873, Hor. Soc. Ent. Ross., 9: 226(66) (E. Siberia).

ventralis BATES, 1873, Ann. Mag. Nat. Hist., (4)12: 388 (China).

abdominalis CHEVROLAT, 1882, Ann. Soc. Ent. France, (6)2: 62.

Distr.: Taiwan; Japan, China, Korea, E. Siberia.

誌謝

本文承蒙國家科發展委員會計劃經費補助 (NSC 75-0409-B002-47)，得以完成，特此誌謝。

臺灣產天牛科目錄 (II)

粗天牛亞科

中村慎吾 楠原 寛 齊藤明子

朱耀沂 林 匡夫

本篇中列出臺灣產天牛科、粗天牛亞科之種類共 311 種。

A. sublusca (THOMSON, 1857)

THOMSON, 1857, Arch. Ent., 1: 293 (*Monochamus*) (Singapore).

curialis PASCOE, 1858, Trans. Ent. Soc. London, (2)4: 246 (*Haplohammus*) (N. China).

Distr.: Taiwan; Japan (Ryukyu Is.), China, Hainan I., Indo-China, Singapore.
subsp. *maculihumera* (MATSUSHITA, 1933) 肩紋絨毛天牛

MATSUSHITA, 1933, J. Fac. Agric. Hokkaido Imp. Univ., 34(2): 328 (*Dihammus*) (Horisha).

Distr.: Taiwan.

A. ikedai MITONO, 1943 池田氏絨毛天牛

MITONO, 1943, Trans. Nat. Hist. Soc. Formosa, 33: 581 (*Dihammus*) (Sozan).

vitalisi GRESSITT (nec PIC), 1951, Longicornia, 2: 404 (*Dihammus*).

Distr.: Taiwan.

A. yokoyamai HAYASHI, 1971 橫山氏絨毛天牛

HAYASHI, 1971, Ent. Rev. Japan, 23(1): 15 (Lanyu I.).

Distr.: Taiwan (Lanyu I.).

Genus *Mimorsidis* BREUNING, 1938 紅翅天牛屬

Mimorsidis BREUNING, 1938, Novit. Ent., 8: 33 (Type-species: *M. lemóulti* BREUNING, 1938).

Mimorsidis taiwanensis HAYASHI, 1974 臺灣深條天牛

HAYASHI, 1974, Ent. Rev. Japan, 27(1/2): 47 (Kenting Park).

Distr.: Taiwan.

Tribe *Batocerini* 白條天牛族Genus *Batocera* CASTELNAU, 1840 白條天牛屬

Batocera CASTELNAU, 1840, Hist. Nat. Col., 2: 470 (Type-species: *Cerambyx rubus* LINNÉ, 1758).

Batocera davidis DEYROLLE, 1878 大衛大天牛

DEYROLLE, 1878, Ann. Soc. Ent. France, (5)8: 131 (E. China).

Dist.: Taiwan; S. China, Tonkin.

B. lineolata CHEVROLAT, 1852 白條天牛

CHEVROLAT, 1852, Revue Mag. Zool., (2)4: 417 (China).

chinensis THOMSON, 1857, Arc. Nat., 1: 170 (N. China).

catenata VOLLENHOVEN, 1871, Tijdschr. Ent., p. 215.

flachi SCHWARZER, 1914, Ent. Mitt., 3: 280 (China).

hauseri SCHWARZER, 1914, Ent. Mitt., 3: 280 (China).

Distr.: Taiwan; Japan, China, Korea, Quelpart I.

B. punctata SCHWARZER, 1925 臺灣大天牛

SCHWARZER, 1925, Ent. Blaett., 21: 60 (Taiwan).

Distr.: Taiwan.