

A SHORT REVIEW ON THE GENUS *PARACLYTUS* BATES, 1884 (COL.: CERAMBYCIDAE: CERAMBYCINAE)

Hüseyin Özdi̇kmen*

* Gazi Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, 06500 Ankara / Türkiye. E-mail: ozdikmen@gazi.edu.tr

[Özdikmen, H. 2009. A short review on the genus *Paraclytus* Bates, 1884 (Coleoptera: Cerambycidae: Cerambycinae). Munis Entomology & Zoology 4 (2): 327-332]

ABSTRACT: All taxa of the genus *Paraclytus* Bates, 1884 in the whole world are evaluated. The genus is also discussed in detail. The main aim of this catalogic work is to clarify current status of the genus in the world.

KEY WORDS: *Paraclytus*, Cerambycinae, Anaglyptini, Cerambycidae.

Subfamily CERAMBYCINAE Latreille, 1802

Tribe ANAGLYPTINI Lacordaire, 1869

= *Anaglypti* LeConte, 1873

The tribe includes currently at least 12 genera as *Anaglyptus* Mulsant, 1839; *Aphysotes* Bates, 1885; *Clytoderus* Linsley, 1935; *Cyrtophorus* LeConte, 1850; *Diphryrama* Bates, 1872; *Hirticlytus* K.Ohbayashi, 1960; *Microclytus* LeConte, 1873; *Miroclytus* Aurivillius, 1910; *Oligoenoplus* Chevrolat, 1863; *Paraclytus* Bates, 1884; *Pempteuryys* Bates, 1885 and *Tilloclytus* Bates, 1885. The genus name *Miroclytus* Aurivillius, 1910 has been given by some authors (e.g. Bisby et al., 2008) as *Microclytus* erroneously. However, the genus *Microclytus* LeConte, 1873 is a neotropical genus and *Miroclytus* Aurivillius, 1910 (original spelling) that was also given by Aurivillius (1912) is a Madagascan genus. The monotypic genus *Microclytus* Aurivillius, 1910 has only the species *Microclytus brunneipennis* Aurivillius, 1910 as the type. But *Microclytus* LeConte, 1873 (type species *Clytus gazellula* Haldeman, 1847) includes two species as *Microclytus compressicollis* (Laporte & Gory, 1835) and *Microclytus gazellula* (Haldeman, 1847) from NE North America: Canada and United States (Monné & Hovore, 2005). So, these genera are not homonyms.

Genus *PARACLYTUS* Bates, 1884

Type species: *Paraclytus excultus* Bates, 1884

Body length is approximately between 10 and 20 mm.

Shortly, the genus *Paraclytus* is characterized by a comparatively long 4th antennal segment (barely shorter than 3rd), convex and compactly punctate pronotum, narrow and long episternum of metathorax, short hind tarsi, and other characters.

Adults differ from other genera of the tribe in relatively long antennae, and structure and pattern of elytra. Pronotum is barely oblong, broadly rounded laterally. Scutellum is generally elongate, triangular, pointed anteriorly. Elytra

with parallel sides, convex, depressed on inner side humeri and on suture behind scutellum.

Larval and pupal developments are in deciduous trees (e.g. *Fagus*, *Crateagus*, *Quercus*, *Morus*, *Sorbus*, *Alnus* etc.) in forests.

The main aim of this catalogic work is to clarify the current status of the relict genus in the world. As commonly accepted that this chiefly Eastern Palaearctic genus *Paraclytus* Bates, 1884 [except the western Palaearctic species *P. luteofasciatus* (Pic, 1905); *P. raddei* (Ganglbauer, 1881); *P. reitteri* (Ganglbauer, 1881) and *P. sexguttatus* (Adams, 1817) which have SW Asiatic chorotype] is represented by 9 species (without subspecies) in the whole world. According to the data on range of these species, distributionally this genus is limited by Bulgaria and Greece in the West (Bulgaria, Greece, Turkey, Iran, Azerbaijan, Armenia) and Japan and Sakhalin Island in the East (China, Japan, Sakhalin Island) (e.g. Kraatz, 1864; Bates, 1884; Winkler, 1924-1932; Villiers, 1967; Demelt, 1972 and 1982; Cherepanov, 1990; Niisato, 2001; Miroshnikov, 2001; Sama, 2002; Holzschuh, 2003; Özdişmen & Demir, 2006; Özdişmen, 2007; Georgiev, 2008; Danilevsky, 2009a,b,c,d).

The most widely distributed species is *Paraclytus sexguttatus* (Adams, 1817) that is given as *P. sexmaculatus* by some authors (e.g. Sama, 2002). *P. raddei* (Ganglbauer, 1881) and *P. reitteri* (Ganglbauer, 1881) are following it. *Paraclytus excultus* Bates, 1884 that is the type species of the genus is only distributed in Far East (Sakhalin Island and Japan). The remaining are endemic taxa. Although *Paraclytus luteofasciatus* (Pic, 1905) is endemic to Greece, 4 species as *Paraclytus apicicornis* (Gressitt, 1937); *Paraclytus emili* Holzschuh, 2003; *Paraclytus primus* Holzschuh, 1992 and *Paraclytus shaanxiensis* Holzschuh, 2003 are endemic to China. The genus is represented only by two species as *Paraclytus luteofasciatus* (Pic, 1905) and *Paraclytus sexguttatus* (Adams, 1817) in Europe. The latter was recorded as a first record for Europe by Georgiev & Stojanova (2003). Very recently, a work on distribution, biology and ecology of *Paraclytus sexguttatus* (Adams, 1817) has been published by Georgiev (2008). *Paraclytus sexguttatus* (Adams, 1817) is only species of this genus in Turkey.

In addition to this, Casey (1912) gave three species as *Paraclytus brevitarsis*; *Paraclytus crucialis* and *Paraclytus lanifer* in the genus *Paraclytus* from California (USA), but all taxa are synonyms of the species *Triodoclytus lanifer* (LeConte, 1873) in the tribe Clytini. Aurivillius (1912) mentioned the species *Oligoenoplus rosti* (Pic, 1911) in the genus *Paraclytus* mistakenly. Also Winkler (1924-1932) placed the species *Anaglyptus thibetanus* Pic, 1914 in the genus *Paraclytus*.

All taxa of this genus in the world are presented as follows:

apicicornis Gressitt, 1937

Original combination: *Aglaophis apicicornis* Gressit, 1937

It is Eastern Palaearctic species.

DISTRIBUTION: China
CHOROTYPE: Chineese endemic

emili Holzschuh, 2003

Original combination: *Paraclytus emili* Holzschuh, 2003

It is Eastern Palaearctic species.

DISTRIBUTION: China
CHOROTYPE: Chineese endemic

excultus Bates, 1884

Original combination: *Paraclytus excultus* Bates, 1884

It is Eastern Palaearctic species.

DISTRIBUTION: Sakhalin Island, Japan
CHOROTYPE: Eastern Palaearctic

luteofasciatus Pic, 1905

Original combination: *Anaglyptus luteofasciatus* Pic, 1905

Other names: *moreanus* Demelt, 1972

It is Western Palaearctic species.

DISTRIBUTION: Greece
CHOROTYPE: Greek endemic

primus Holzschuh, 1992

Original combination: *Paraclytus primus* Holzschuh, 1992

It is Eastern Palaearctic species.

DISTRIBUTION: China
CHOROTYPE: Chineese endemic

raddei Ganglbauer, 1881

Original combination: *Anaglyptus raddei* Ganglbauer, 1881

It is Western Palaearctic species.

DISTRIBUTION: Caucasus (Azerbaijan), Iran
CHOROTYPE: SW-Asiatic (Irano-Caucasian)

reitteri Ganglbauer, 1881

Original combination: *Anaglyptus reitteri* Ganglbauer, 1881

It is Western Palaearctic species.

DISTRIBUTION: Caucasus (Azerbaijan), Iran

CHOROTYPE: SW-Asiatic (Irano-Caucasian)

sexguttatus Adams, 1817

Original combination: *Clytus sexguttatus* Adams, 1817

Other names: *caucasicus* Motschulsky, 1839; *bruckii* Kraatz, 1864; *disjunctus* Pic, 1909

It is Western Palaearctic species.

MATERIAL EXAMINED: Bolu province: Near Abant Lake, 2007, 1 specimen.

RECORDS IN TURKEY: Northern Turkey (Danilevsky & Miroshnikov, 1985); Artvin prov.: Saçinka (Alkan, 2000); Kırklareli prov.: Demirköy, Bolu prov.: Abant lake (Akçaalan) (Özdikmen & Demir, 2006).

DISTRIBUTION: Europe (Bulgaria), Caucasus (Georgia, Armenia, Azerbaijan), Turkey, Iran

CHOROTYPE: SW-Asiatic (Anatolo-Caucasian + Irano-Caucasian + Irano-Anatolian) + Turano-Mediterranean (Balkano-Anatolian)

shaanxiensis Holzschuh, 2003

Original combination: *Paraclytus shaanxiensis* Holzschuh, 2003

It is Eastern Palaearctic species.

DISTRIBUTION: China

CHOROTYPE: Chinese endemic

An important output: As this work also revealed that apparently the distributional areas of almost all species of this genus are much narrower than the past. So, these relict and rare forest species of the genus *Paraclytus* must be taken under protection for the future.

LITERATURE CITED

Aurivillius, C. 1910. Neue oder wenig bekannt Coleoptera Longicornia. Arkiv. För Zoologi 7 (3): 17-18.

Aurivillius, C. 1912. Coleopterorum Catalogus, pars 39 [vol. 22], Cerambycidae: Cerambycinae. Berlin. W. Junk & S. Schenckling. 574 pp.

Bates, H. W. 1884. Longicorn beetles of Japan. Additions, chiefly from the later collections of Mr. George Lewis; and notes on the synonymy, distribution, and habits on the previously known species. The Journal of the Linnean Society of London. Zoology 18: 205-262.

Bisby, F. A., Roskov, Y. R., Orrell, T. M., Nicolson, D., Paglinawan, L. E., Bailly, N., Kirk, P. M., Bourgoin, T. & van Hertum, J. eds. 2008. Species 2000 & ITIS Catalogue of Life: 2008 Annual Checklist. Digital resource at www.catalogueoflife.org/annual-checklist/2008/. Species 2000: Reading, U.K.

Casey, T. L. 1912. III - Studies in the Longicornia of North America. Memoirs on the Coleoptera, Lancaster 3: 215-376.

Danilevsky, M. L. 2009a. A check-list of Longicorn Beetles (Coleoptera, Cerambycoidea) of Europe. Available from: <http://www.cerambycidae.net/> (Updated 29.01.2009).

Danilevsky, M. L. 2009b. Systematic list of Longicorn Beetles (Cerambycoidea) of the territory of the former USSR. Available from: <http://www.cerambycidae.net/> (Updated 29.01.2009).

Danilevsky, M. L. 2009c. A check-list of Longicorn Beetles (Cerambycoidea) of Russia. Available from: <http://www.cerambycidae.net/> (Updated 29.01.2009).

Danilevsky, M. L. 2009d. A check-list of Longicorn Beetles (Cerambycoidea) of Russia, Ukraine, Moldova, Transcaucasia, Central Asia, Kazakhstan and Mongolia. Available from: <http://www.cerambycidae.net/> (Updated 29.01.2009).

Demelt, C. 1972. Beschreibung des Männchens von *Anaglyptus moreanus* Dem. Entomologische Blätter für Biologie und Systematik der Käfer 68 (3): 175-176.

Demelt, C. 1982. Nachtrag zur Kenntnis der Cerambycidenfauna Griechenlands. Entomofauna Zeitschrift für Entomologie, Linz 92 (17): 240-242.

Georgiev, G. & Stojanova, A. 2003. New and rare Longhorn Beetles (Coleoptera: Cerambycidae) in the Strandzha Mountain, Bulgaria. Acta Zoologica Bulgarica 55 (2): 105-109.

Georgiev, G. 2008. Notes on distribution, biology and ecology of *Paraclytus sexguttatus* (Coleoptera, Cerambycidae). Fragmenta entomologica, Roma 40 (1): 115-117.

Holzschuh, C. 2003. Beschreibung von 72 neuen Bockkäfern aus Asien, vorwiegend aus China, Indien, Laos und Thailand (Coleoptera, Cerambycidae). Entomologica Basiliensa, 25: 147-241.

Kraatz, G. 1864. *Clytus bruckii* n. sp. Berliner Entomologische Zeitschrift 8 (3-4): 389-390.

LeConte, J. L. 1873. Classification of the Coleoptera of North America. Prepared for the Smithsonian Institution. Part II. Smithsonian Miscellaneous Collections. Washington, D. C. 11 (265): 279-348.

Mirosnikov, A. I. 2001. New data on the Longicorn Beetles (Coleoptera, Cerambycidae) of the Talysh Mountains. Bulletin de la Société des Naturalistes de Moscou 106 (1): 49-50.

Monné, M. A. & Hovore, F. T. 2005. Checklist of the Cerambycidae, of the Western Hemisphere. Electronic Version 2005 (updated through 01 January 2006). Available from http://www.cerambycids.com/checklist/Monne&Hovore_2005.pdf

Niisato, T. 2001. Collecting Records of Cerambycid Beetles (Coleoptera, Cerambycidae) from the Kuril Archipelago (3) Elytra, Tokyo 29 (1): 7-15.

Özdikmen, H. 2007. The Longicorn Beetles of Turkey (Coleoptera: Cerambycidae) Part I - Black Sea Region. *Munis Entomology & Zoology* 2 (2): 179-422.

Özdikmen, H. & Demir, H. 2006. Notes on longicorn beetles fauna of Turkey (Coleoptera: Cerambycidae). *Munis Entomology & Zoology*, 1 (1): 157-166.

Sama, G. 2002. Atlas of the Cerambycidae of Europe and the Mediterranean Area. Volume 1: Northern, Western, Central and Eastern Europe. British Isles and Continental Europe from France (excl. Corsica) to Scandinavia and Urals. Vít Kabourek, Zlín, 173 pp.

Villiers, A. 1967. Contribution à la faune de l'Iran. I. - Coléoptères Cerambycidae. *Annales de la Société Entomologique de France* (N.S.) 3 (2): 327-379.

Winkler, A. 1924-1932. Catalogus Coleopterorum regionis palaearcticae. Verlag von Albert Winkler, 1135-1226.