

ART. 28.—*An Index of New Zealand Beetles.*

By G. V. HUDSON, F.E.S., F.N.Z.Inst.

[Read before the Wellington Philosophical Society, 7th December, 1921; received by Editor, 20th October, 1921; issued separately, 30th April, 1923.]

I HAVE undertaken the preparation of the following index, which is outside the scope of my usual entomological work, under the conviction that such an index is urgently required, and is not otherwise likely to be compiled. It is perhaps desirable for me to state that the list is simply a systematic index. It is neither a revision nor a synonymic list. No doubt a properly revised synonymic list of our Coleoptera is a great desideratum at the present time, but the production of such a list would require a much fuller knowledge of our beetles than I can claim to possess, and must, so far as I can see, be left to some future worker. In the meantime the present list may be regarded as a preliminary step. It will assist collectors in arranging their collections, and will enable them to refer to a description of any species they may desire to consult.

The list of Coleoptera contained in Captain Hutton's invaluable *Index Faunae Novae Zealandiae* is now very much out of date, over 1,400 species having been described by the late Major Broun since that work was published in 1904. Captain Hutton's list was prepared by Major Broun himself, and I have assumed that it is a complete index up to the end of the year 1903. In compiling the following list I have consulted all available publications issued subsequent to that date, and in this way have endeavoured to attain a measure of completeness. In the course of my work I have found that certain species mentioned in Major Broun's *Manual of New Zealand Coleoptera* do not appear in Captain Hutton's list; and, unless these species are mentioned in publications later than Hutton's index, I have not inserted them, as I assume that Major Broun must have regarded them as synonyms and intentionally omitted them.

The scheme of the classification followed in my index is that in general use in England at the present time. It is regarded as a more natural and satisfactory system than that contained in the *Cambridge Natural History*, which formed the basis of Captain Hutton's list. For assistance and advice in connection with this part of my work I am much indebted to my kind friend Commander J. J. Walker. Within each family I have arranged the genera, as far as possible, in accordance with the views expressed in Major Broun's publications.

As regards the question of synonymy, it should be stated at the outset that Major Broun is not responsible for the creation of anything like the number of synonyms that some of his critics have assumed. The coleopterous fauna of New Zealand is undoubtedly a very rich one, and, like the Lepidoptera, contains a very large proportion of difficult forms whose slight and elusive differences are extremely difficult to appreciate. The whole subject of synonymy is, however, very slightly touched upon in Major Broun's works, and it certainly appears probable that some of the descriptions contained in the very early publications may relate to insects which have been renamed and redescribed in later publications. For example, *Uloma tenebrionides* White and *Uloma laevicostata* Blanchard

may be names referring to the same species, and this may also apply to *Teramocerus barbicornis* Fabricius and *Teramocerus cylindricornis* Fabricius. These and similar questions can only be settled by reference to the original types. Again, Major Broun repeatedly states he has not seen the types of certain species described by Dr. Sharp and other authors, and it would appear likely that a certain amount of synonymy may have resulted in at least some of these cases. Furthermore, it is noteworthy that Major Broun has described a great number of species from single specimens, and in these cases the question of possible intermediate forms connecting some of the species cannot have been fully taken into account.

I am much indebted to the Hon. G. M. Thomson, M.L.C., for a list of the introduced beetles. These have, as far as possible, been inserted in systematic order. They are in every case distinguished by an asterisk placed before the name of each species.

Beetles found in the Kermadec Islands which do not also occur in New Zealand are not included in the index.

In referring to so many scattered publications, extending over a period of nearly twenty years, it is inevitable that some errors and omissions have occurred, and in this connection I must ask for the kind indulgence of all those who may use the index.

ABBREVIATIONS.

- A. *Annals and Magazine of Natural History.*
- B. *Bulletin, New Zealand Institute.*
- C. *Manual of New Zealand Coleoptera, 1880-93.*
- E. *Transactions of the Entomological Society of London.*
- S. *Subantarctic Islands of New Zealand.*
- T. *Transactions of the New Zealand Institute.*

Order COLEOPTERA.

Group GEODEPHAGA.

Family CICINDELIDAE.

- Cicindela* Linnaeus, 1735.
 - tuberculata* Fabricius. C. 2.
 - buttoni* Broun. C. 2.
 - laticeincta* White. C. 3.
 - wakefieldi* Bates. C. 3.
 - douei* Chenu. C. 3.
 - parryi* White. C. 4.
 - dunedinensis* Castelnau. C. 4.
 - jeredaijii* Bates. C. 4.
 - austromontana* Bates. C. 5.
 - perhispida* Broun. C. 4.
 - campbelli* Broun. C. 817.
 - helmsi* Sharp. C. 975.
 - incognita* Horn. *Deutsche Ent. Zeit.*, 1892, 82.
 - novaseelandica* Horn. *Deutsche Ent. Zeit.*, 1892, 83.
 - circumpictoides* Horn. *Deutsche Ent. Zeit.*, 1900, 207.
 - warioruensis* Broun. B. 1, iii, 146.
 - halli* Broun. B. 1, v, 350.
 - hamiltoni* Broun. B. 1, vii, 594.
 - ezonata* Broun. B. 1, vii, 594.

Family CARABIDAE.

Subfamily MIGADOPINAE.

- Amarotypus* Bates, 1872.
- edwardsii* Bates. C. 6.

Subfamily SCARILINAE.

- Clivina* Latreille, 1810.
- rugithorax* Putzeys. C. 7.

Subfamily CNEMACANTHINAE.

- Mecodema* Blanchard, 1853.
- sculpturatum* Blanchard. C. 8.
- howitti* Castelnau. C. 8.
- rectolineatum* Castelnau. C. 8.
- impressum* Castelnau. C. 9.
- lucidum* Castelnau. C. 9.
- crenicolle* Castelnau. C. 9.
- simplex* Castelnau. C. 9.
- alternans* Castelnau. C. 10.
- crenaticolle* Redtenbacher. C. 10.
- spinifer* Broun. C. 10.
- fulgidum* Broun. C. 653.
- constrictum* Broun. C. 653.

Group GEODEPHAGA—continued.

Mecodema—continued.

- rugicollis* Broun. C. 745.
venator Broun. C. 817.
litoreum Broun. C. 875.
ducale Sharp. C. 976.
rugiceps Sharp. C. 976.
metallicum Sharp. C. 977.
suteri Broun. C. 977.
scitulum Broun. A. 6, xiv, 302.
lineatum Broun. A. 6, xiv, 303.
punctifer Broun. C. 745.
aeneoniger Broun. C. 818.
bullatum Lewis. T. xxxiv, 202.
infimatum Lewis. T. xxxiv, 202.
laeviceps Broun. A. 7, xiv, 41.
striatum Broun. A. 7, xiv, 42.
walkeri Broun. A. 7, xiv, 43.
costellatum Broun. A. 7, xi, 451.
intricatum Broun. A. 7, xi, 451.
nitidum Broun. A. 7, xi, 452.
variolum Broun. A. 7, xi, 453.
costipenne Broun. B. 1, ii, 82.
angustulum Broun. B. 1, ii, 82.
dissimum Broun. B. 1, ii, 83.
acuductum Broun. A. 8, ii, 335.
cognatum Broun. A. 8, ii, 336.
lewisi Broun. A. 8, ii, 337.
seriatum Broun. A. 8, ii, 338.
attenuatum Broun. A. 8, ii, 339.
puncticolle Broun. B. 1, iii, 146.
hallsi Broun. B. 1, iv, 270.
huttense Broun. B. 1, iv, 270.
persculptum Broun. B. 1, iv, 271.
ambiguum Broun. B. 1, iv, 272.
laterale Broun. B. 1, v, 350.
chiltoni Broun. B. 1, v, 351.
politatum Broun. B. 1, v, 352.
rubripes Broun. B. 1, v, 353.
erraticum Broun. B. 1, v, 353.
gratum Broun. B. 1, v, 354.
latulum Broun. B. 1, v, 355.
affinum Broun. B. 1, v, 355.
indiscretum Broun. B. 1, v, 356.
trailii Broun. B. 1, v, 356.
mutabile Broun. B. 1, v, 357.
gordonense Broun. B. 1, v, 357.
o'connori Broun. T. xliv, 382.
bryobium Broun. T. xliv, 383.
laevicollis Broun. T. xliv, 384.
quoinense Broun. T. xliv, 384.
arcuatum Broun. T. xliv, 385.
hudsoni Broun. S. 83.
femorale Broun. B. 1, vii, 595.
veratrum Broun. B. 1, vii, 596.
punctellum Broun. B. 1, vii, 596.
insulare Broun. B. 1, vii, 597.
validum Broun. B. 1, viii, 670.
occiputale Broun. B. 1, viii, 670.
cassense Broun. B. 1, viii, 671.
antennale Broun. B. 1, viii, 672.
longicollis Broun. B. 1, viii, 672.
philotti Broun. B. 1, viii, 672.
ventriculum Broun. B. 1, viii, 673.
subaeneum Broun. B. 1, viii, 674.

- Metaglymma* Bates, 1867.
tibiale Castelnau. C. 11.
monilifer Bates. C. 11.
punctatum Putzeys. C. 11.
morio Castelnau. C. 11.
elongatum Castelnau. C. 12.
aberrans Putzeys. C. 12.
dyschirioidea Castelnau. C. 12.
oblongum Broun. C. 746.
asperum Broun. C. 978.
junctum Broun. C. 979.
sulcatum Sharp. C. 979.
tersatum Broun. C. 980.
rugipenne Broun. C. 1321.
thoracicum Broun. C. 1322.
modicum Broun. A. 6, xiv, 305.
oregoide Broun. A. 6, xiv, 305.
rufipes Broun. C. 876.
rugiceps Broun. A. 7, xi, 454.
calcaratum Broun. A. 7, xi, 455.
doulli Broun. A. 7, xv, 543.
minor Broun. A. 7, xv, 543.
ovicolle Broun. B. 1, iii, 147.
curvipes Broun. B. 1, iv, 273.
Oregus Putzeys, 1868.
aereus White. C. 13.
inaequalis Castelnau. C. 13.
Brullea Castelnau, 1867.
antarctica Castelnau. C. 14.
Diglymma Sharp, 1886.
ovipenne Sharp. C. 981.
dubium Sharp. C. 981.
punctipenne Broun. C. 982.
nigripes Broun. C. 982.
oblusum Broun. C. 819.
clivinoides Castelnau. C. 12.
tarsalis Broun. A. 8, ii, 340.
marginale Broun. B. 1, iii, 148.
basale Broun. B. 1, v, 358.
thoracicum Broun. B. 1, v, 359.
castigatum Broun. S. 84.
Snofru Broun, 1908.
aemulator Broun. A. 8, ii, 341.
Acallistus Sharp, 1886.
simplex Sharp. C. 983.

Subfamily LICININAE.

- Rembus* Latreille, 1822.
zeelandicus Redtenbacher. C. 15.
Physolaesthus Chaudoir, 1850.
insularis Bates. C. 15.
limbatus Bates. C. 17.
Dichrochile Guérin, 1846.
subopaca Bates. C. 16.
blanipes Broun. B. 1, v, 360.
aterrima Bates. C. 16.
ovicollis Motschulsky. C. 16.
maura Broun. C. 18.
nitida Broun. C. 746.
cinctiger Broun. C. 747.
cephalotes Broun. A. 6, xiv, 306.
fabrei Guérin. Ann. Fr. 1846, Bull., 103.

Group GEODEPHAGA—continued.

Dichrochile—continued.

- anthracina* Broun. A. 6, xii, 161.
cordicolle Broun. A. 7, xi, 455
thoracica Broun. A. 8, ii, 345.
insignis Broun. B. 1, v, 359.
rugicollis Broun. B. 1, v, 360.

Subfamily ANCHOMENINAE.

- Calathus* Bonelli, 1809.
rubromarginatus Blanchard. S. 96.
Anchomenus Bonelli, 1809.
novae-zealandiae Fairmaire. Ann. Soc. Ent., 1843.
elevatus White. C. 21.
montivagus Broun. C. 22.
feredayi Bates. C. 23.
lawsoni Bates. C. 23.
submetallicus White. C. 24.
deplanatus White. C. 25.
coletonensis White. C. 25.
otagoensis Bates. C. 25.
edwardsii Bates. C. 26.
sulcifarius Broun. C. 27.
helmisi Sharp. C. 749.
haasti Broun. C. 747.
sandageri Broun. C. 748.
oreobius Broun. C. 876.
atratus Blanchard. Voy. Pole. Sud, iv, 21.
raptor Redtenbacher, Reise Novara, Coleop., 18.
marginellus Erichson. Archives, 1842, 130.
walkeri Broun. A. 7, xi, 456.
libitus Broun. B. 1, ii, 84.
macrocaelis Broun. A. 8, ii, 346.
xanthomelas Broun. A. 8, ii, 346.
intermedius Broun. A. 8, ii, 347.
integratus Broun. A. 8, ii, 348.
sophronitis Broun. A. 8, ii, 349.
chalhamensis Broun. T. xl, 147.
arnaudensis Broun. B. 1, vii, 598.
hallianus Broun. B. 1, vii, 598.
Ctenognathus Fairmaire, 1843.
latipennis Sharp. C. 984.
parabilis Broun. C. 20.
politulus Broun. C. 20.
batesi Broun. C. 20.
defoimipes Broun. C. 20.
cheesemani Broun. C. 20.
suborbithorax Broun. C. 20.
perrugithorax Broun. C. 20.
lucifugus Broun. C. 20.
adamsi Broun. C. 20.
pictonensis Sharp. C. 985.
bidens Chaudoir. Ann. Soc. Ent. Fr., 1875, 303.
munroi Broun. C. 984.
actochares Broun. A. 6, xiv, 307.
novae-zealandiae Fairmaire. Ann. Soc. Ent. Fr., 1843, 12.
neozelandicus Chaudoir. Ann. Soc. Ent. Fr., 1875, 294.

Ctenognathus—continued.

- crenatus* Chaudoir. Ann. Soc. Ent. Fr., 1875, 304.
cardiophorus Chaudoir. Ann. Soc. Ent. Fr., 1875, 305.
macropterus Chaudoir. Ann. Soc. Ent. Fr., 1875, 370.
littorellus Broun. A. 8, ii, 349.
simmonsi Broun. T. xliv, 386.
Pristancylus Blanchard, 1853.
brevis Blanchard. S. 95.
castaneus Blanchard. S. 95.
Pristonychus.
**terricola* Herbst.
Drimostoma Dejean, 1831.
antarctica Castelnau. C. 30.
- Subfamily FERONINAE.
- Prosopogmus* Chaudoir, 1865.
impressifrons Chaudoir. C. 30.
Trichosternus Chaudoir, 1865.
antarcticus Chaudoir. C. 31.
guernii Chaudoir. C. 32.
capito White. C. 32.
syurus Bates. C. 35.
aucklandicus Broun. C. 33.
diformipes Bates. C. 33.
lobipes Bates. C. 34.
temukensis Bates. C. 34.
rectangulus Chaudoir. C. 35.
planiusculus White. C. 36.
humeralis Broun. C. 750.
enys Broun. C. 752.
cephalotes Broun. C. 825.
alternus Broun. C. 877.
urquharti Broun. C. 877.
virens Broun. C. 937.
compressus Sharp. C. 987.
polychaetus Broun. C. 987.
smithii Broun. C. 1322.
hampdenensis Broun. C. 1323.
crassalis Broun. C. 1395.
walkeri Broun. A. 7, xi, 602.
akaroensis Broun. A. 7, xi, 603.
bucolicus Broun. A. 7, xi, 604.
rectalus Broun. C. 1146.
haplopus Broun. C. 1323.
hudsoni Broun. A. 7, xiv, 44.
dissentaneus Broun. B. 1, i, 6.
coelocephalus Broun. A. 8, ii, 405.
hammerensis Broun. A. 8, ii, 406.
ordinarius Broun. A. 8, ii, 407.
halli Broun. B. 1, iii, 149.
blandellus Broun. B. 1, iv, 274.
wallacei Broun. T. xliv, 390.
- Zeopoecilus* Sharp, 1886.
calcaratus Sharp. C. 989.
achilles Sharp. C. 990.
putus Broun. C. 750.
opulentus Broun. C. 820.
optandus Broun. A. 8, ii, 408

Group GEODEPHAGA—continued.

Pterostichus Bonelli, 1809.

- (Species like *Trichosternus*, but without prosternal setae.)
- fultoni* Broun. C. 751.
 - waikorensis* Broun. C. 821.
 - monticola* Broun. C. 821.
 - aeruginosus* Broun. C. 822.
 - asperatus* Broun. C. 822.
 - chloris* Broun. C. 823.
 - grassator* Broun. C. 823.
 - maiaei* Broun. B. I, v, 363.
 - amicus* Broun. C. 824.
 - fusulus* Broun. C. 824.
 - curvipes* Broun. C. 878.
 - optabilis* Broun. C. 913.
 - meritus* Broun. C. 914.
 - angulatus* Broun. C. 914.
 - melsusculus* Broun. C. 915.
 - agriolus* Broun. C. 915.
 - erraticus* Broun. C. 915.
 - decepitus* Broun. C. 916.
 - curlulus* Broun. C. 916.
 - suspicax* Broun. C. 917.
 - amplicollis* Broun. C. 918.
 - sandageri* Broun. C. 988.
 - haplopus* Broun. C. 1323.
 - vagans* Broun. C. 825.
 - dentiferus* Broun. C. 36.
 - kirkianus* Broun. A. 7, xi, 604.
 - memes* Broun. A. 7, xi, 605.
 - prasignis* Broun. A. 7, xi, 606.
 - setiventris* Broun. A. 7, xi, 606.
 - oneroaensis* Broun. A. 8, ii, 413.
 - flexipes* Broun. A. 8, ii, 413.
 - fossalis* Broun. B. I, iii, 149.
 - pascoi* Broun. B. I, iv, 275.
 - aciphylla* Broun. B. I, v, 361.
 - hamiltoni* Broun. T. xliv, 390.

(Thorax with two setae on each side.)

- prolixus* Broun. C. 35.
- integratus* Bates. C. 40.
- cavelli* Broun. C. 991.
- procerus* Broun. C. 827.
- ovatellus* Chaudoir. C. 39.
- perbonus* Broun. A. 8, ii, 418.
- placidulus* Broun. C. 658.
- impiger* Broun. C. 879.
- mordax* Broun. C. 938.
- longiformis* Sharp. C. 991.
- sternalis* Broun. C. 658.
- oscillator* Sharp. C. 995.
- lepidulus* Broun. A. 8, ii, 419.
- perfidiosus* Broun. C. 995.
- hunuensis* Broun. C. 996.
- maorinus* Bates. C. 41.
- disparalis* Broun. C. 1324.
- edax* Broun. C. 1326.
- arduus* Broun. C. 1395.
- scispennis* Broun. C. 1396.
- sharpianus* Broun. A. 6, xii, 163.
- chalmeri* Broun. A. 8, ii, 420.
- bullatus* Broun. B. I, iv, 275.
- sculpturalis* Broun. B. I, v, 363.

Pterostichus—continued.

- fenwicki* Broun. B. I, vii, 599.
- truncatus* Broun. B. I, viii, 674.
- (Thorax with three setae on each side.)
- elongellus* White.
- (Thorax with four setae on each side.)
- hispidulus* Broun. C. 40.
- cribralis* Broun. C. 752.
- myrmidon* Sharp. C. 993.
- hybridus* Broun. C. 826.
- fieldi* Broun. B. I, iv, 276.
- praecox* Broun. C. 827.
- burrowsi* Broun. B. I, iii, 150.
- brownianus* Sharp. C. 992.
- constrictellus* Sharp. C. 994.
- detractus* Broun. C. 996.
- subaeneus* Guérin. C. 38.
- oedicenemus* Bates. C. 39.
- pastorius* Broun. C. 994.
- inconstans* Broun. C. 999.
- philpotti* Broun. A. 8, ii, 418.
- egmontensis* Broun. C. 997.
- irregularis* Broun. C. 998.
- suteri* Broun. C. 1324.
- sinuellus* Broun. C. 1325.
- delator* Broun. C. 1397.
- rugifrons* Sharp. C. 1397.
- ithaginus* Broun. A. 6, xii, 162.
- obsoletus* Broun. A. 6, xii, 163.
- vexatus* Broun. A. 8, ii, 417.
- levisi* Broun. A. 6, xiv, 310.
- turgidiceps* Broun. A. 8, ii, 409.
- odontellus* Broun. A. 8, ii, 410.
- antennalis* Broun. A. 8, ii, 412.
- oxymelus* Broun. A. 8, ii, 415.
- sinuiventris* Broun. A. 8, ii, 416.
- (Thorax with five or more setae on each side.)
- catenulatus* Broun. C. 753.
- muuronatus* Broun. C. 826.
- insidiosus* Broun. C. 999.
- sculptipes* Broun. C. 1325.
- egregialis* Broun. B. I, v, 362.
- (Species of doubtful position.)
- thoracicus* Broun. C. 657.
- angustulus* Chaudoir. C. 38.
- sylvaticus* Chaudoir. C. 38.
- vigil* White. C. 41.
- politissima* White. C. 41.
- pella* White. C. 657.
- reflexum* Chaudoir. *Bull. Mosc.*, xxxviii, 74.
- convergadorsis* Tschitschérine. *Hör. En. Ross.*, 1891, 166.
- adoxus* Broun. A. 8, ii, 414.
- Alogus* Motschulsky, 1865.
- monachicus* Motschulsky. C. 44.
- Cerabilia* Castlenau, 1868.
- maori* Castlenau. C. 44.
- punctigera* Broun. C. 753.
- ruficornis* Broun. C. 754.

Group GEODEPHAGA—continued.

- Rhytidernus* Chaudoir, 1865.
rugifrons Broun. C. 41.
erythrognathus Broun. C. 986.
**puella* Chaudoir.
- Steropus* Megerle, 1821.
helmsi Sharp. C. 989.
- Argutor* Megerle, 1837.
erythropus Blanchard. C. 42.
piceus Blanchard. C. 43.
pantomelas Blanchard. *Voy. Pole Sud*, Zool., iv, 27.
calathoides Broun. C. 879.
- Subfamily ANISODACTYLINAE.
- Triplosarus* Bates, 1874.
fulvescens Bates. C. 45.
novae-zealandiae Castelnau. C. 659.
- Loxomerus* Chaudoir.
nebrioides Guérin. S. 92.
ambiguus Broun. S. 92.
fossilatus Broun. S. 93.
cilicollis Broun. S. 94.
huttoni Broun. S. 94.
- Lecanomerus* Chaudoir, 1850.
latimanus Bates. C. 45.
insignitus Broun. C. 47.
obesus Bates. C. 47.
fuliginosus Broun. C. 48.
fallax Broun. C. 48.
stenopus Broun. C. 880.
marginatus Sharp. C. 1000.
palipes Broun. A. 6, xiv, 379.
labialis Broun. B. 1, iii, 151.
incertus Broun. B. 1, iii, 151.
- Gaioxenus* Broun, 1910.
pilipalpis Broun. B. 1, i, 8.
- Taenarthrus* Broun, 1914.
philpotti Broun. B. 1, ii, 85.
- Zabronothus* Broun, 1893.
striatulus Broun. C. 1327.
rufipes Broun. C. 1328.
oblongus Broun. B. 1, i, 8.
major Broun. T. xliv, 393.
aphelus Broun. T. xliv, 394.
- Hypharpax* MacLeay, 1825.
antarcticus Castelnau. C. 49.
abstrusus Bates. C. 49.
- **australasiae* Dejean.
**australis* Dejean.
- Syllectus* Bates, 1878.
anomalus Bates. C. 52.
- Subfamily HARPALINAE.
- Harpalus* Latreille, 1804.
novae-zealandiae Castelnau. *Trans. R. Soc. Vict.*, 1867, 108.
- antarcticus* Castelnau. *Trans. R. Soc. Vict.*, 1867, 107.
- Allocinopus* Broun, 1903.
sculpticollis Broun. A. 7, xi, 608.
ocularius Broun. A. 8, ii, 344.
smithi Broun. T. xliv, 391.
castaneus Broun. T. xliv, 392.
latitarsis Broun. T. xliii, 95.
angustulus Broun. T. xliv, 392.
- Euthenarus* Bates, 1874.
brevicollis Bates. C. 53.
puncticollis Bates. C. 53.
cilicollis Broun. T. xxxiv, 176.
huttoni Broun. T. xxxiv, 177.
- Subfamily ZOLINAE.
- Zolus* Sharp, 1886.
helmsi Sharp. C. 1001.
atratus Broun. C. 1002.
femoralis Broun. A. 6, xiv, 311.
carinatus Broun. C. 754.
subopacus Broun. B. 1, iv, 277.
ocularius Broun. B. 1, v, 365.
labialis Broun. B. 1, vii, 599.
- Megadromus* Motschulsky, 1865.
viridilimbatus Motschulsky. *Bull. Mosc.*, iv, 251.
- Synteratus* Broun, 1909.
ovalis Broun. S. 85.
- Subfamily POGONINAE.
- Oöpterus* Guérin, 1841.
laevicollis Bates. C. 54.
patulus Broun. C. 655.
fulvipes Broun. C. 936.
sobrinus Broun. C. 936.
collaris Broun. C. 1002.
pallidipes Broun. C. 1003.
puncticeps Broun. C. 1398.
clivinotus Guérin. *Voy. Pole Sud*, Zool., iv, 43, and S. 86.
- plicaticollis* Blanchard. *Voy. Pole Sud*, Zool., iv, 44, and S. 87.
- latipennis* Broun. A. 7, xi, 609.
probus Broun. A. 7, xi, 610.
parvulus Broun. A. 7, xi, 610.
pygmeatus Broun. A. 7, xix, 56.
nigritulus Broun. A. 8, ii, 342.
frontalis Broun. A. 8, ii, 342.
sculpturalis Broun. A. 8, ii, 343.
basalis Broun. B. 1, iv, 273.
minor Broun. B. 1, v, 364.
latifossus Broun. B. 1, v, 364.
suavis Broun. B. 1, v, 365.
laevigatus Broun. B. 1, vi, 389.
tripunctatus Broun. S. 87.
marrineri Broun. S. 88.
tarsalis Broun. S. 89.
elongellus Broun. S. 89.
- Kenodactylus* Broun, 1909.
capito Broun. S. 91.
- Cyclothorax* MacLeay, 1873.
rotundicollis White. C. 54.
insularis Motschulsky. C. 29.
- Tarastethus* Sharp, 1883.
puncticollis Sharp. C. 1004.
laeviventris Sharp. C. 1004.
alpinus Broun. C. 1005.
debilis Sharp. C. 1005.
cinctus Broun. C. 1005.
scuticollis Bates. C. 28.
seriatoporus Bates. C. 28.
placens Broun. C. 28.
simplex Broun. A. 7, xi, 457.

Group GEODEPHAGA—continued.

Tarastethus—continued.

- laevicollis* Broun. A. 7, xi, 458.
pretiosus Broun. B. 1, i, 6.
strenuus Broun. A. 6, xiv, 308.
dubius Broun. A. 6, xiv, 309.
similans Broun. A. 6, xiv, 309.
southlandicus Broun. A. 8, ii, 350.
carbonarius Broun. A. 8, ii, 351.
amplipennis Broun. T. xliv, 386.
phylocharis Broun. T. xliv, 387.
lewisi Broun. T. xliv, 388.
cordipennis Broun. T. xliv, 388.
convexus Broun. B. 1, v, 366.
diversus Broun. B. 1, v, 366.
halli Broun. B. 1, v, 367.
fovealis Broun. B. 1, v, 367.
longulus Broun. B. 1, v, 368.
propinquus Broun. B. 1, v, 369.
optatus Broun. B. 1, v, 369.
robustus Broun. B. 1, vii, 600.
eplicatus Broun. B. 1, viii, 675.
fuscipes Broun. B. 1, viii, 675.
insularis Broun. B. 1, viii, 676.
Sympiestus Sharp, 1866.
syntheticus Sharp. C. 1006.
fallax Broun. C. 1007.
oculator Broun. C. 1398.
modestus Broun. A. 6, xiv, 311.
frontalis Broun. B. 1, v, 370.

Subfamily BEMBIDIINAE.

- Bembidium* Latreille, 1804.
charile Bates. C. 55.
moorinum Bates. C. 55.
tairuense Bates. C. 55.
rotundicolle Bates. C. 56.
parviceps Bates. C. 56.
anchonoderum Bates. C. 57.
eustichum Bates. C. 57.
callipeplum Bates. C. 57.
orbiferum Bates. C. 58.
chalcipes Bates. C. 58.
holitikense Bates. C. 59.
musae Broun. C. 755.
latiusculum Broun. C. 755.
nesophilum Broun. C. 828.
tantillum Broun. C. 828.
tekapoense Broun. C. 880.
attenuatum Broun. C. 881.
clevedonense Broun. C. 1007.
diaphanum Broun. C. 1007.
dehiscens Broun. C. 1009.
antipodum Broun. C. 1009.
tersatum Broun. C. 1010.
actuarium Broun. A. 7, xi, 611.
waikatoense Broun. B. 1, i, 9.
tinctulum Broun. B. 1, i, 9.
Cillenum Samouelle, 1819.
albescens Bates. C. 59.
chalmeri Broun. C. 881.
subcaecum Sharp. C. 1010.
alacris Broun. B. 1, vii, 601.
Tachys Shau, 1860.
antarcticus Bates. C. 60.
latipennis Sharp. C. 1011.

Tachys—continued.

- oreobius* Broun. C. 1399.
cavelli Broun. C. 1400.
coriaceus Broun. A. 8, ii, 421.
Anillus Duval, 1851.
pallidus Broun. C. 918.
phyllobius Broun. A. 6, xii, 164.
monticola Broun. B. 1, i, 10.
marginatus Broun. B. 1, iii, 152.
pinctigerus Broun. B. 1, iii, 153.

Subfamily LEBIINAE.

- Actenonyx* White, 1846.
bembidioides White. C. 61.
Lebia Latreille, 1804.
bembidioides Fairmaire. *Rev. Zool.* 1849, 32.
posticalis Guérin. *Voy. Coquille, Coleop.* 58.
Wakefieldia Broun, 1880.
vittata Broun. C. 62.
Agonochila Chaudoir, 1848.
**binotata* White. C. 63.
Demetrida White, 1846.
lineella White. C. 64.
nasuta White. C. 64.
picea Chaudoir. C. 65.
ater Broun. C. 66.
moesta Sharp. C. 66.
lateralis Broun. B. 1, i, 10.
sinuata Broun. B. 1, v, 370.
Pedalopia Castelnau, 1867.
novae-zelandiae Castelnau. C. 67.

Subfamily PERICALINAE.

- Scopodes* Erichson, 1842.
fossulatus Blanchard. C. 68.
elaphroides White. C. 68.
edwardsi Bates. C. 68.
versicolor Bates. C. 68.
prasinus Bates. C. 69.
multipunctatus Bates. C. 69.
laevigatus Bates. C. 70.
venustus Broun. C. 756.
planus Broun. C. 756.
tardus Broun. C. 757.
pustulatus Broun. C. 757.
laevistriatus Broun. C. 829.
bryophilus Broun. C. 882.
cognatus Broun. C. 882.
antennalis Broun. C. 882.
nigrinus Sharp. C. 1011.
basalis Broun. C. 1012.
speciosus Broun. C. 1012.
viridis Broun. A. 7, xi, 612.
nigripes Broun. B. 1, i, 11.
instabilis Broun. B. 1, v, 371.

Family RHYSODIDAE.

- Rhysodes* Dalman, 1823.
aterrimus Broun. C. 214.
eminens Broun. C. 215.
orbitosus Broun. C. 215.
proprius Broun. C. 216.
pensus Broun. C. 215.

Group HYDRADEPHAGA.

Family DYTISCIDAE.

- Homoeodytes* Sharp, 1882.
hookeri White. C. 75.
scutellaris Sharp. C. 1331.
Rhanthus Sharp, 1882.
pulverosus Stephens = *C. rufimanus* White. C. 74; C. 1333.
plantaris Sharp. C. 1333.
Lances Sharp, 1882.
lanceolatus Clark. C. 1335.

Family HYDROPORIDAE.

- Antiporus* Sharp, 1882.
uncifer Sharp. C. 1330.
wakefieldi Sharp. C. 71.
duplex Sharp. C. 72.

Antiporus—continued.

- strigosulus* Broun. C. 72.
nitidicornis Broun. C. 73.
Bidessus Sharp, 1882.
impressus Sharp. C. 1329.
plicatus Sharp. C. 1329.
huttoni Sharp. C. 1329.
Huxelhydrus Sharp, 1882.
syntheticus Sharp. *Trans. Roy. Dub. Soc.*, 1882.
virgatus Broun. C. 1014.
Copelatus Erichson. Sharp, 1882.
sharpi Broun. C. 1014 and 939.

Family GYRINIDAE.

- Gyrinus*, Linnaeus, 1733.
huttoni Pascoe. C. 76.

Group PHILHYDRIDA.

Family HYDROPHILIDAE.

- Philhydrus* Solier, 1834.
tritus Broun. C. 78.
variolosus Broun. C. 79.
abditus Sharp. C. 1018.
Berosus Leach, 1817.
mergus Broun. C. 883.
Phelerosus Sharp, 1884.
pallidipennis Sharp. C. 1023.
Saphydrus Sharp, 1884.
oblongus Broun. C. 81.
suffusus Sharp. C. 1015.
obesus Sharp. C. 1016.
antennatus Sharp. C. 1016.
longulus Sharp. C. 1016.
monticola Broun. C. 1401.
consonus Broun. B. 1, vi, 478.
coloris, Broun. B. 1, vi, 478.
Rygnodus White, 1846.
modestus White. C. 80.
pedinoides White. C. 80.
incertus Broun. C. 81.
optimus Broun. C. 82.
cyanus Broun. C. 659.
puncticeps Broun. C. 757.
femoratus Sharp. C. 1016.
tibialis Broun. C. 1017.
ovalis Sharp. C. 1017.
limbatis Broun. C. 1335.
aleniensis Broun. C. 1401.
nigripennis Broun. A. 7, xi, 612.
Tormus Sharp, 1884.
helmsi Sharp. C. 1019.
nitidus Broun. C. 1402.
Tormissus Broun, 1893.
marginatus Broun. C. 1021.
magnulus Broun. C. 1021.
Thomosis Broun, 1904.
†guanicola Broun. A. 7, xiv, 274, and S. 100.
Hydrostygnus Sharp, 1884.
linsi Sharp. C. 1020.
pontalis Broun. C. 84.
minor Broun. C. 1020.
bifoveatus Broun. C. 1021.

Hydrobius.

- **assimilis* Hope.
Exydrus Broun, 1886.
flavicornis Broun. C. 84 and C. 940.
gibbosus Broun. C. 84 and C. 940.
Stygnohydrus Broun, 1893.
nitidus Broun. C. 1336.
femoralis Broun. B. 1, i, 12.
posticalis Broun. B. 1, v, 371.
Giocyloma Broun, 1915.
nigratus Broun. B. 1, iv, 278.
Cyloma Sharp, 1872.
lawsonius Sharp. C. 85.
thomsonius Sharp. C. 1022.
guttulatus Sharp. C. 1022.
stewarti Broun. A. 6, xiv, 379.
Namostygynus Broun, 1909.
rufipes Broun. S. 99.
Cylomissus Broun, 1903.
glabratus Broun. A. 7, xi, 613.
Cyclonotum Dejean, 1833.
**marginale* Sharp. C. 83.
Psephoboragus Broun, 1893.
signatus Broun. C. 1403.
lineatus Broun. C. 1404.
dispar Broun. B. 1, i, 13.
Zeadolopus Broun, 1903.
spinipes Broun. A. 7, xi, 614.
Adolopus Sharp, 1884.
helmsi Sharp. C. 1023.
badius Broun. C. 86.
altulus Broun. C. 86.
vicus Broun. C. 939.
rugipennis Broun. C. 939.
montanus Broun. C. 1336.
converus Broun. C. 1404.
australis Broun. A. 8, iii, 223.
tibialis Broun. A. 8, iii, 224.
Cercyodes Broun, 1893.
laevigatus Broun. C. 941.
Cercyon.
**flavipes* Fabricius.
Hydraenodes Broun, 1919.
spinospennis Broun. B. 1, vi, 479.
Paracymus.
*nitidiusculus Broun.

Group BRACHELYTRA.

Family STAPHYLINIDAE.

Subfamily ALBOCHARINAE.

- Falagria* Mannerheim, 1830.
micans Broun. C. 96.
subopaca Broun. C. 1405.
- Ocalea* Erichson, 1837.
socialis Broun. C. 90.
crepera Broun. A. 6, xiv, 380.
- Baeostethus* Broun, 1909.
chiltoni Broun. S. 97.
- Aleochara* Gravenhorst, 1802.
puber Broun. C. 90.
semifusca Broun. A. 6, xiv, 386.
subaenea Fauvel. C. 91.
- Prototristus* Broun. 1909.
minutus Broun. A. 8, iii, 226.
- Caloderia* Mannerheim, 1830.
flavescens Broun. C. 93.
optabilis Broun. C. 94.
aeraria Broun. C. 94.
thoracica Broun. C. 95.
fulgens Broun. C. 95.
alophaula Broun. C. 941.
sericophora Broun. A. 6, xiv, 381.
wallacei Broun. T. xliv, 596.
granifer Broun. A. 6, xiv, 381.
diversa Broun. A. 6, xiv, 382.
fultoni Broun. T. xliv, 596.
vestita Broun. A. 6, xiv, 383.
tumidella Broun. A. 6, xiv, 383.
bituberculata Broun. A. 6, xiv, 384.
fungicola Broun. A. 6, xiv, 384.
rufipennis Broun. C. 92.
- Tachysa* Erichson, 1837.
fuscicornis Broun. C. 92.
- Homalota* Mannerheim, 1825.
cornigera Broun. C. 91.
atriceps Broun. C. 91.
**sordida* Marsham.
- Polylobus* Solsky, 1851.
sternalis Broun. C. 88.
- Aphytopus* Sharp, 1886.
gracilis Sharp. C. 1025.
pictulus Broun. B. 1, ii, 87.
porosus Broun. T. xliv, 394.
granifer Broun. T. xliv, 395.
guinnessi Broun. T. xliv, 395.
- Gyrophaena* Mannerheim, 1830.
punctata Broun. C. 87.
nugae Broun. C. 88.
densicornis Broun. C. 89.
versicolor Broun. C. 89.
atriceps Broun. C. 89.
carnulata Broun. B. 1, ii, 86.
- Encephalus* Stephens, 1832.
latulus Broun. A. 6, xiv, 386.
- Myrmecopora* Sauley, 1864.
funesta Broun. T. xliv, 397.
granulata Broun. T. xliv, 398.

Subfamily DIGRAMMINAE.

- Digrammus* Fauvel, 1900.
miricollis Fauvel. *Rev. Ent. Fr.*, xix, 124.

Subfamily TACHYPORINAE.

- Conurus* Stephens, 1832.
languidus Broun. C. 97.
austerus Broun. C. 97.
subruber Broun. C. 98.
badius Broun. C. 98.
flavithorax Broun. C. 98.
acerbus Broun. C. 99.
atricapillus Broun. C. 99.
auricomus Broun. C. 100.
nubilus Broun. C. 100.
maculosus Broun. C. 100.
maorinus Broun. C. 1025.
brevicornis Broun. C. 1405.
niticollis Broun. C. 1406.
asperillus Broun. B. 1, ii, 87.
laetulus Broun. B. 1, ii, 88.
morosus Broun. B. 1, viii, 676.
antennalis Broun. B. 1, viii, 677.
seminudus Broun. B. 1, viii, 677.

Subfamily STAPHYLININAE.

- Quedius* Stephens, 1832.
ruficornis Broun. C. 111.
aeneocepis Broun. C. 111.
enodis Broun. C. 111.
veterarius Broun. C. 112.
vividus Broun. C. 112.
arctifrons Broun. C. 112.
fuscatus Broun. C. 1027.
antipodus Sharp. C. 1028.
edwardsi Sharp. C. 1029.
insolitus Sharp. C. 1029.
latifrons Sharp. C. 1030.
discrepans Broun. C. 1030.
latimanus Broun. C. 1031.
collinus Broun. C. 1031.
cavelli Broun. C. 1032.
agathis Broun. C. 1407.
sciticollus Broun. A. 6, xiv, 420.
conspicuellus Broun. A. 6, xiv, 421.
scutellaris Broun. A. 6, xiv, 422.
puncticollis Broun. A. 6, xiv, 422.
ambiguus Broun. A. 6, xiv, 423.
taieriensis Broun. A. 6, xiv, 424.
tinctellus Broun. B. 1, i, 14.
aeneiventris Broun. B. 1, i, 14.
longiceps Broun. B. 1, i, 15.
hilaris Broun. A. 8, iii, 226.
burrowsi Broun. B. 1, iv, 278.
hirtipennis Broun. B. 1, iv, 279.
hallianus Broun. B. 1, v, 372.
recticeps Broun. B. 1, v, 373.
megophthalmus Broun. B. 1, v, 373.
urbanus Broun. B. 1, vi, 480.
iridescens Broun. B. 1, vi, 480.
eruensis Broun. T. xliv, 398.
xenophaeus Broun. T. xliv, 399.
**fulgidus* Fabricius.
badius Broun. B. 1, viii, 678.
bryocaris Broun. B. 1, viii, 678.
maorinus Broun. B. 1, viii, 679.
ohaensis Broun. B. 1, viii, 679.
slipsensis Broun. B. 1, viii, 680.

Group BRACHELYTRA—continued.

- Cafroquedus* Sharp, 1866.
gularis Sharp. C. 1033.
Staphylinus Linnaeus, 1758.
oculatus Fabricius. C. 107.
huttoni Broun. C. 108.
Xantholinus Serville, 1825.
sharpae Broun. C. 102.
cultus Broun. C. 103.
arecae Broun. C. 103.
mediocris Broun. C. 104.
labralis Broun. C. 104.
socius Fauvel. C. 104.
puncticeps Broun. C. 104.
**punctulatus* Paykull.
Pachycorynus Motschulsky, 1858.
dimorphus Fauvel. *Rev. Ent. Soc. Fr.*, 1892, 92.
Metoponcus Kraatz, 1857.
brouni Sharp. C. 106.
fulvipes Broun. C. 106.
rufulus Broun. C. 106.
Cafius Leach, 1830.
litoreus Broun. C. 108.
maritimus Broun. C. 109.
quadri-impressus (fem.), White. C. 105.
No. 194 = *caviceps* (male) Broun. C. 942.
algophilus Broun. A. 6, xiv, 419.
Othius Stephens, 1832.
adustus Broun. C. 101.
angustus Broun. C. 1026.
puncticeps Broun. A. 6, xiv, 419.
Philonthus.
**aeneus* Rossi.
**scutellaris* Nordmann.
**affinis* Roth.
**nigritulus* Gravenhorst.

Subfamily PAEDERINAE.

- Lithocaris* Lacordaire, 1835.
zealandicus Redtenbacher. C. 114.
mandibularis Broun. C. 114.
ventralis Broun. C. 115.
caecus Broun. A. 6, xiv, 424.
granipennis Broun. B. 1, i, 16.
longipennis Broun. T. xliv, 399.
Phanophilus Sharp, 1886.
comptus Broun. C. 1026 and 114.
Hyperomma Fauvel, 1878.
dispersum Fauvel. C. 1408.
mandibulare Broun. C. 1409.
duplicatum Broun. C. 1409.
sanguineum Broun. A. 6, xiv, 425.
tenellum Broun. A. 8, iii, 227.
subcaecum Broun. B. 1, vi, 481.
lobatum Broun. B. 1, vi, 481.
picipenne Broun. B. 1, vi, 482.
discrepans Broun. B. 1, vi, 483.
flavipes Broun. B. 1, viii, 681.
Dimerus Fauvel.
stibus Broun. B. 1, i, 15.
bisulciceps Broun. B. 1, v, 374.
vicinus Broun. B. 1, vi, 483.
whitehorni Broun. T. xliv, 400.

- Subfamily OXYTELINAE.
- Coprostygynus* Sharp, 1886.
sculptipennis Sharp. C. 1027.
optandus Broun. C. 1409.
curvipes Broun. B. 1, vi, 484.
picipennis Broun. B. 1, vi, 485.
Holotrochus Erichson, 1840.
ferrugineus Broun. C. 1034.
brevipennis Broun. C. 1034.
laevigatus Broun. B. 1, i, 16.
caecus Broun. B. 1, i, 17.
gracilis Broun. B. 1, i, 17.
nitidorsis Broun. B. 1, i, 17.
montanus Broun. B. 1, i, 18.
setigerus Broun. T. xliv, 401.
Phloeoneus.
cordicollis Broun. A. 7, xix, 57.
Bledius Leach.
bideefrons Broun. T. xliv, 401.
Oxytelus.
**rugosus* Fabricius.
- Subfamily OMALINAE.
- Omalium* Gravenhorst, 1802.
hebes Broun. C. 116.
suclithorax Broun. C. 116.
agrestis Broun. C. 117.
spadix Broun. C. 117.
pullum Broun. C. 120.
tibiale Broun. C. 120.
littorum Broun. C. 942.
sagoloide Sharp. C. 1035.
actobium Broun. C. 1035.
cariniferum Broun. C. 1036.
antipodum Broun. C. 1036.
chalmeri Broun. C. 1037.
micrathrum Broun. C. 1037.
cognatum Broun. C. 1038.
monilifer Broun. C. 1038.
debile Broun. C. 1039.
prolixum Broun. C. 1039.
stenosoma Broun. C. 1040.
tenellum Broun. C. 1410.
fusciventre Broun. A. 6, xiv, 426.
australe Broun. A. 6, xiv, 426.
philpotti Broun. A. 6, xiv, 427.
perplexum Broun. A. 6, xiv, 428.
nemobium Broun. B. 1, i, 18.
laetipenne Broun. B. 1, i, 19.
longiceps Broun. B. 1, ii, 89.
hilarum Broun. B. 1, ii, 90.
sanguineum Broun. B. 1, ii, 90.
planimarginatum Broun. A. 8, iii, 229.
setipes Broun. A. 8, iii, 230.
robustum Broun. T. xlvi, 96.
fossigerum Eppelshiem. T. xlvi, 93.
venator Broun. S. 98.
flavipalpi Broun. A. 8, iii, 228.
Eupsorus Broun, 1904.
costatus Broun. A. 7, xiv, 46.
Ischnoderus Fauvel, 1867.
tectus Broun. C. 117.
genalis Broun. C. 118.

Group BRACHELYTRA—continued.

Ischnoderus—continued.

- pictulus* Broun. C. 1040.
morosus Broun. C. 1041.
fultoni Broun. C. 1041.
politulus Broun. C. 119.
cognatus Broun. B. 1, i, 20.
arecubius Broun. B. 1, i, 20.
curtipennis Broun. B. 1, iv, 280.
Omalissus Broun, 1893.
castaneus Broun. C. 1042.

Omalissus—continued.

- scutosus* Broun. B. 1, iv, 281.
Trogophlaeus Mannerheim, 1830.
zealandicus Sharp. *Ent. Mag.*, xxxvi, 233.
maritimus Broun. A. 7, xi, 615.
Blediotrogus Sharp, 1900.
guttiger Sharp. *Ent. Mag.*, xxxvi, 234.
cribricollis Fauvel. *Rev. Ent. Fr.*, xix, 184.

Group CLAVICORNES.

Family CLAMBIDAE.

- Clambus* Fisher, 1820.
domesticus Broun. C. 762.
vestitus Broun. C. 762.
suffusus Broun. C. 762.

Family SILPHIDAE.

- Necrophilus* Latreille, 1829.
prolongatus Sharp. C. 760.
Camiarus Sharp, 1876.
thoracicus Sharp. C. 148.
indiscretus Broun. C. 149.
convexus Sharp. C. 150.
estriatus Broun. T. xliv, 415.
Zenocolon Broun, 1917.
laevicollis Broun. B. 1, v, 387.
Inocatops Broun, 1893.
compactus Broun. C. 1066.
flectipes Broun. C. 1067.
nigrescens Broun. C. 1339.
concininus Broun. C. 150.
spinifer Broun. B. 1, v, 383.
granipennis Broun. B. 1, v, 384.
elongellus Broun. B. 1, v, 384.
separatus Broun. B. 1, v, 385.
impressus Broun. B. 1, vi, 520.

- Microsilpha* Broun, 1886.
litorea Broun. C. 895.
Asphaerites Broun, 1893.
nitidus Broun. C. 1068.
Catopsolius Sharp, 1886.
laevicollis Sharp. C. 1069.
Isocolon Broun, 1893.
hilaris Broun. C. 1070.
frontale Broun. B. 1, vi, 521.
modestum Broun. B. 1, vi, 521.
orruensis Broun. B. 1, viii, 683.
Choleva Latreille, 1796.
antennalis Broun. C. 152.
alacris Broun. C. 152.
granifer Broun. C. 890.
fulvitaris Broun. C. 946.
lugubris Sharp. C. 1069.
relata Broun. C. 1339.
phyllobia Broun. C. 1433.
monticola Broun. C. 1434.
hunuensis Broun. C. 1434.
marginalis Broun. A. 6, xv, 87.
suturalis Broun. A. 6, xv, 88.

Choleva—continued.

- nemoralis* Broun. A. 8, iii, 230.
caeca Broun. T. xliv, 414.
castanea Broun. T. xliv, 415.
brunneipes Broun. T. xlvi, 97.
avivorus Broun. S. 101.
crenata Broun. B. 1, vii, 609.
flectipes Broun. B. 1, vii, 609.
curvicerca Broun. B. 1, vii, 610.
Zeagyrtes Broun, 1917.
vitticollis Broun. B. 1, v, 386.
Allotropis Broun, 1893.
ovalis Broun. C. 1435.
Mesocolon Broun, 1880.
clathrata Broun. C. 153.
liturata Broun. C. 154.
puncticeps Broun. C. 154.
undulata Broun. C. 154.
bicolor Broun. C. 155.
nebulosa Broun. C. 155.
maculifer Broun. C. 156.
hirtalis Broun. C. 156.
punctulata Broun. C. 157.
torva Broun. C. 157.
domestica Broun. C. 158.
varia Broun. C. 946.
strigicollis Broun. B. 1, vii, 610.
nitidulus Broun. B. 1, vii, 611.
labralis Broun. B. 1, vii, 611.
Baeosilpha Broun, 1895.
rufescens Broun. A. 6, xv, 85.
Mesagyrtes Broun, 1895.
scabripes Broun. A. 6, xv, 86.
Silphotelus Broun, 1895.
nitidus Broun. A. 6, xv, 84.
obliquus Broun. T. xliv, 416.

Family SCYDMAENIDAE.

- Scydmaenus* Latreille, 1802.
edwardsi Sharp. C. 144.
ambiguus Broun. C. 145.
puncticollis Broun. C. 146.
princeps Broun. C. 924.
galeatus Broun. C. 924.
elongellus Broun. C. 1338.
laetans Broun. C. 663.
ciliatus Broun. A. 6, xii, 178.
cedius Broun. A. 6, xii, 179.
allocerus Broun. A. 6, xii, 179.

Group CLAVICORNES—continued.

Scydmmaenus—continued.

- brachycerus* Broun. A. 6, xii, 180.
xanthopus Broun. A. 6, xii, 181.
heterarthus Broun. A. 6, xii, 181.
relatus Broun. A. 6, xii, 182.
insignis Broun. A. 6, xii, 182.
angulifrons Broun. B. 1, iv, 307.
sulciferus Broun. B. 1, iv, 308.
latuliceps Broun. B. 1, iv, 309.
fragilis Broun. B. 1, iv, 309.

Adrastia Broun, 1881.

- laetans* Broun. C. 663.
clavatella Broun. B. 1, iv, 310.

Phaganophana King, 1864.

- selosa* Sharp. C. 146.
picicollis Broun. C. 147.
calva Broun. C. 147.
oreas Broun. C. 925.
lanosa Broun. C. 925.
angustata Broun. C. 926.
planiceps Broun. C. 1063.
murroi Broun. C. 1063.
antennalis Broun. C. 1064.
stenocera Broun. C. 1064.
latiuscula Broun. C. 1064.
sanguinea Broun. C. 1065.
erythronota Broun. C. 1065.
ovipennis Broun. C. 1339.
russata Broun. C. 1432.
monstifer Broun. C. 1432.
angulata Broun. C. 1433.
alacer Broun. B. 1, iv, 310.
halli Broun. B. 1, iv, 311.
palpalis Broun. B. 1, iv, 312.
sinuata Broun. B. 1, iv, 313.

Sciacharis Broun, 1893.

- fulva* Broun. C. 1062.

Family PSELAPHIDAE.

Plesiotyrus Broun, 1914.

- crassipes* Broun. C. 1337.

Euglyptus Broun, 1893.

- elegans* Broun. C. 1412.
costifer Broun. A. 6, xii, 166.
foveicollis Broun. T. xliv, 403.
longicornis Broun. T. xliv, 404.
sublaevis Broun. B. 1, vii, 603.
longiceps Broun. B. 1, vii, 604.
abnormis Broun. B. 1, vii, 604.

Microtyrus Broun, 1893.

- punctatus* Broun. C. 1413.

Phormiobius Broun, 1917.

- halli* Broun. B. 1, v, 382.

Agatyrus Broun, 1917.

- fulvihirtus* Broun. B. 1, v, 383.
Tyrogetus Broun, 1893.

- oplandus* Broun. A. 6, xii, 166.

- palpalis* Broun. B. 1, i, 23.

Hamotulus Schaufuss, 1887.

- mutandus* Sharp. C. 121.

- armatus* Broun. C. 1043.

- spinipes* Broun. C. 1411.

- sternalis* Broun. A. 6, xii, 168.

- curvipes* Broun. A. 6, xii, 170.

Hamotulus—continued.

- frontalis* Broun. B. 1, ii, 94.

- angulipes* Broun. B. 1, iii, 171.

- robustus* Broun. B. 1, iv, 305.

- fuscipalpis* Broun. B. 1, iv, 306.

- cornutus* Broun. B. 1, iv, 307.

Pselaphus Herbst, 1792.

- pauper* Sharp. C. 122.

- pilosistratus* Broun. C. 123.

- dulcis* Broun. C. 660.

- caecus* Broun. C. 943.

- delicatus* Broun. C. 943.

- fuscopilus* Broun. C. 944.

- citimus* Broun. C. 1043.

- meliusculus* Broun. C. 1044.

- cavelli* Broun. C. 1414.

- sulcicollis* Broun. C. 1415.

- pilifrons* Broun. B. 1, ii, 93.

- trifoveatus* Broun. B. 1, iii, 172

- urquharti* Broun. B. 1, v, 379.

- oviceps* Broun. B. 1, v, 380.

- caridorsis* Broun. B. 1, viii, 681.

Gastrobothrus Broun, 1886 = *Physa* Rafra, 1890 (not of Draparnaud, 1801).

- inflata* Sharp. C. 124.

- sharp* Broun. C. 124.

- abdominalis* Broun. C. 125 and 758.

Eupines King, 1864.

- rudicornis* Broun. C. 759.

- micans* Sharp. C. 126.

- clavata* Broun. C. 126.

- picipes* Broun. C. 127.

- dispar* Sharp. C. 127.

- impressifrons* Broun. C. 128.

- mundula* Broun. C. 129.

- crassicornis* Broun. C. 129.

- deformis* Sharp. C. 130.

- impar* Sharp. C. 130.

- grata* Sharp. C. 131.

- altula* Broun. C. 131.

- nasuta* Broun. C. 132.

- sanguinea* Broun. C. 132.

- pagana* Broun. C. 660.

- ignota* Broun. C. 661.

- glabrata* Broun. C. 830.

- nemoralis* Broun. C. 831.

- punctata* Broun. C. 831.

- calcarata* Broun. C. 831.

- sylvicola* Broun. C. 919.

- fraudulenta* Broun. C. 944.

- epulectoides* Broun. C. 1045.

- decens* Broun. C. 1046.

- murroi* Broun. C. 1046.

- forficulata* Broun. C. 1047.

- foveatissima* Broun. C. 1047.

- platynota* Broun. C. 1338.

- conspicua* Broun. C. 1415.

- costata* Broun. C. 1416.

- platyarthra* Broun. C. 1417.

- mundula* Broun. C. 1417.

- fulvitarsis* Broun. C. 1418.

- sternalis* Broun. A. 6, xii, 171.

- allocera* Broun. A. 6, xii, 172.

- setifer* Broun. A. 6, xii, 173.

- diversa* Broun. A. 6, xii, 174.

Group CLAVICORNES—continued.

Eupines—continued.

- lewisi* Broun. B. 1, i, 24.
simplex, Broun. A. 8, xi, 206.
nesobia Broun. B. 1, ii, 92.
clemens, Broun. B. 1, vii, 607.
Parmipalpus Broun, 1881.
montivagus Broun. C. 662.
Startes Broun, 1886.
sculpturata Broun. C. 830.
foveata Broun. A. 6, xii, 170.
Dalmus Sharp, 1874.
pubescens Sharp. C. 133.
gigantea Broun. B. 1, iii, 164.
graniceps Broun. B. 1, vii, 605.
Dalmis Sharp, 1886.
batriscodes Sharp. C. 1049.
Sagola Sharp, 1874.
major Sharp. C. 135.
prisca Sharp. C. 136.
misella Sharp. C. 136.
parva Sharp. C. 136.
pulcher Broun. C. 137.
notabilis Broun. C. 137.
deformipes Broun. C. 138.
denticollis Broun. C. 138.
genalis Broun. C. 663.
terricola Broun. C. 832.
castanea Broun. C. 884.
excavata Broun. C. 884.
sulcator Broun. C. 885.
osculans Broun. C. 885.
fovealis Broun. C. 886.
elevata Broun. C. 886.
brevitarsis Broun. C. 887.
bispunctata Broun. C. 887.
duplicata Broun. C. 888.
tenuis Broun. C. 888.
convexa Broun. C. 889.
insignis Broun. C. 1049.
hirtalis Broun. C. 1050.
sobrina Broun. C. 1050.
rectipes Broun. C. 1051.
insolens Broun. C. 1051.
punctata Broun. C. 1052.
fulva Broun. C. 1052.
ruficeps Broun. C. 1053.
parallelia Broun. C. 1053.
anisarthra Broun. C. 1053.
macronyx Broun. C. 1418.
mimica Broun. C. 1419.
robusta Broun. C. 1420.
luminata Broun. C. 1421.
immota Broun. C. 1422.
flavipes Broun. C. 1422.
elongata Broun. C. 1423.
lineata Broun. A. 6, xii, 175.
pertinax Broun. A. 6, xii, 176.
citima Broun. A. 6, xii, 177.
rugifrons Broun. A. 6, xv, 73.
spinifer Broun. A. 6, xv, 75.
eminens Broun. A. 6, xv, 75.
punctulata Raffray. Rev. Ent. Fr., 1893,
 22.
frontalis Raffray. Rev. Ent. Fr., 1893,
 23.

Sagola—continued.

- sharpi* Raffray. Rev. Ent. Fr., 1893, 26.
brevicornis Raffray. Rev. Ent. Fr., 1893,
 27.
carinata Broun. A. 8, x, 622.
ventralis Broun. A. 8, x, 623.
occipitalis Broun. A. 8, x, 624.
longula Broun. A. 8, x, 625.
pallidula Broun. A. 8, x, 626.
spiniventris Broun. A. 8, x, 627.
grata Broun. A. 8, x, 628.
bifoveiceps Broun. A. 8, x, 629.
biimpressa Broun. A. 8, x, 630.
clavatella Broun. A. 8, x, 631.
lawsoni Broun. A. 8, x, 632.
latula Broun. A. 8, x, 633.
puncticeps Broun. A. 8, viii, 489.
laticeps Broun. A. 8, viii, 490.
angulifer Broun. A. 8, viii, 491.
nitida Broun. A. 8, viii, 492.
cognata Broun. A. 8, viii, 494.
diversa Broun. A. 8, viii, 495.
latistriata Broun. A. 8, viii, 495.
basalis Broun. A. 8, viii, 496.
longicollis Broun. A. 8, viii, 498.
puncticollis Broun. A. 8, viii, 499.
auripila Broun. A. 8, viii, 500.
guinessi Broun. A. 8, viii, 502.
longipennis Broun. A. 8, viii, 504.
electa Broun. B. 1, ii, 91.
halli Broun. B. 1, iii, 155.
colorata Broun. B. 1, iii, 156.
dilucida Broun. B. 1, iii, 157.
insueta Broun. B. 1, iii, 157.
suturalis Broun. B. 1, iii, 158.
fuscipalpis Broun. B. 1, iii, 159.
disparata Broun. B. 1, iii, 160.
bituberata Broun. B. 1, iii, 160.
socia Broun. B. 1, iv, 281.
setiventris Broun. B. 1, iv, 282.
crassulipes Broun. B. 1, iv, 283.
opercularis Broun. B. 1, iv, 284.
brevisternis Broun. B. 1, iv, 284.
rustica Broun. B. 1, iv, 285.
confusa Broun. B. 1, iv, 286.
longipes Broun. B. 1, iv, 287.
indiscreta Broun. B. 1, iv, 288.
rotundiceps Broun. B. 1, iv, 289.
fulvipennis Broun. B. 1, iv, 289.
bifida Broun. B. 1, iv, 290.
posticalis Broun. B. 1, iv, 291.
laetula Broun. B. 1, iv, 291.
concolorata Broun. B. 1, iv, 292.
unicalis Broun. B. 1, v, 376.
dickensis Broun. B. 1, v, 377.
robustula Broun. B. 1, v, 378.
hectorii Broun. B. 1, v, 378.
monticola, Broun. T. xliv, 402.
bilobata Broun. B. 1, vi, 486.
tenebrica Broun. B. 1, vi, 487.
subcuneata Broun. B. 1, vi, 488.
strialis Broun. B. 1, vi, 489.
rectipennis Broun. B. 1, vi, 489.
valida Broun. B. 1, vi, 490.
ciliipes Broun. B. 1, vi, 491.

Group CLAVICORNES—continued.

Sagola—continued.

- joveiventris* Broun. B. 1, vi, 492.
striatifrons Broun. B. 1, vi, 492.
cordiceps Broun. B. 1, vi, 493.
fagicola Broun. B. 1, vi, 494.
ignota Broun. B. 1, vi, 495.
furcata Broun. B. 1, vi, 495.
aemula Broun. B. 1, vi, 496.
fusciculata Broun. B. 1, vi, 497.
minuscula Broun. B. 1, vi, 497.
acuminata Broun. B. 1, vi, 498.
rufescens Broun. B. 1, vi, 499.
bipuncticeps Broun. B. 1, vi, 499.
affinis Broun. B. 1, vi, 500.
planipennis Broun. B. 1, vi, 500.
brevifossa Broun. B. 1, vi, 501.
arboricola Broun. B. 1, vi, 502.
remixta Broun. B. 1, vi, 502.
planicula Broun. B. 1, vi, 503.
lineiceps Broun. B. 1, vi, 504.
disonans, Broun. B. 1, vii, 601.
distorta Broun. B. 1, vii, 602.

Stenosagola Broun, 1919.

- gracilis* Broun. C. 1424.
crassicornis Broun. A. 8, viii, 501.
connata Broun. A. 8, viii, 503.
planiocula Broun. B. 1, vi, 505.
oblongiceps Broun. B. 1, vi, 505.
griseipila Broun. B. 1, vi, 506.

Exeirarthra Broun, 1893.

- enigma* Broun. C. 1054.
palida Broun. C. 1424.
angustula Broun. B. 1, v, 374.
longiceps Broun. B. 1, v, 375.
parviceps Broun. B. 1, vi, 486.
- Vidamus* Raffray, 1898.
- convexus* Sharp. C. 139.
modestus Broun. A. 8, xi, 203.
sternalis Broun. A. 8, xi, 204.
trochanteralis Broun. A. 8, viii, 691.
armiferus Broun. A. 8, viii, 692.
congruus Broun. B. 1, iv, 301.
bryophilus Broun. B. 1, iii, 166.
clavipes Broun. B. 1, iv, 302.
punctulatus Broun. B. 1, iv, 302.
gracilipes Broun. B. 1, v, 379.
simplex Broun. B. 1, vi, 511.
ovicollis Broun. B. 1, vi, 511.
muscicola Broun. B. 1, vi, 512.
nitidus Broun. B. 1, vi, 513.
fossalis Broun. B. 1, vi, 513.
calcaratus Broun. T. xliv, 411.
incertus Reitter. T. xliv, 411.

Vidamodes Broun, 1919.

- furvis* Broun. B. 1, vii, 606.
Sagolonus Raffray, 1898.
patronus Broun. C. 1060.
impressus Broun. B. 1, i, 21.

Gabata Raffray.

- bifoveata* Broun. B. 1, vi, 514.
parallela Broun. B. 1, vi, 515.
spinipes Broun. B. 1, i, 21.

Zelandius Raffray, 1898.

- asper* Broun. C. 140.
usitatus Broun. B. 1, i, 22.

Zelandius—continued.

- fovealis* Broun. A. 8, xi, 205.
raffrayi Broun. A. 8, viii, 693.
illustris Broun. A. 8, viii, 695.
fulgens Broun. A. 8, viii, 696.
basalis Broun. B. 1, iii, 167.
spinifer Broun. B. 1, iii, 167.
brookesi Broun. B. 1, iv, 303.
tuberculata Broun. B. 1, iv, 304.

Whitea Hutton, 1894 = *Brouna*, Raffray
1898 (not of Sharp, 1878).*laevifrons* Broun. C. 1425.*Plectomorphus* Raffray, 1898.*spinifer* Broun. C. 1426.*longiceps* Broun. A. 8, xi, 200.*brevicornis* Broun. A. 8, xi, 201.*egenus* Broun. A. 8, xi, 202.*laminifer* Broun. B. 1, iv, 300.*scitiventris* Broun. B. 1, vi, 507.*anguiferus* Broun. B. 1, vi, 508.*rugiceps* Broun. B. 1, vi, 509.*insignis* Broun. B. 1, vi, 509.*collinus* Broun. B. 1, vi, 510.*optandus* Broun. T. xliv, 412.*longipes* Broun. T. xliv, 412.*Euplectus* Leach, 1817.*trisulcicollis* Broun. C. 140.*longulus* Broun. C. 141.*opacus* Sharp. C. 141.*sculpturatus* Broun. C. 142.*frontalis* Broun. C. 142.*brevitarsis* Broun. C. 143.*ovicollis* Broun. C. 143.*foveolatus* Broun. C. 143.*tuberigerus* Broun. C. 760.*crassipes* Broun. C. 919.*patruelis* Broun. C. 919.*monticola* Broun. C. 920.*ovithorax* Broun. C. 920.*obnusus* Broun. C. 921.*vacuus* Broun. C. 921.*incomptus* Broun. C. 921.*mirificus* Broun. C. 922.*u-impressus* Broun. C. 923.*careus* Broun. C. 923.*eminens* Broun. C. 945.*auripilus* Broun. C. 945.*morens* Broun. C. 1055.*lepidophorus* Broun. C. 1056.*validus* Broun. C. 1056.*scruposus* Broun. C. 1057.*allocephalus* Broun. C. 1057.*acuminatus* Broun. C. 1058.*clevedonensis* Broun. C. 1058.*sandageri* Broun. C. 1059.*personatus* Broun. C. 1059.*unicus* Broun. C. 1060.*verticalis* Broun. C. 1061.*antiquus*, Broun. C. 1061.*munroi* Broun. C. 1426.*obscurus* Broun. C. 1427.*coeruleus* Broun. C. 1428.*inscitus* Broun. C. 1428.*iracundus* Broun. C. 1429.*caudatus* Broun. C. 1429.

Group CLAVICORNES—continued.

Euplectus—continued.

- tumipes* Broun. A. 6, xv, 76.
parvulus Broun. A. 6, xv, 77.
modestus Broun. A. 6, xv, 78.
arohaensis Broun. A. 6, xv, 79.
foveiceps Broun. A. 6, xv, 80.
semiopacus Broun. A. 6, xv, 81.
pusillus Broun. A. 6, xv, 82.
caviceps Broun. A. 7, xiv, 48.
sulciceps Broun. A. 7, xiv, 49.

Pycnoplectus Reitter.

- cephalotes* Reitter. T. xliv, 410.

Byraxis Reitter.

- illustris* Broun. B. 1, iii, 168.
anisarthra Broun. B. 1, iii, 169.
bisulcifrons Broun. B. 1, iii, 170.
lewisi Broun. B. 1, iii, 170.
monstrosa Reitter. T. xliv, 413.
rhyssarthra Broun. T. xliv, 413.
halli Broun. B. 1, vii, 608.
acceptus Broun. B. 1, viii, 683.

Alloplectus Broun, 1911.

- claviger* Broun. C. 1480.
picipennis Broun. A. 8, viii, 698.
subcaecus Broun. A. 8, viii, 699.

Kenoccelus Broun, 1911.

- dimorphus* Broun. A. 8, viii, 701.
Eleusomatus Raffray.

- ovicollis* Broun. B. 1, iv, 305.
vidamoides Broun. B. 1, vi, 516.
oculatus Broun. B. 1, vi, 516.
subcaecus Broun. B. 1, vi, 516.

Placodium.

- zenarthrum* Broun.

Patreus Broun, 1904.

- lewisi* Broun. A. 7, xiv, 47.

Euplectopsis Raffray.

- sanguineus* Broun. A. 8, xi, 196.
duplicatus Broun. A. 8, xi, 197.
clavatulus Broun. A. 8, xi, 199.
granulatus Broun. A. 8, viii, 686.
fastigiatus Broun. A. 8, viii, 687.
eruensis Broun. B. 1, iv, 294, and
T. xliv, 408.

- mucronellus* Broun. A. 8, viii, 689.
tumidus Broun. A. 8, viii, 690.
tibialis Broun. B. 1, iii, 161.
terrestris Broun. B. 1, iii, 162.
femoralis Broun. B. 1, iii, 163.
curvipennis Broun. B. 1, iii, 164.
perpunctatus Broun. B. 1, iv, 293.
nitipennis Broun. B. 1, iv, 294.
duplex Broun. B. 1, iv, 295.
cuneiceps Broun. B. 1, iv, 296.
bryocharis Broun. B. 1, iv, 296.
blandiatus Broun. B. 1, iv, 297.
impressus Broun. B. 1, iv, 298.
dorsalis Broun. B. 1, iv, 299.
elongellus Broun. B. 1, iv, 299.
crassulus Broun. B. 1, vi, 506.
longicollis Reitter. T. xliv, 405.
macrocephalus Reitter. T. xliv, 405.
brevicollis Reitter. T. xliv, 405.
rotundicollis Reitter. T. xliv, 405.

Euplectopsis—continued.

- trichonyformis* Reitter. T. xliv, 406.
schizocnemis Broun. T. xliv, 406.
carinatus Broun. T. xliv, 407.
antennalis Broun. T. xliv, 407.
heterarthrus Broun. T. xliv, 409.
biimpressus Broun. T. xliv, 410.
Eupinogitus Broun, 1919.
sulcipennis Broun. B. 1, vi, 518.
Neosampa Broun, 1919.
granulata Broun. B. 1, vi, 519.
Anabasis Raffray.
minor Broun. B. 1, vi, 519.

Family TRICHOPTERYGIDAE.

- Trichopteryx* Kirby, 1826.
inconspicua Matthews. C. 1436.
Ptenidium Erichson, 1845.
lawsoni Matthews. C. 1437.
posticale Broun. C. 1437.
proprium Broun. C. 1438.
cavelli Broun. C. 1438.
Actidiump Matthews, 1863.
lineare Matthews. C. 1439.
Actinopteryx Matthews, 1872.
australis Matthews. C. 1440.
Ptinella Motschoulski, 1844.
fauveli Matthews. Mon. *Trichop.*
Suppl., 19.

Family CORYLOPHIDAE.

- Anisomeristes* Matthews, 1886.
sharpi Matthews. Ent. Mon. Mag.,
1886, xxii, 425.
ater Matthews. Ent. Mon. Mag., 1886,
xxii, 425.
Sacina Broun, 1893.
oblonga Broun. C. 1071.
laetans Broun. B. 1, ii, 101.
curlula Broun. B. 1, iii, 173.
Sericoderus Stephens, 1829.
fulvicollis Broun. C. 1072.
scutellaris Broun. C. 1073.
basalis Broun. C. 1073.

Family COCCINELLIDAE.

- Coccinella* Linnaeus, 1735.
tasmanii White. C. 645.
coriacea Broun. C. 1319.
novae-zelandiae Colenso. T. xx, 4.
**11-punctata* Linnaeus.
**californica* Mannerheim.
**sanguinea* Linnaeus.
Leis Mulsant, 1851.
antipodum Mulsant. C. 645.
**conformis* Boisduval.
Rhizobius.
**ventralis* Erichson.
Cranophorus Mulsant, 1851.
venustus Pascoe. C. 650.
Orcus.
**chalybeus* Boisduval.
**australasiae* Boisduval.

Group CLAVICORNES—continued.

- Scymnus* Kugelann, 1794.
eximius Broun. C. 646.
fiavikirtus Broun. C. 646.
acceptus Broun. C. 647.
macrostictus Broun. T. xlii, 114.
consors Broun. C. 647.
tristis Broun. C. 647.
pallidiceps Broun. C. 648.
fagus Broun. C. 648.
terrenus Broun. C. 648.
rarus Broun. C. 649.
suffusus Broun. C. 649.
minutulus Broun. C. 649.
piceinus Broun. C. 649.
sedatus Broun. C. 813.
villosum Broun. C. 814.
circularis Sharp. *Insect Life*, 1889,
 365.
prolongatus Broun. B. 1, iii, 264.
halli Broun. B. 1, iii, 265.
circularis Sharp. B. 1, iii, 265.
nigritulus Broun. B. 1, iii, 265.
Novius.
**cardinalis* Mulsart.
Hippodamia.
**convergens* Guérin.
Holopsis Broun, 1886.
nigellus Broun. C. 814.
pallidus Broun. C. 815.
lawsoni Broun. C. 815.
pictulus Broun. C. 1503.
rotundatus Broun. C. 1503.
Cryptolaemus.
**montrouzieri* Mulsart.
Veronicobius Broun, 1893.
hirtus Broun. C. 1393.

Family EROTYLIDAE.

- Cryptodacne* Sharp, 1878.
synthetica Sharp. C. 641.
lenis Broun. C. 641.
vagepunctata Broun. C. 813.
vittata Broun. C. 873.
pubescens Broun. C. 1319.
ferrugata Broun. B. 1, i, 78.
ocularia Broun. T. xlv, 163.
Triplax Paykull, 1798.
brouni Pascoe. C. 642.
Thallis Erichson, 1842.
polita White. C. 643.
Tritomidea.
rubripes Reitter. B. 1, i, 78.

Family COLYDIIDAE.

Subfamily COLYDIINAE.

- Tarphiomimus* Wollaston, 1873.
indentatus Wollaston. C. 182.
wollastoni Sharp. C. 1080.
tuberculatus Broun. T. xliv, 417.
Dryptope Broun, 1886.
dorsalis Broun. C. 763.
undosus Broun. C. 764.
acuminatus Broun. C. 183.

- Ulonotus* Erichson, 1845.
antarcticus White. C. 187.
discedens Sharp. C. 187.
brouni Sharp. C. 188.
viridipictus Wollaston. C. 188.
asper Sharp. C. 189.
integer Sharp. C. 189.
lawsoni Wollaston. C. 189.
aberrans Broun. C. 189.
atratus Broun. C. 190.
punctatus Broun. C. 894.
isostictus Broun. C. 926.
carus Broun. C. 947.
fuscatus Broun. C. 948.
rufescens Broun. C. 948.
cinereus Broun. C. 948.
integratus Broun. C. 949.
fulvus Broun. C. 1080.
dissimilis Sharp. C. 1081.
philpotti Broun. B. 1, ii, 95.
proprius Broun. B. 1, iii, 174.
planiceps Broun. B. 1, iv, 315.
discalis Broun. B. 1, vi, 525.
uropterus Broun. T. xliv, 418.
wallacei Broun. T. xliv, 419.
plagiatus Broun. T. xlii, 98.
affinis Broun. B. 1, viii, 683.
aenulus Broun. B. 1, viii, 684.
Phormesa Pascoe, 1863.
costicollis Reitter. B. 1, i, 38.
Notonulus Broun, 1886 (substituted for
Ablabus).
ornatus Broun. C. 184.
pallidipictus Broun. C. 185.
scabrus Broun. C. 185.
fervidulus Broun. C. 186.
punctipennis Broun. C. 186.
brevis Broun. C. 763.
nodosus Broun. C. 894.
sparsus Broun. C. 947.
libentus Broun. C. 947.
facetus Broun. C. 1341.
varicornis Broun. B. 1, i, 38.
crassulus Broun. B. 1, ii, 96.
truncatus Broun. B. 1, iii, 175.
longipes Broun. B. 1, iii, 176.
discors Broun. B. 1, vi, 526.
demissus Broun. T. xliv, 419.
sellata Sharp. C. 1081.
Allobitoma Broun, 1919.
halli. B. 1, vi, 527.
Bitoma Herbst, 1793.
insularis White. C. 192.
vicina Sharp. C. 193.
distans Sharp. C. 193.
rugosa Sharp. C. 193.
nana Sharp. C. 194.
distincta Broun. C. 194.
discoidea Broun. C. 195.
lobata Broun. C. 893.
scita Broun. C. 895.
guttata Broun. C. 895.
auriculata Sharp. C. 1082.
serraticula Sharp. C. 1083.
mundula Sharp. C. 1083.

Group CLAVICORNES—continued.

Bitoma—continued.*picicorne* Broun. A. 8, iii, 285.*obsoleta* Broun. B. 1, iii, 176.*maura* Broun. T. xliv, 420.*morosa* Broun. B. 1, vii, 613.*Cozelus* Latreille, 1829.*dubius* Sharp. C. 196.*similis* Sharp. C. 196.*robustus* Broun. C. 197.*clarus* Broun. C. 766.*grossanus* Broun. C. 927.*chalmeri* Broun. C. 949.*rufus* Broun. C. 1084.*posticalis* Broun. C. 1084.*ovicollis* Broun. C. 1084.*longulus* Broun. C. 1085.*oculator* Broun. C. 1342.*picicornis* Broun. C. 1342.*graniceps* Broun. C. 1343.*regularis* Broun. C. 1440.*thoracicus* Broun. A. 6, xv, 194.*helmsi* Reitter. B. 1, i, 37.*elongatus* Broun. A. 8, iii, 386.*variegatus* Broun. A. 8, iii, 387.*bicavus* Broun. A. 8, iii, 388.*instabilis* Broun. B. 1, iii, 177.*muconatus* Broun. T. xlvi, 98.*Heterargus* Sharp, 1886.*rudis* Sharp. C. 1086.*serricollis* Broun. C. 1441.*subaequus* Broun. B. 1, ii, 97.*parallelus* Broun. B. 1, iii, 178.*Gathocles* Broun, 1893.*nodosus* Broun. C. 1086.*obliquicostatus* Broun. A. 8, iii, 388*angulifer* Broun. B. 1, iii, 178.*fuscus* Broun. B. 1, viii, 684.*interruptus* Broun. B. 1, viii, 685*Vitiacus* Broun, 1893.*costatus* Broun. C. 1087.*costicollis* Broun. C. 1442.*incurvus* Broun. A. 6, xv, 195.*subcaecus* Broun. B. 1, vi, 528.*posticalis* Broun. B. 1, vi, 529.*purus* Broun. B. 1, vi, 529.*suturalis* Broun. B. 1, vi, 530.*lateralis* Broun. B. 1, vi, 531.*Enarsus* Pascoe, 1866.*bakewellii* Pascoe. C. 199.*wakefieldi* Sharp. C. 199.*rudis* Sharp. C. 200.*contractifrons* Broun. C. 200.*probus* Broun. C. 1088.*cucullatus* Sharp. C. 1089.*Rhitidinotus* Broun, 1880.*squamulosus* Broun. C. 204.*Synclus* Sharp, 1876.*optatus* Sharp. C. 201.*politus* Broun. C. 201.*hystrix* Sharp. C. 202.*piciceps* Broun. C. 1092.*monroi* Broun. C. 1442.*explanatus* Broun. T. xliv, 417.*solidus* Broun. B. 1, viii, 685*Acosmetus* Broun, 1880.*oblongus* Broun. C. 197.*granulatus* Broun. C. 197.*Protarphius* Broun, 1893.*ruficornis* Broun. A. 6, xii, 184.*indentatus* Broun. A. 6, xii, 185.*crassus* Broun. C. 669.*decorus* Broun. B. 1, ii, 97.*tricarus* Broun. A. 8, iii, 389.*posticalis* Broun. A. 8, iii, 390.*palleus* Broun. B. 1, iii, 179.*Recyntus* Broun, 1886.*exiguus* Broun. C. 765.*tuberculatus* Broun. C. 191.*insignis* Broun. C. 191.*salebrosum* Broun. C. 192.*reitteri* Broun. B. 1, viii, 686.*Sympsius* Broun, 1909.*serratus* Broun. A. 8, iii, 391.*lobifer* Broun. A. 8, iii, 392.*Epistranus* Sharp, 1876.*lawsoni* Sharp. C. 203.*humeralis* Broun. C. 203.*parvus* Broun. C. 950.*optabilis* Broun. C. 1343.*hirtalis* Broun. A. 6, xii, 189.*valens* Broun. C. 670.*sharpi* Reitter. B. 1, i, 37.*fulvus* Reitter. B. 1, i, 37*Noris* Broun, 1893.*crassus* Broun. C. 1091.*Glenentela* Broun. 1893.*serrata* Broun. C. 1090.*costata* Broun. B. 1, vi, 527.*Ithris* Pascoe, 1864.*gracilis* Sharp. C. 205.*Rhizonium* Sharp, 1877.*antiquum* Sharp. C. 206.*Chorasus* Sharp, 1882.*subcaecus* Sharp. C. 1093.*Archaeoglenea* Broun, 1893.*costipennis* Broun. A. 6, xii, 189.*Lithostygus* Broun, 1886.*costatus* Broun. C. 950.*serripennis* Broun. B. 1, iii, 185.*cuneiceps* Broun. B. 1, iii, 186.*Ciconissus* Broun, 1893.*granifer* Broun. A. 6, xii, 186.

Subfamily BOTRIDERINAE.

Bothrideres Erichson, 1845.*moestus* Sharp. C. 207.*cognatus* Sharp. C. 1096.*obsoletus* Broun. A. 6, xv, 196.*picipes* Broun. A. 7, xi, 618.*diversus* Broun. T. xliv, 422.*paynteri* Broun. T. xlii, 100.

Subfamily PYCNOTERINAE.

Pycnoterus Erichson, 1842.*sophorae* Sharp. C. 208.*simulans* Sharp. C. 209.*simplex* Broun. C. 209.

Group CLAVICORNES—continued.

- Pycnomerus*—continued.
minor Sharp. C. 209.
ellipticus Broun. C. 210.
rufescens Broun. C. 766.
basalis Broun. C. 766.
carinellus Broun. C. 896.
caecus Broun. C. 896.
hirtus Broun. C. 897.
lateralis Broun. C. 897.
arboreus Broun. C. 927.
cognatus Broun. C. 951.
marginalis Broun. C. 1093.
sinuatus Broun. C. 1094.
impressus Broun. C. 1094.
longulus Sharp. C. 1095.
helmsi Sharp. C. 1095.
sulcatus Sharp. C. 1095.
latitans Sharp. C. 1096.
angulatus Broun. C. 1443.
frontalis Broun. C. 1443.
elongellus Broun. C. 1444.
longipes Broun. C. 1444.
depressiusculus White. *Voy. Ereb. and Terr., Ins.* 18.
nitiventris Broun. A. 7, xi, 617.
arcuatus Broun. B. 1, ii, 98.
ocularius Broun. B. 1, ii, 99.
suteri Broun. A. 8, ii, 393.
ruficollis Broun. A. 8, iii, 394.
tenueculus Broun. B. 1, iii, 180.
reversus Broun. T. xliv, 421.
canidus Broun. T. xliv, 421.
mediocris Broun. T. xliii, 99.
parvulus Broun. B. 1, vii, 614.
Pycnomerodes Broun, 1886.
peregrinus Broun. C. 952.

Subfamily CERYLONINAE.

- Philothermus* Aubé, 1843.
nitidus Sharp. C. 211.
sanguineus Broun. C. 211.
notabilis Broun. C. 211.
bicavus Reitter. B. 1, i, 39.

Family HISTERIDAE.

- Platysoma* Leach, 1817.
**baleucelli* Marseul. C. 161.
cinnamomeus White. C. 161.
Sternaulax Marseul, 1862.
zealandicus Marseul. C. 162.
Epiurus Erichson, 1834.
sylvanus Broun. C. 163.
purus Broun. C. 163.
abrogatus Broun. C. 892.
planiceps Broun. C. 892.
crenulatus Broun. C. 892.
simplex Broun. C. 893.
rusticus Broun. C. 893.
rufescens Reitter. B. 1, i, 24.
spinellus Broun. B. 1, vii, 612.
Spininus Erichson, 1834.
pseudocyaneus White. C. 164.
punctulipennis Broun. C. 165.
latipes Broun. C. 666.

- Pachylopus* Erichson, 1834.
pedator Sharp. C. 165.
lepidulus Broun. C. 665.
Abraeus Leach, 1817.
brouni Lewis. C. 166.
vividulus Broun. C. 166.
brunneus Broun. C. 666.
phyllobius Broun. B. 1, ii, 101.
Carcinops.
**14-striata* Stephens.

Family NITIDULIDAE.

- Brachypeplus* Erichson, 1842.
brevicornis Sharp. C. 168.
Epuraea Erichson, 1843.
antarctica White. C. 169.
zealandica Sharp. C. 169.
signata Broun. C. 169.
Homepuraea Broun, 1893.
amoena Broun. C. 171 and 1073.
Nitidula Fabricius, 1775.
lateralis White. C. 170.
abbreviata Fabricius. C. 171.
Soronia Erichson, 1843.
hystrix Sharp. C. 172.
optata Sharp. C. 172.
morosa Broun. C. 1074.
micans Broun. C. 1074.
oculator Reitter. B. 1, i, 24.

- Carpopophilus.*
**hemipterus* Linnaeus.
**mutilatus* Erichson.
Omosita Erichson, 1843.
spinipes Broun. C. 173.
scutellaris Broun. C. 173.
**colon* Linnaeus.
Neocercus Broun, 1919.
electus Broun. B. 1, vi, 523.
Xenoscelis Wollaston, 1864.
prolixus Sharp. C. 174.
Priates Broun, 1881.
optandus Broun. C. 669.
Ips Fabricius, 1776.
minimus Sharp. C. 1078.

- Cyclaryra* Broun, 1893 (substituted for *Cyclomorpha*).
politula Broun. C. 668.
impressa Broun. B. 1, iv, 314.
Platipidia Broun, 1893.
asperella Broun. C. 1075.
Priasilpha Broun, 1893.
obscura Broun. C. 1078.

- Inopria* Broun, 1919.
halli Broun. B. 1, vi, 524.
notata Broun. B. 1, vi, 525.

Subfamily RHIZORHAGINAE.

- Lenax* Sharp, 1877.

- mirandus* Sharp. C. 175.

Family TROGOSITIDAE.

- Trogosita*, Olivier, 1790.
affinis White. C. 177.
**mauritanica* Linnaeus.

Group CLAVICORNES—continued.

- Leperina* Erichson, 1844.
nigrosparsa White. C. 178.
brouni Pascoe. C. 178.
sobrina White. C. 178.
wakefieldi Sharp. C. 179.
ambigua Broun. C. 179.
farinosa Broun. C. 179.
shandi Broun. T. xlvi, 307.
Promanus Sharp, 1877.
depressus Sharp. C. 180.
auripilus Broun. C. 1079.
subcostatus Broun. A. 8, iii, 232.
pulchellus Broun. B. 1, iv, 314.
Grynomus Sharp, 1877.
fusca Sharp. C. 181.
diluta Sharp. C. 181.
viridescens Broun. C. 893.
regularis Sharp. C. 1079.
varians Broun. C. 1340.
rugosa Broun. C. 1341.
albosparsa Broun. A. 8, iii, 231.
setigera Broun. B. 1, v, 387.
clavalis Broun. B. 1, v, 388.
proxima Broun. B. 1, v, 388.
pallidula Broun. B. 1, v, 389.
Phycosecis Pascoe, 1875.
discoidea Pascoe. C. 359.
atomaria Pascoe. C. 359.

Family LATHRIDIIDAE.

- Holoparamecus* Curtis, 1893.
lucidus Broun. C. 232.
castaneus Broun. C. 1448.
tenuis Reitter. B. 1, i, 26.
Lathridius Herbst, 1793.
antipodeus White. C. 233.
sculpturatus Broun. C. 233.
marginalis Broun. C. 233.
costulatus Broun. C. 234.
floridus Broun. C. 234.
bifoveatus Broun. C. 834.
castaneus Broun. C. 834.
proprius Broun. C. 898.
puncticeps Broun. C. 898.
caviceps Broun. C. 1447.
rufifrons Broun. B. 1, iii, 181.
dialis Broun. B. 1, iii, 181.
Corticaria Marsham, 1802.
angusticollis Broun. C. 235.
pubera Broun. C. 235.
fasciata Broun. C. 235.
hirtalis Broun. C. 236.
finitima Broun. C. 236.
variegata Broun. C. 236.
discoidea Broun. C. 236.
obesa Broun. C. 237.
pudibunda Broun. C. 237.
alacer Broun. C. 237.
tarsalis Broun. C. 768.
pacata Broun. C. 899.
erythrocephala Broun. C. 899.
platyptera Broun. C. 899.
semirufa Broun. C. 952.
gilvipes Broun. C. 953.

Corticaria—continued.

- amplipennis* Broun. C. 1104.
clarula Broun. A. 6, xv, 197.
illustris Reitter. B. 1, i, 26.
longula Broun. B. 1, i, 26.
melasoma Broun. B. 1, i, 27.
terricola Broun. A. 6, xii, 189.
picicornis Broun. B. 1, ii, 100.
unicolor Broun. B. 1, iii, 182.
vagepunctata Broun. B. 1, iii, 182.
castanea Broun. B. 1, iii, 183.
robusta Broun. B. 1, iii, 183.
latilippennis Broun. B. 1, iii, 184.
clavatula Broun. B. 1, iii, 185.
fuscicollis Broun. T. xliv, 423.
Rethusus Broun, 1886.
pictulus Broun. C. 835.
lachrymosus Broun. C. 836.
fulvescens Broun. B. 1, vi, 532.
Diarthrocerus Broun, 1893.
formicaephila Broun. C. 1348.
Myrmeccoxenus.
atomarooides Reitter. B. 1, i, 27.
Monotoma.
**picipes* Herbst.
**sub-A-foveolata* Watson.
**spinicollis* Aubé.
Coniciomus.
**nodifer* Westwood.

Family CUCUJIDAE.

- Chaetosoma* Westwood, 1851.
scaritides Westwood. C. 767.
Chaetosomodes Broun, 1919.
halli Broun. B. 1, vii, 615.
Diagrypnodes Waterhouse, 1876.
wakefieldi Waterhouse. C. 217.
Dryocora Pascoe, 1868.
howitti Pascoe. C. 218.
Dendrophagus Schönherr, 1809.
umbrinus Smith. Cat. Coleop. Brit. Mus., 12.
capito Pascoe. C. 219.
Parabrontes Redtenbacher, 1867.
silvanoides Redtenbacher. C. 220.
setiger Broun. C. 768.
Cryptamorpha Wollaston, 1854.
brevicornis White. C. 221.
curvipes Broun. C. 221.
suturalis White. C. 222.
lateritia Broun. C. 222.
rugicollis Broun. B. 1, i, 39.
picturatus Reitter. B. 1, i, 39.
Picrotus Sharp, 1886.
thoracicus Sharp. C. 1098.
sanguineus Broun. C. 1098.
pensus Broun. B. 1, i, 40.
Bronlopiscus Sharp, 1886.
sinuatus Sharp. C. 1099.
pleuralis Sharp. C. 219.
Cathartocryptus Sharp, 1886.
obscurus Sharp. C. 1100.
Saphophagus Sharp, 1886.
minutus Sharp. C. 1101.
ferrugineus Broun. C. 1102.

Group CLAVICORNES—continued.

Tularthrum Broun, 1893.
lineatum Broun. C. 1103.

Thortus Broun, 1893.
ovalis Broun. C. 1345.

Silvanus Latreille.
**surinamensis* Linnaeus.
**unidentatus* Fabricius.

Family CRYPTOPHAGIDAE.

Telmatophilus Heer, 1841.
depressus Sharp. C. 224.
nitens Sharp. C. 224.
olivaceus Broun. C. 1104.
vestitus Broun. B. 1, i, 25.
Cryptophagus Herbst, 1792.
australis Redtenbacher. C. 225.
rubellus Broun. C. 225.
rutilus Broun. C. 226.
vestitus Broun. C. 226.
sylvanus Broun. C. 227.
angulifer Broun. C. 227.
hispidulus Broun. C. 227.
discoideus Broun. C. 1345.
distinctus Broun. C. 1346.
tumidus Broun. C. 1445.
obscurus Broun. C. 1446.
anthracinus Broun. C. 1446.
rufescens Blanchard. *Voy. Pole Sud*, iv, 53, pl. 4.
amoenus Broun. T. xliv, 423.
Saltilius Broun. 1893.
ruficeps Broun. C. 228 and 1347.
Paramecosoma Curtis, 1833.
maculosum Broun. C. 670.
Micrambina.
helmsi Reitter. B. 1, i, 25.
insignis Reitter. B. 1, i, 25.

Family SCAPHIDIIDAE.

Scaphisoma Leach, 1812.
concinnum Broun. C. 158.
scutellare Redtenbacher. C. 159.
tenellum Pascoe. C. 159.
apicellum Broun. C. 160.
actuosum Broun. C. 664.
Baeocera Erichson, 1845.
rufa Broun. C. 665.
rufipes Broun. C. 833.
fulvicolis Broun. C. 891.
armata Broun. C. 891.
sternalis Broun. B. 1, iii, 173.
Brachynopus Broun, 1881.
latus Broun. C. 664.

Family MYCETOPHAGIDAE.

Typhaea Stephens, 1830.
curvipes Broun. C. 238.
hirta Broun. C. 239.
Triphyllus Latreille, 1829.
integratus Broun. C. 1105.
fuliginosus Broun. C. 228.
substratus Broun. C. 229.
aciculatus Broun. C. 229.
serratus Broun. C. 229.
punctulatus Broun. C. 230.

Triphyllus—continued.
adspersus Broun. C. 230.
hispidellus Broun. C. 231.
constans Broun. B. 1, ii, 100.
pubescens Broun. A. 8, iii, 195.

Family DERMESTIDAE.

Trogoderma Latreille, 1892.
serigerum Sharp. C. 240.
signatum Sharp. C. 240.
maestum Broun. C. 241.
granulatum Broun. C. 900.
punctatum Broun. C. 953.
suffusum Broun. C. 953.
fusciatum Broun. C. 1105.
antennale Broun. C. 1106.
puncticollis Broun. B. 1, iii, 187.
pictulum Broun. T. xlvi, 100.
Dermestes, Linnaeus.
** vulpinus*, Fabricius.
** lardarius* Linnaeus.
Anthrenus, Geoffroy.
** musaeorum* Linnaeus.

Family BYRRHIDAE

Nesodendron Latreille.
ovatum Broun. B. 2, 6.
zealandicum Sharp. B. 2, 6.
seriatum Broun. B. 1, vi, 532.
Curimus Erichson, 1846.
squamiger Broun. B. 2, 7.
striatus Broun. B. 2, 7.
vestitus Broun. B. 2, 8.
zeelandicus Redtenbacher. B. 2, 8
Synorthus Broun, 1910.
sternalis Broun. B. 2, 9.
orbicularis Broun. B. 2, 10.
nigricans Broun. B. 2, 10.
insuetus Broun. B. 2, 10.
nigralis Broun. B. 2, 11.
mixtus Broun. B. 2, 11.
mandibularis Broun. B. 2, 12.
laevigatus Broun. B. 2, 12.
setarius Broun. B. 2, 13.
rotundus Broun. B. 2, 13.
villosus Broun. B. 2, 14.
pygmaeus Broun. B. 2, 14.
anomalus Broun. B. 2, 15.
granulatus Broun. B. 2, 15.
substriatus Broun. B. 1, iii, 187.
vesipilus Broun. B. 1, iii, 188.
rectifrons Broun. B. 1, iv, 316.
Pedilophorus Steffahnay, 1843.
foveigerus Broun. B. 2, 16.
gemmeus Broun. B. 2, 17.
laevipennis Broun. B. 2, 17.
probus Broun. B. 2, 17.
puncticeps Broun. B. 2, 18.
tibialis Broun. B. 2, 18.
creperus Broun. B. 2, 19.
puicherrimus Broun. B. 2, 19.
picipes Broun. B. 2, 20.
laetus Broun. B. 2, 20.
lewisi Broun. B. 2, 21.
sculpturatus Broun. B. 2, 21.
cognatus Broun. B. 2, 22

Genus CLAVICORNES—continued.

- Pedilophorus*—continued.
coruscans Pascoe. B. 2, 22.
helmsi Reitter. B. 2, 22.
bryobius Broun. B. 2, 23.
ornatus Broun. B. I, ii, 102.
aemulator Broun. B. I, iii, 189.
nigrescens Broun. B. I, iv, 316.
opaculus Broun. T. xliv, 424.
tumidulus Broun. S. 102.
Liochoria Pascoe, 1875.
huttoni Pascoe. B. 2, 23.
sumptuosa Broun. S. 103.
longula Broun. S. 104.
Cylindrus Broun, 1893.
claviger Broun. B. 2, 24.
Limnichus Latreille, 1829.
decorus Broun. B. 2, 24.
simplex Broun. B. 2, 24.
pictinus Broun. B. 2, 25.
nigripes Broun. B. 2, 25.

Family PARNIDAE.

- Potaminus* Sturm, 1837.
angusticollis Pascoe. C. 248.
Alloparnus Broun, 1893 (substituted for
Parnida).
agrestis Broun. C. 1107 and 249.
scutellaris Broun. B. I, i, 11.
Protoparnus Sharp, 1886.
longulus Sharp. C. 1107.

Family ELMIDAE.

- Hydora* Broun, 1886 (substituted for
Pachycephala).
picea Broun. C. 672.
nitida Broun. C. 927.
obsoleta Broun. C. 928.
vestita Broun. B. I, iii, 153.
lanigera Broun. B. I, iii, 154.
subaenea Broun. B. I, iii, 154.

Group PECTICORNES.

Family LUCANIDAE.

- Lissotes* Westwood, 1855.
reticulatus Westwood. C. 255.
rufipes Sharp. C. 1109.
elegans Broun. C. 1110.
squamidorsis White. C. 251.
punctulatus White. C. 251.
planus Broun. C. 252.
stewarti Broun. C. 673.
capito Deyrolle. E. 1873, 339, and T
 xi, 308.
abditus Broun. C. 673.
demaresti Deyrolle. Ann. Soc. Ent.
 France, 1881, 239.
helmsi Sharp. C. 770.
ithaginus Broun. C. 1108.
aemulus Broun. C. 1109.
acmenus Lewis. T. xxxiv, 203.
menalus Westwood. N.Z. Jour. Sci.,
 ii, 221.
philpotti Broun. B. I, ii, 103.
dispar Broun. T. xlvi, 307.

Ceratognathus Westwood, 1838.

- helotooides* Thomson. C. 254.
alboguttatus Bates. C. 253.
dispar Sharp. C. 1111.
Mitophyllus Parry, 1845.
irroratus Parry. C. 253.
parryanus Hope. C. 253.
marmoratus Watson. Ent. Mo. Mag.,
 xi, 8.
foveolatus Broun. C. 253.
macrocerus Broun. C. 837.
gibbosus Broun. C. 928.
tuberculatus Broun. C. 1111.
cylindricus Broun. A. 6, xv, 199.
angusticeps Broun. A. 6, xv, 199.
fusculus Broun. C. 837.
auriculatus Broun. A. 7, xi, 615.
comognathus Broun. A. 7, xi, 616.
curvifrons Broun. A. 7, xiv, 50.
cristatellus Broun. B. I, v, 391.
reflexus Broun. T. xli, 148.
insignis Broun. B. I, viii, 687.
Dendroblax White, 1836.
earlianus White. C. 251.

Group LAMELLICORNES.

Family COPRIDAE.

- Saphobius* Sharp, 1873.
edwardsi Sharp. C. 256.
wakefieldi Sharp. C. 256.
nitidulus Broun. C. 256.
squamulosus Broun. C. 954.
inflatipes Broun. C. 1112.
fucus Broun. C. 1113.
setosus Sharp. C. 1113.
fulvipes Broun. C. 1114.

Saphobius—continued.

- curvipes* Broun. C. 1349.
tibialis Broun. A. 6, xv, 198.
laticollis Broun. B. I, ii, 104.
lepidus Broun. T. xliv, 425.
Onthophagus Latreille.
**granulatus* Boh.
**posticus* Erichson.
Trox Fabricius.
**sp.*

Group LAMELLICORNES—continued.

Family APHODIIDAE.

- Aphodius* Illiger, 1798.
esculptus White. C. 258.
candens Broun. C. 258.
suspectus Sharp. C. 259.
pascœi Sharp. C. 259.
communis Broun. C. 260.
browni Sharp. C. 260.
fortipes Broun. C. 954.
distant Sharp.
sulcatus Broun. T. xlivi, 101.
**granarius* Linnaeus.
Proctophanes.
**sculptus* Hope.
Phycocnus Broun. 1886.
graniceps Broun. C. 771.
lobatus Broun. C. 1114.

Family MELOLONTHIDAE.

- Stethaspis* Hope, 1837.
suratalis Hope. C. 261.
prasinus Broun. C. 1115.
Costleya Broun, 1886.
discoidea Broun. C. 1116.
simmondsi Broun. T. xliv, 427.
Poecilodiscus Broun, 1895.
pulcher Broun. A. 6, xv, 202.
Scythrodes Broun, 1886.
squalidus Broun. C. 955, and A. 7, xix 59.
Xylostygnum Broun, 1886.
piceus Broun. C. 956.
brookesi Broun. B. 1, vi, 534.
Psilodontria Broun, 1895.
viridescens Broun. A. 6, xv, 201.
Mycernus Broun, 1904.
elegans Broun. A. 7, xiv, 52.
Pyronota Hope, 1837.
festiva Fabricius. C. 262.
laeta Fabricius. C. 262.
edwardsi Sharp. C. 263.
sobrina Sharp. C. 263.
munda Sharp. C. 264.
electa Broun. C. 1116.
regalis Broun. C. 1116.
dives Broun. C. 1117.
aurata Broun. C. 1117.
purpurata Broun. C. 1117.
caerulea Broun. C. 1117.
lugubris Sharp. C. 1118.
pallida Broun. C. 1118.

- Heteronyx* Guerin-Meneville, 1830.
pumilus Sharp. C. 264.
Eusoma White, 1846.
rossii White. C. 268.
costella Broun. C. 269.
aenealis Broun. A. 8, iii, 397.
eximia Broun. B. 1, v, 389.
piliviridis Broun. B. 1, vii, 616.
Sericopilus Sharp, 1882.
advena Sharp. C. 1119.
Prodontria Broun, 1904.
lewisi Broun. A. 7, xiv, 54.
Odontria White, 1846.
striata White. C. 265.
suavis Broun. C. 266.
punctulata Broun. C. 266.
cinnamomea White. C. 267.
xanthosticta White. C. 267.
sandageri Broun. C. 929.
albonotata Broun. C. 1118.
smithii Broun. C. 1450.
piciceps Broun. C. 1450.
epomeas Lewis. T. xxxv, 272.
occipitalis Broun. C. 1451.
marmorata Broun. C. 1451.
fusca Broun. C. 1452.
glabrata Broun. C. 1452.
obscura Broun. A. 6, xv, 203.
zealandica White. C. 270.
brunnea Broun. C. 270.
svylvatica Broun. C. 268.
praelatella Broun. A. 8, iii, 400.
puncticollis Broun. B. 1, iv, 317.
obsolete Broun. B. 1, v, 390.
halli Broun. B. 1, vi, 533.
nitidula Broun. T. xliv, 425.
monticola Broun. T. xliv, 426.
similis Broun. T. xliv, 427.
longitarsis Broun. S. 105.
nesobia Broun. B. 1, vii, 615.
Lewisella, Broun, 1909.
modesta Broun. A. 8, iii, 399.
capito Broun. A. 8, iii, 400.

Family DYNASTIDAE.

- Pericoptus* Burmeister, 1847.
truncatus Fabricius. C. 271.
punctatus White. C. 272.
stupidus Sharp. C. 272.
nitidulus Broun. C. 273.
frontalis Broun. A. 7, xiv, 55.

Group SERRICORNIA.

Family BUPRESTIDAE.

- Nascio* Castelnau and Gory, 1835.
eremita White. C. 276.
enysii Sharp. C. 276.
Buprestis Linnaeus.
**lauta* Leconte.
Stigmodera.
**gulielmi* White.

Family EUCMENIDAE.

- Neocharis* Sharp, 1877.
varia Sharp. C. 277.
pubescens Sharp. C. 278.
simplex Sharp. C. 278.
concolor Sharp. C. 278.
osculans Broun. C. 674.
lobitarsis Broun. B. 1, i, 27.

Group SERRICORNIA—continued.

- Talerax* Sharp, 1877.
distans Sharp. C. 279.
capax Broun. C. 674.
niger Broun. C. 675.
rusticus Broun. C. 675.
foveatus Broun. C. 676.
tenuis Broun. C. 772.
micans Broun. C. 1350.
spinistrasis Broun. B. 1, i, 28.
dorsalis Broun. T. xliv, 428.
Drasterius Eschscholtz, 1829.
nigellus White. C. 279.
Melanus Broun, 1881.
sculptus Broun. C. 677.
Agalba Broun, 1893.
ruficornis Broun. C. 1120.
nigrescens, Broun. C. 1121.
cylindrata Broun. C. 771.
- Family ELATERIDAE.**
- Protelater* Sharp, 1877.
elongatus Sharp. C. 305.
huttoni Sharp. C. 305.
guttatus Sharp. C. 305.
picticornis Sharp. C. 306.
opacus Sharp. C. 306.
nigricans Sharp. C. 772.
vitticollis Broun. C. 956.
atriceps Broun. C. 957.
urquharti Broun. C. 1121.
costiceps Broun. C. 1121.
pubescens Broun. C. 1122.
diversus Broun. T. xliv, 428.
Geranus Sharp, 1877.
collaris Pascoe. C. 302.
lineicollis White. C. 304.
fulvus Sharp. C. 303.
similis Sharp. C. 303.
Parinus Sharp, 1877.
villosus Sharp. C. 301.
Corymbites Latreille, 1834.
antipodus Candèze. C. 299.
dubius Sharp. C. 299.
strangulatus White. C. 300.
myops White. C. 300.
olivasicens White. C. 300.
agriotoides Sharp. C. 301.
irregularis Sharp. C. 1128.
mundus Sharp. C. 1129.
canaliculatus Broun. C. 1129.
munroi Broun. C. 1130.
fulvescens Broun. T. xliv, 429.
vitticollis Broun. T. xliv, 630.
approximans Broun. T. xliv, 430.
sternalis Broun. T. xliv, 431.
Asymphus Sharp, 1886.
insidiosus Sharp. C. 1132.
Agriotes Eschscholtz.
**lineatus* Linnaeus.
Exocelus Broun, 1893.
rufescens Broun. C. 1134.
punctatus Broun. C. 1134.
obsoletus Broun. C. 1135.
- Ochosternus* Candèze, 1863.
realandicus White. C. 298.
parryi Candèze. Mon., iv, 447.
Chrosis Candèze 1863.
polita Sharp. C. 296.
reversa Sharp. C. 297.
barbata Candèze. C. 297.
elongata Sharp. C. 298.
livens Broun. C. 681.
valida Broun. C. 681.
setigera Broun. C. 682.
brevicollis Broun. C. 682.
castanea Broun. C. 683.
certa Broun. C. 683.
fulvipes Broun. C. 683.
impressa Broun. C. 1133.
aeneola Candèze. Mem. Belg., xvii, 1865, 54.
eximia Broun. A. 6, xii, 288.
dubitans Broun. T. xliv, 429.
Cryptohypnus Eschscholtz, 1833.
powelli Sharp. C. 295.
humilis Sharp. C. 295.
frontalis Sharp. C. 295.
longicornis Sharp. C. 296.
thoracicus Sharp. C. 296.
meinertzhageni Broun. C. 680.
montanus Broun. C. 773.
sandageri Broun. C. 774.
pallipes Sharp. C. 1122.
picticornis Broun. C. 1122.
monticola Broun. C. 1124.
sternalis Broun. C. 1124.
basalis Broun. C. 1125.
fuscipennis Broun. C. 1125.
attenuatus Broun. C. 1453.
deterius Broun. C. 1454.
Oxylasme Broun, 1881.
pannosum Broun. C. 679.
tectum Broun. C. 680.
basale Broun. C. 957.
carinale Broun. C. 1130.
vittiger Broun. C. 1452.
Silene Broun, 1893.
brunnea Broun. C. 1136.
Monocrepidius Eschscholtz, 1829.
**exsul* Sharp. C. 294.
subrufus Broun. C. 294.
maritimus Broun. C. 1135.
Mecastrus Sharp, 1877.
convexus Sharp. C. 293.
vicinus Sharp. C. 293.
discedens Sharp. C. 293.
lateristrigatus White. C. 306.
intermedius Broun. C. 1128.
Zeuglophus Broun, 1895.
pilicornis Broun. A. 6, xv, 235.
Lomemus Sharp, 1877.
pilicornis Sharp. C. 290.
pictus Sharp. C. 290.
suffusus, Sharp. C. 291.
flavipes Sharp. C. 291.
similis Sharp. C. 291.
obscuripes, Sharp. C. 292.

Group SERRICORNIA—continued.

- Lomemus*—continued.
elegans Sharp. C. 292.
collaris Sharp. C. 292.
rectus Broun. C. 774.
vittatus Broun. C. 775.
fulvipennis Broun. C. 1126.
fuscicornis Broun. C. 1126.
maurus Broun. C. 1127.
sculpturatus Broun. C. 1127.
frontalis Broun. C. 1128.
fusciipes Broun. C. 1454.
puncticollis Broun. A. 6, xv, 234.
vittipennis Broun. B. 1, i, 29.
- Lacon.*
**variabilis* Candèze.
Psorochroa Broun, 1886.
granulata Broun. C. 773.
- Aglophus* Sharp, 1877.
modestus Sharp. C. 289.
- Panspoeus* Sharp, 1877.
guttatus Sharp. C. 288.
tenebrosus Broun. C. 288.
- Betarmon* Kiesenwetter, 1858.
gracilipes Sharp. C. 287.
frontalis Sharp. C. 287.
laetus Sharp. C. 287.
obscurus Sharp. C. 287.
flavipilus Broun. C. 1350.
- Amphiplatys* Sharp, 1877.
lawsoni Sharp. C. 286.
- Metablaes* Sharp, 1877.
brouni Sharp. C. 284.
acutipennis White. C. 284.
approximans White. C. 285.
cinctiger White. C. 285.
- Thoramus* Sharp, 1877.
wakefieldi Sharp. C. 280.
rugipennis Broun. C. 281.
feredayi Sharp. C. 282.
laevithorax White. C. 282.
per blandus Broun. C. 282.
foveolatus Broun. C. 283.
cervinus Broun. C. 677.
angustus Broun. C. 678.
parvulus Broun. C. 678.
huttoni Sharp. C. 1122.
- GENERA OF DOUBTFUL POSITION.**
- Brounia* Sharp, 1878.
thoracica Sharp. C. 274.
- Byrrhocryptus* Broun, 1893.
urquharti Broun. C. 1137.
- Amplectopus* Sharp, 1886.
ovalis Sharp. C. 1138.
latulus Broun. C. 1351.
fuscus Broun. C. 1351.
pallicornis Broun. C. 1456.
- Amychus.*
candezei Pascoe. T. ix, 416.
- Family DASCILLIDAE.**
- Byrrhodes* Sharp, 1878.
gravidus Sharp. C. 307.
- Cyprobius* Sharp, 1878.
nitidus Sharp. C. 308.
undulatus Broun. C. 775.
terrenus Broun. A. 6, xv, 236.
- Cyphanus* Sharp, 1878.
laticeps Sharp. C. 309.
punctatus Sharp. C. 309.
capax Broun. C. 309.
mollis Sharp. C. 310.
debilis Sharp. C. 310.
granulatus Broun. C. 311.
medius Broun. C. 311.
mandibularis Broun. C. 776.
dubius Broun. C. 1139.
scaber Broun. C. 1139.
ocularius Broun. B. 1, i, 29.
maculifer Broun. B. 1, i, 29.
granulosus Broun. B. 1, i, 30.
- Atopida* White, 1846.
lawsoni Sharp. C. 312.
suturalis White. C. 338.
brouni Sharp. C. 312.
prota Sharp. C. 313.
castanea White. C. 313.
hirta Broun. C. 313.
testacea Broun. C. 314.
dorsalis Broun. C. 1140.
sinuata Broun. C. 1455.
grahami Broun. B. 1, i, 31.
suffusa Broun. B. 1, i, 31.
impressa Broun. B. 1, ii, 105.
villosa Broun. B. 1, vi, 534.
pallidula Broun. B. 1, vi, 535.
montana Broun. B. 1, vi, 536.
basalis Broun. T. xliv, 431.
- Veronatus* Sharp, 1878.
longicornis Sharp. C. 314.
longipalpis Sharp. C. 315.
sharpae Broun. C. 315.
frontalis Broun. C. 316.
scabiosus Broun. C. 316.
capito Broun. C. 316.
antennalis Broun. C. 776.
nubilus Broun. C. 1455.
amplus Broun. A. 6, xv, 237.
granicollis Broun. B. 1, i, 32.
punctipennis Broun. B. 1, ii, 105.
tarsalis Broun. B. 1, iv, 318.
fulgidulus Broun. B. 1, iv, 318.
sternalis Broun. B. 1, vi, 536.
reversus Broun. B. 1, vi, 537.
apterus Broun. B. 1, vi, 537.
brevicollis Broun. B. 1, vi, 538.
versicolor Broun. B. 1, vi, 539.
vestitus Broun. B. 1, vi, 539.
- Mesocypion* Sharp, 1878.
marmoratus Sharp. C. 317.
setiger Sharp. C. 317.
wakefieldi Sharp. C. 318.
divergens Sharp. C. 318.
monticola Broun. C. 958.
granulatus Broun. C. 958.
pallidus Broun. C. 1141.
laticeps Broun. C. 1142.

Group SERRICORNIA—continued.

Mesocypphon—continued.

- tristis* Broun. B. I, i, 33.
capito Broun. B. I, ii, 106.
vestitus Broun. B. I, ii, 106.
longicornis Broun. B. I, ii, 107.
bifoveatus Broun. B. I, iii, 189.
lateralis Broun. B. I, iii, 190.
mandibularis Broun. T. xliv, 432.

Cyphon Paykull, 1798.

- huttoni* Sharp. C. 320.
parviceps Sharp. C. 320.
pumilio Sharp. C. 320.
arduus Sharp. C. 320.
oscillans Sharp. C. 321.
aequalis Sharp. C. 321.
amplus Broun. C. 322.
viridipennis Broun. C. 322.
graniger Sharp. C. 322.
pictulus Sharp. C. 322.
zealandicus Sharp. C. 323.
suffusus Sharp. C. 323.
laticeps Sharp. C. 323.
genalis Sharp. C. 324.
remotus Broun. C. 776.
nigropictus Broun. C. 777.
laticollis Broun. C. 777.
rectangulus Broun. C. 777.
plagiatus Broun. C. 778.
propinquus Broun. C. 778.
acerbus Broun. C. 778.
crassus Broun. C. 779.
suburalis Broun. C. 779.
nitidus Broun. C. 779.
lateralis Broun. C. 780.
molestus Broun. C. 780.
dilutus Broun. C. 780.
rectalis Broun. C. 837.
waikatoensis Broun. C. 838.
signatus Broun. C. 958.
aethiops Broun. C. 959.
trivialis Broun. C. 959.
pauper Broun. C. 1143.
discedens Broun. C. 1143.
flavescens Broun. C. 1143.
nigrifrons Broun. C. 1144.
fuscifrons Broun. C. 1144.
fulvicornis Broun. C. 1144.
ornatus Broun. C. 1145.
variegatus Sharp. C. 1145.

- mackerowii* Broun. A. 6, xv, 237.
deterius Broun. B. I, ii, 107.
cincticollis Broun. B. I, iv, 319.
granulicollis Broun. B. I, iv, 320.
plumatellus Broun. B. I, iv, 320.
burrowsi Broun. B. I, iv, 320.
pachymerus Broun. T. xliv, 432.

Metriorhynchus.

- rufipennis*
- Fabricius.

Cyphotelus Sharp, 1878.

- angustifrons*
- Sharp. C. 325.

Family CLERIDAE.

Paupris Sharp, 1877.

- aptera*
- Sharp. C. 331.

- Metaxina* Broun, 1909.
ornata Broun. A. 8, iii, 408.
Parmius Sharp, 1877.
longipes Sharp. C. 331.
debilis Sharp. C. 332.
rugosus Broun. C. 1148.
violaceus Broun. T. xliv, 434.
Pelonium Spinola, 1844.
pustuliferum Westwood. P.Z.S., 1852,
 52, pl. 24, fig. 8.
Balcus Sharp, 1877.
niger Sharp. C. 333.
signatus Broun. C. 333.
Eurnede Pascoe, 1876.
aeraria Pascoe. C. 334.
Phymatophaea Pascoe, 1876.
electa Pascoe. C. 334.
violacea Fabricius. C. 335.
hilaris Sharp. C. 335.
pantomelas Boisduval. C. 335.
longula Sharp. C. 336.
opiloides Pascoe. C. 336.
fulvipalpis Broun. C. 336.
picta Broun. C. 685.
dorsalis Broun. C. 685.
viridans Broun. C. 686.
testacea Broun. C. 686.
atrata Broun. C. 686.
ignea Broun. C. 784.
opacula Broun. C. 1352.
abnormis Broun. B. I, i, 37.
sculptipennis Broun. B. I, ii, 108.
fusciatris Broun. B. I, ii, 109.
brevisetula Broun. B. I, ii, 109.
lugubris Broun. A. 8, iii, 405.
apicale Broun. A. 8, iii, 406.
o'connori Broun. B. I, iii, 194.
griseipennis Broun. T. xliv, 433.
hudsoni Broun. B. I, viii, 688.
nigricornis Broun. B. I, viii, 689.
brevicollis Broun. B. I, viii, 689.
formosa Broun. B. I, viii, 689.
Mathesis Waterhouse, 1877.
guttigera Waterhouse. E. 1877, 8.
Necrobia Oliver.
**ruficollis*, Fabricius.
**rufipes* De Geer.

Family MELYRIDAE.

- Dasytes* Paykull, 1798.
subcyanus Broun. C. 328.
wakefieldi Sharp. C. 329.
cinerohirtus Broun. C. 329.
obscuricollis Broun. C. 330.
laticeps Broun. C. 330.
stewarti Broun. C. 684.
constrictus Broun. C. 781.
occipitalis Broun. C. 782.
cheesemani Broun. C. 902.
opaculus Broun. C. 202.
helmsi Sharp. C. 1147.
oreocharis Broun. C. 1148.
littoralis Broun. C. 1351.
aethiops Broun. C. 1352.

Group SERRICORNIA—continued.

Dasytes—continued.

- nigripes* Broun. A. 6, xv, 238.
veronicae Broun. B. 1, i, 36.
laevifrons Broun. B. 1, ii, 109.
aurisetifer Broun. A. 8, iii, 404.
anacharis Broun. A. 8, iii, 405.
fuscularis Broun. B. 1, iii, 192.
philotti Broun. B. 1, iv, 321.
pittensis Broun. T. xlvi, 101.
violascens Broun. B. 1, vii, 617.
clavatus Broun. B. 1, vii, 617.

Arthracanthus Broun, 1886.

- planifrons* Broun. C. 781.
fulvipes Broun. B. 1, iii, 192.
fossicollis Broun. B. 1, iii, 193.
atriceps Broun. B. 1, iii, 193.
foveicollis Broun. T. xlvi, 433.

Hyles Broun, 1886.

- nigrescens* Broun. C. 783.
brevicornis Broun. C. 783.
semidilutus Broun. C. 783.

Family TELEPHORIDAE.

Asilis Broun, 1893.

- fulvithorax* Broun. C. 326.
nigriceps Broun. C. 327.
subnudus Broun. C. 327.
tenuiculus Broun. C. 327.
striatus Broun. C. 328.
tumidus Broun. C. 684.
piliventer Broun. C. 684.
laevigatus Broun. C. 901.
coloratus Broun. C. 1147.
collaris Broun. B. 1, i, 33.
brevicornis Broun. B. 1, i, 34.
opaculus Broun. B. 1, i, 34.
egmontensis Broun. B. 1, i, 35.
punctipennis Broun. B. 1, i, 35.
parallelus Broun. B. 1, i, 36.
plicornis Broun. A. 8, iii, 401.
sinuellus Broun. A. 8, iii, 402.
granipennis Broun. A. 8, iii, 402.
interstitialis Broun. A. 8, iii, 403.
apicalis Broun. A. 8, iii, 404.
flavipennis Broun. B. 1, iii, 191.

Family PTINIDAE.

Subfamily PTININAE.

- Ptinus* Linnaeus, 1767.
speciosus Broun. C. 338.
marinus White. C. 338.
pilosus White. C. 338.
littoralis Broun. C. 1353.
plagiatus Broun. B. 1, iii, 194.
**fur* Linnaeus.
**tectus* Boisduval.

Subfamily ANOBINAE.

- Anobium* Fabricius, 1775.
amplicolle Broun. C. 339.

Anobium—continued.

- ruficorne* Broun. C. 340.
tricostatum White. C. 341.
undulatum Broun. C. 687.

inaequale Broun. T. xliv, 434.

niticole Broun. T. xliv, 435.

**tesselatum* Linnaeus.

**domesticum* Linnaeus.

**paniceum* Linnaeus.

Xenogonus Broun, 1882.

notatus Broun. C. 340.

granulatus Broun. C. 341.

sericeus Broun. C. 341.

pullus Broun. C. 688.

furcatus Broun. C. 689.

versutus Broun. C. 689.

plagiatus Broun. C. 689.

ambiguus Broun. C. 689.

Ochina Sturm, 1837.

vulgaris Broun. C. 342.

Sphindoteles Broun, 1881.

atriventris Broun. C. 687.

Methemus Broun, 1903.

griseipilus Broun (*Capnodes*). C. 690.

Dorcatoma Herbst, 1790.

oblonga Broun. C. 343.

illustris Broun. C. 343.

lata Broun. C. 690.

Euderia Broun, 1880.

squamosa Broun. C. 344.

Family BOSTRICHIDAE.

Apate Fabricius, 1775.

minuta Fabricius. C. 346.

Lyctus.

**brunneus* Stephens.

Family CIOIDAE.

Cis Latreille, 1796.

undulatus Broun. C. 347.

assimilis Broun. C. 347.

rufulus Broun. C. 348.

lineicollis Broun. C. 348.

flavitarsis Broun. C. 348.

asperimus Broun. C. 349.

cornuticeps Broun. C. 349.

illustris Broun. C. 349.

perpinguis Broun. C. 350.

anthracinus Broun. C. 350.

picturatus Broun. C. 358.

obesulus Broun. C. 839.

recurvatus Broun. C. 784.

picicollis Broun. C. 785.

viridiflavus Broun. C. 785.

fultoni Broun. C. 904.

piciceps Broun. C. 904.

zeelandicus Reitter. B. 1, i, 41.

Ennearthron Mellie.

boettgeri Reitter. B. 1, i, 41.

obsoletum Reitter. B. 1, i, 41.

Group LONGICORNES.

Family PRIONIDAE.

- Prionoplus* White, 1846.
reticularis White. C. 566.
Ochrocydus Pascoe, 1876.
huttoni Pascoe. C. 567.

Family CERAMBYCIDAE.

- Liogramma* Bates, 1874.
zealandica Blanchard. C. 568.
Didymocantha Newman, 1840.
sublineata White. C. 568.
diversicornis White. C. 569.
picta Bates. C. 569.
aegrota Bates. C. 569.
brevicornis Broun. C. 570.
clavipes Broun. C. 808.
robusta Sharp. C. 1271.
quadriguttata Sharp. C. 1272.
fucunda Broun. C. 1272.
binotata Broun. C. 1273.
vittata Broun. C. 1273.
pallida Broun. A. 6, xii, 388.
cognata Broun. B. 1, i, 73.
hudsoni Broun. B. 1, ii, 140.
media Broun. T. xlvi, 152.
oedemera Broun. T. xlvi, 153.
fuscolobus Broun. T. xlvi, 153.
Aemonia Newman, 1842.
hirta Fabricius. C. 570 and 1275.
villosa Fabricius. C. 1275.
simplicollis Broun. C. 571 and 1276.
humilis Newman. C. 1274.
inaequalis Sharp. C. 1275.
mutica Sharp. C. 1276.
plicicollis Sharp. C. 1276.
debilis Sharp. C. 1277.
sublineata Broun. A. 8, iv, 281.
separata Broun. B. 1, vii, 659.
Leptachinus Bates, 1874.
strigipennis Westwood. C. 752.
Votum Broun, 1880.
mundum Broun. C. 753.
Pseudocalliprason Broun, 1880.
marginatum White. C. 574.
Ambeodontus.
tristis Fabricius. C. 574.
Agapanthida White, 1846.
pulchella White. C. 575.
scutellaris Pascoe. C. 575.
Ophryops White, 1846.
pallidus White. C. 576.
dispar Sharp. C. 1277.
nigropictus Broun. C. 1278.
testaceus Broun. C. 1278.
lentiginosus Broun. C. 1279.
Pseudosemnus Broun, 1893.
amabilis Broun. C. 1280.
Hesperophanes Mulsant, 1839.
unicolor Fabricius. *Mantissa Insectorum*, 147.
Astetholea Bates, 1874.
pauper Bates. C. 577.
lepturoides Bates. C. 577.
aubreyi Broun. C. 577.
Astetholida Broun, 1880.
lucida Broun. C. 578.

- Epheus* Broun, 1886.
costifer Broun. C. 871.
Blosyropus Redtenbacher, 1868.
spinosis Redtenbacher. C. 579.
Ptychopterus Broun, 1886.
rugosus Broun. C. 809.
Psilotomorpha Saunders, 1850.
tenuipes Saunders. C. 580.
Xuthodes Pascoe, 1875.
punctipennis Pascoe. C. 580.
apicalis Sharp. C. 580.
batesi Sharp. C. 581.
divergens Broun. C. 581.
lepidus Broun. C. 1280.
Calliprason White, 1846.
sinclairi White. C. 582.
Stenopotes Pascoe, 1875.
pallidus Pascoe. C. 583.
Cacodrotus Broun, 1893.
bifasciatus Broun. C. 1281.
Drotus Sharp, 1877.
elegans Sharp. C. 583.
Drototelus Broun, 1903.
politus Broun. A. 7, xii, 82.
Zorion Pascoe, 1867.
minutum Fabricius. C. 584.
guttigerum Westwood. C. 585.
castum Broun. C. 1281.
opacum Sharp. T. ali, 150.
Gnomodes Broun, 1893.
piceus Broun. C. 1282.
Gastrosarus Bates, 1874.
nigricollis Bates. C. 586.
urbanus Broun. C. 1283.
lautus Broun. C. 1283.
picticornis Broun. C. 1284.
Elurida Broun, 1880.
sericea Smith. C. 587.
Acrocyrta Pascoe, 1856.
spinicornis Newman. C. 587.
Clytus Laichorting, 1784.
rugulosus Broun. C. 588.
Coptomma Newman, 1840.
variegatum Fabricius. C. 589.
Nasomorpha Blanchard, 1853.
lineatum Fabricius. C. 590.
sulcatum Fabricius. C. 590.
neglectum Broun. C. 591.
sticticum Broun. C. 1284.
Ceresium Newman, 1842.
zealandicum Blanchard. *Voy. Pole Sud*, iv, 272.
Aphanasius Thomson, 1860.
australis Boisduval. *Voy. Astrolabe*, ii, 480.
Phlyctaenodes Newman, 1840.
retifer Lacordaire. *Gen Col.*, 1869, 374.
Hylotrupes.
**bajulus* Linnaeus.
Phoracantha.
**recurrens* Newman.
Callirhoe.
**allaspa* Newman.
Tessaromma.
**undatum* Newman.
**sulcatum*.

Group LONGICORNES—continued.

Family LAMIIDAE.

- Hexatricha* Lacordaire, 1869.
pulverulenta Westwood. C. 592.
heteromorpha Boisduval, *Voy. Astrolabe*,
 ii, 21.
Xylotoles Newman, 1840.
lynceus Fabricius. C. 593.
griseus Westwood. C. 593
humeralis Bates. C. 593.
subpinguis White. C. 594.
nudus Bates. C. 594.
rugicollis Bates. C. 594.
inornatus Broun. C. 595.
laetus White. C. 595.
gratus Broun. C. 595.
nanus Bates. C. 596.
aegrotus Bates. C. 597.
pulchellus Bates. C. 597.
scissicauda Bates. C. 597.
bullatus Sharp. C. 598.
pictulus Bates. C. 598.
sandageri Broun. C. 912.
maculosus Broun. C. 913.
angustulus Broun. C. 973.
buttoni Sharp. C. 1287.
abnormalis Sharp. T. xli, 150.
gaudens Broun. C. 1288.
germanus Sharp. C. 1289.
traversi Pascoe. C. 599.
fasciatus Sharp. C. 1289.
phormiobius Broun. C. 1388.
prolongatus Broun. C. 1389.
costatus Pascoe. C. 599.
parvulus White. *Voy. Ereb. and Terror*,
 Insects, 22.
gracilis Broun. B. 1, i, 75.
schaubinslandi Sharp. T. xli, 150.
submicans Broun. B. 1, vii, 659.
apicalis Broun. B. 1, viii, 705.
pygmaeus Broun. B. 1, viii, 705.
Microlamia Bates, 1874.
pygmaea Bates. C. 599.
aemula Broun. C. 810.
Mesolamia Sharp, 1882.
marmorata Sharp. C. 1295.
aerata Broun. C. 1296.
Somatidia Thomson, 1864.
antarctica White. C. 600.
assimilata Broun. C. 600.
terrestris Broun. C. 601.
ptinoides Bates. C. 601.
angusta Broun. C. 601.
pictipes Broun. C. 602.
longipes Sharp. C. 602.
diversa Broun. C. 603.
nitida Broun. C. 603.
crassipes Broun. C. 810.
elongata Broun. C. 811.
grandis Broun. C. 1290.
helmsi Sharp. C. 1290.
spinicollis Broun. C. 1291.
fulvipes Broun. B. 1, viii, 706.
costifer Broun. C. 1291.
variegata Broun. C. 1292.

- Somatidia*—continued.
simplex Broun. C. 1292.
signata Broun. C. 1293.
convexa Broun. C. 1498.
latula Broun. C. 1499.
laevior Broun. C. 1499.
tenebria Broun. C. 1500.
albicomata Broun. C. 1500.
proxima Broun. A. 6, xii, 390.
picticornis Broun. A. 6, xv, 418.
testudo Broun. A. 7, xiv, 127.
websteriana Broun. A. 8, iv, 281.
heterothra Broun. A. 8, iv, 282.
crassicollis Broun. B. 1, viii, 706.
testacea Broun. A. 8, iv, 283.
maculata Broun. B. 1, vi, 587.
sericophora Broun. A. 8, iv, 283.
lineifera Broun. A. 8, iv, 284.
rubella Broun. B. 1, iii, 254.
suturalis Broun. B. 1, iii, 255.
ruficornis Broun. B. 1, iii, 255.
halli Broun. B. 1, iii, 256;
waitei Broun. T. xlvi, 113.
vicina Broun. T. xlvi, 114.
spectabilis Broun. B. 1, v, 456.
flavidorsis Broun. B. 1, v, 457.
suffusa Broun. B. 1, v, 457.
femoralis Broun. B. 1, v, 458.
oscillans Broun. B. 1, v, 458.
obesula Broun. B. 1, v, 459.
laevinotata Broun. B. 1, v, 460.
parvula Broun. B. 1, v, 461.
longula Broun. B. 1, vi, 588.
vittigera Broun. B. 1, vi, 588.
pennulata Broun. B. 1, vi, 589.
thoracica Broun. T. xlvi, 154.
nodularia Broun. T. xlvi, 155.
discoidea Broun. T. xlvi, 155.
posticalis Broun. T. xlvi, 156.
corticola Broun. T. xlvi, 156.
pinguis Broun. T. xlvi, 157.
placita Broun. B. 1, vii, 660.
origana Broun. B. 1, vii, 661.
oedemera Broun. B. 1, vii, 661.
o'connori Broun. B. 1, vii, 662.
commoda Broun. B. 1, vii, 662.
Stenellipasis, Bates, 1874,
bimaculata White. C. 604.
gracilis White. C. 604.
latipennis Bates. C. 605.
pumila Pascoe. C. 605.
cuneata Sharp. C. 1289.
Psilocnæia Bates, 1874.
linearis Bates. C. 606.
brouni Bates. C. 606.
Spilotrogia Bates, 1874.
maculata Bates. C. 607.
kilarula Broun. C. 607.
Eurychaena Bates, 1874.
fragilis Bates. C. 608.
feredai Bates. C. 608.
acutula Broun. C. 608.
Tetrorea White, 1846.
ciliipes White. C. 609.

Group LONGICORNES—continued.

Tetrorea—continued.
discedens Sharp. C. 1294.
longipennis Sharp. C. 1293.
sellata Sharp. C. 1294.
maculata Broun. T. xlvi, 157.
Hybolasius Bates, 1874.
cristus Fabricius. C. 610.
lanipes Sharp. C. 610.
pedator Sharp. C. 610.
wakefieldi Bates. C. 611.
concolor Broun. C. 611.
viridescens Bates. C. 611.
promissus Broun. C. 612.
simplex Bates. C. 612.
modestus Broun. C. 613.
variegatus Broun. C. 613.
cristatellus Bates. C. 614.
bellicosus Broun. C. 614.
parvus Broun. C. 614.
vegetus Broun. C. 744.
fasciatus Broun. C. 744.
brevicollis Broun. C. 811.
pictaris Broun. C. 812.
pusillus Broun. C. 812.
apicalis Broun. C. 872.
piceus Broun. C. 872.
deplanatus Sharp. C. 1297.
castaneus Broun. C. 1297.
finitimus Broun. C. 1298.
abshirtus Broun. C. 1298.
gnarus Broun. C. 1298.

Hybolasius—continued.
femoralis Broun. C. 1299.
thoracicus Broun. C. 1299.
rufescens Broun. C. 1300.
dubius Broun. C. 1389.
optatus Broun. A. 6, xii, 389.
trigonellaris Hutton. T. xxx, 158.
cognatus Broun. A. 7, xii, 82.
laticollis Broun. A. 7, xi, 83.
gracilipes Broun. A. 7, xii, 84.
genalis Broun. A. 7, xii, 84.
varipes Broun. A. 8, iv, 284.
ciliatus Broun. B. 1, ii, 140.
vittiger Broun. B. 1, ii, 141.
lineiceps Broun. B. 1, iii, 257.
sculpturatus Broun. B. 1, iv, 341.
cupiendus Broun. T. xlvi, 158.
tumidulus Broun. T. xlvi, 159.
rugicollis Broun. T. xlvi, 159.
Poecilippe Bates, 1874.
stictica Bates. C. 615.
flavipes White. C. 615.
medialis Sharp. C. 1301.
femoralis Sharp. C. 1301.
simplex Bates. C. 612.
Diastamerus Redtenbacher, 1868.
tomentosus Redtenbacher. C. 616.
Tympanopalpus Redtenbacher, 1868.
dorsalis Redtenbacher. C. 617.
Adriopea Broun. 1910.
pallidata Broun. B. 1, i, 74.

Group PHYTOPHAGA.

Family CHRYSOMELIDAE.

Subfamily EUMOLFINAE.

Eucolaspis Sharp, 1886.
brunneus Fabricius. C. 622.
pallidipennis White. C. 622.
puncticollis Broun. C. 623.
jucundus Broun. C. 623.
subaeneus Broun. C. 624.
plicatus Broun. A. 8, iv, 285.
sculptus Broun. C. 624.
merus Broun. C. 624.
brevicollis Broun. C. 625.
atroceruleus Broun. C. 625.
ochraceus Broun. C. 1303.
coloratus Broun. C. 1303.
montanus Broun. C. 1304.
vittiger Broun. A. 6, xii, 391.
picticornis Broun. A. 6, xii, 391.
Atrichatus Sharp, 1886.
ochraceus Broun. C. 625 and 1305.
aeneicollis Broun. A. 6, xv, 419.
nitidulus Broun. A. 8, iv, 286.
Pilacolaspis Sharp, 1886.
wakefieldi Sharp. C. 1306.
huttoni Broun. C. 626.
rugiventris Broun. B. 1, iii, 258.
angulatus Broun. T. xlvi, 160.
latipennis Broun. T. xlvi, 160.

Peniticus Sharp, 1876.
suffusus Sharp. C. 627.
antiquus Sharp. C. 627.
robustus Broun. C. 628.
plicatus Broun. B. 1, vii, 663.

Subfamily CRYPTOCEPHALINAE.

Arnomus Sharp, 1876.
browni Sharp. C. 619.
curtipes Broun. C. 1390.
marginalis Broun. C. 1390.
viridicollis Broun. A. 8, iv, 286.
signatus Broun. A. 8, iv, 287.
fulvus Broun. B. 1, iv, 342.
vicus Broun. B. 1, iv, 342.
Alema Sharp, 1876.
paradoxa Sharp. C. 620.
puncticollis Broun. C. 620.
spatiosa Broun. C. 621.
Eualema Broun, 1903.
walkeri Broun. A. 7, xii, 86.
Scaphodius Chapuis, 1874.
compactus Sharp. C. 1302.

Subfamily CHRYSOMELINAE.

Allocharis Sharp, 1882.
marginata Sharp. C. 1306.
morosa Broun. C. 1307.

Group PHYTOPHAGA—continued.

Allocharis—continued.

- limbata* Broun. C. 1307.
praestans Broun. B. 1, v, 462.
subscutata Broun. B. 1, v, 462.
nigricollis Broun. B. 1, v, 463.
picticornis Broun. B. 1, v, 463.
media Broun. B. 1, v, 463.
fuscinervis Broun. B. 1, v, 464.
robusta Broun. B. 1, v, 464.
tarsalis Broun. B. 1, v, 465.

Chalcolumpra.

- speculifera*
- Sharp. B. 1, i, 77.

Aphilon Sharp, 1876.

- enigma* Sharp. C. 629.
pretiosa Broun. C. 630.
punctata Broun. C. 630.
minuta Broun. C. 631.
monstrosa Broun. C. 874.
sobrina Broun. C. 875.
praestans Broun. C. 1309.
convexa Broun. C. 1310.
latulum Broun. C. 1310.
scutellaris Broun. C. 1501.
laticollis Broun. A. 6, xii, 392.
impressa Broun. B. 1, iii, 258.
sternalis Broun. B. 1, vii, 664.

Caccomolpus Sharp, 1886.

- globosus* Sharp. C. 1308.
plagiatus Sharp. C. 1308.
pullatus Broun. C. 1309.
maculatus Broun. C. 1309.
ornatus Broun. B. 1, i, 77.
flectipes Broun. B. 1, iii, 259.
nigristernis Broun. B. 1, v, 465.
hallianus Broun. B. 1, v, 466.
fuscicornis Broun. B. 1, v, 466.
substriatus Broun. B. 1, v, 467.
tibialis Broun. B. 1, v, 467.
viridescens Broun. B. 1, v, 468.
subcypreus Broun. B. 1, vi, 589.
amplus Broun. B. 1, vi, 590.
cinctiger Broun. B. 1, vi, 590.
montanus Broun. B. 1, vii, 665.
Cyrtonegetus Broun, 1915.
crassus Broun. B. 1, iv, 343.

Subfamily GALERUCINAE.

Luperus Geoffroy, 1764.

- vulgaris* Broun. C. 632.
attenuatus Broun. C. 633.
thoracicus Broun. C. 633.
nigripes Broun. C. 634.
nigricans Broun. C. 634.
nitidicollis Broun. C. 634.
aeneus Broun. C. 635.
viridis Broun. C. 635.
oleareae Broun. C. 1312.
rugicollis Broun. C. 1313.
brevicollis Broun. C. 1313.
suicifer Broun. C. 1313.
rectipes Broun. C. 1314.
calcarius Broun. C. 1314.
princeps Broun. C. 1314.
monticola Broun. C. 1315.
fuscatus Broun. C. 1315.
sordidus Broun. C. 1316.
truncatus Broun. C. 1316.
nigricornis Sharp. C. 1316.

Luperus—continued.

- aenescens* Sharp. C. 1317.
puncticollis Sharp. C. 1317.
mollis Broun. C. 1502.
cheesemani Broun. B. 1, i, 75.
obscurus Broun. B. 1, i, 76.
diversus Broun. B. 1, i, 76.
angularis Broun. A. 8, iv, 288.
scutellaris Broun. A. 8, iv, 289.
lewisi Broun. A. 8, iv, 289.
axyrocharis Broun. A. 8, iv, 290.
palialis Broun. A. 8, iv, 290.
asperellus Broun. A. 8, iv, 290.
insolitus Broun. B. 1, iii, 260.
dilutipes Broun. B. 1, iv, 344.
anthracinus Broun. B. 1, iii, 260.
bullatus Broun. B. 1, iii, 261.
discrepans Broun. B. 1, iii, 261.
dilatatus Broun. B. 1, iii, 262.
iridescentis Broun. B. 1, iii, 263.
aurellus Broun. B. 1, iii, 263.
nodicollis Broun. B. 1, iv, 344.
pubicollis Broun. B. 1, iv, 345.
halli Broun. B. 1, v, 469.
dilucidus Broun. B. 1, v, 470.
cyanescens Broun. B. 1, v, 470.
gracilipes Broun. B. 1, v, 471.
minor Broun. B. 1, v, 471.
quadricollis Broun. B. 1, v, 472.
xenoscelis Broun. B. 1, v, 472.
perplexus Broun. B. 1, v, 473.
pygidialis Broun. B. 1, v, 473.
medioiris Broun. B. 1, v, 474.
simmondsi Broun. T. xlvi, 161.
foveigerus Broun. T. xlvi 161.
o'connori Broun. T. xlvi, 162.
atribennis Broun. T. xlvi, 162.
Allastena Broun, 1893.
nitida Broun. C. 1318.
quadrata Broun. C. 1319.
piliventris Broun. B. 1, iv, 346.
eminens Broun. B. 1, v, 468.

Bryobates Broun, 1886.

- coniformis* Broun. C. 874.
aeratus Broun. B. 1, ii, 141.
nigricans Broun. B. 1, ii, 142.
rugidorsis Broun. B. 1, v, 461.

Subfamily HALTICINAE.

Trachytetra Sharp, 1886.

- rugulosa* Broun. C. 636 and 1311.
frontalis Broun. B. 1, viii, 707.
robusta Broun. B. 1, viii, 707.

Inopelonia Broun, 1893.

- testacea* Broun. C. 637.
fuliginosa Broun. C. 637.

Pleuraltica Sharp, 1886.

- cyanea* Broun. C. 1312 and 638.

Phyllotreta Chevrolat, 1839.

- nitida* Broun. C. 636.
graminicola Broun. C. 1391.
littoralis Broun. C. 1391.
vittigera Broun. C. 1392.

- passpala* Broun. B. 1, viii, 708.

Family BRUCHIDAE.

Bruchus.**rufimanus* Bohem.

Group HETEROMERA.

Family TENEBRIONIDAE.

Subfamily OPATRINAE.

- Opatrum* Fabricius, 1775.
laevigatum Fabricius. *Ent. Syst.*, i, 89, 5.
Pseudopatrum Sharp, 1886.
tuberculostatum White. C. 353.
sordidum Sharp. C. 1152.
Mesopatrum Broun, 1893.
granulosum Broun. C. 1355.
dubium Broun. B. 1, v, 396.
Periatrum Sharp, 1886.
helmsi Sharp. C. 1153.
tumipes Broun. C. 1456.
Syphethodes Pascoe, 1875.
marginatus Pascoe. C. 352.
crenatus Broun. C. 352.
decoratus Broun. C. 353.
bullatus Sharp. C. 1150.
sylvius Broun. C. 1150.
cordipennis Broun. C. 1457.
dorsalis Broun. C. 1458.
punctatus Broun. C. 1458.
nodosalis Broun. A. 7, xiv, 56.
simplex Broun. A. 7, xii, 69.
variegatus Broun. B. 1, v, 392.
thoracicus Broun. B. 1, vi, 540.
pensus Broun. B. 1, vi, 540.
truncatus Broun. T. xliv, 435.
Paraphylax Broun, 1880.
squamiger Broun. C. 355.
varius Broun. C. 355.
binodosus Broun. A. 6, xv, 240.
sternalis Broun. A. 7, xiv, 57.
exiguus Broun. B. 1, iii, 195.
Exohadrus Broun, 1893.
volutithorax Broun. C. 356 and 1153.

Subfamily TRACHYSCELINAE.

- Chaeodes* White, 1846.
trachyscelides White. C. 357.
laetus Broun. C. 358.
concolor Sharp. C. 358.
fuscatus Broun. A. 6, xv, 241.
Actizeta Pascoe, 1875.
ammobiooides Pascoe. C. 359.
albata Pascoe. C. 360.

Subfamily DIAPERINAE.

- Menimus* Sharp, 1876.
batesi Sharp. C. 361.
puncticeps Broun. C. 361.
oblongus Broun. C. 362.
caecus Sharp. C. 362.
crinalis Broun. C. 363.
crassus Sharp. C. 363.
dubius Broun. C. 363.
obscurus Broun. C. 364.
thoracicus Broun. C. 364.
curtulus Broun. C. 785.
piceus Broun. C. 786.
striatulus Broun. C. 842.
sinuatus Broun. C. 842.
vicinus Broun. A. 6, xii, 288.

Menimus—continued.

- laevicollis* Broun. A. 6, xv, 242.
humeralis Broun. B. 1, i, 41.
aemulator Broun. B. 1, i, 42.
pubiceps Broun. B. 1, vi, 541.
lineatus Broun. T. xliv, 436.

- Bolitophagæ* Illiger, 1798.
angulifer Blanchard. *Voy. Pole Sud*, iv, 167.

- Uloma* Castelnau, 1840.
tenebrionoides White. C. 366.
laevicostata Blanchard. *Voy. Pole Sud*, iv, 165.

Tenebrio Linnaeus.

- **obscurus* Fabricius.
**militaris* Linnaeus.

- Uломотипус* Broun, 1886.
laevigatus Broun. C. 841.

- Aphthora* Bates, 1872.
ruficeps Bates. C. 367.
glabritarsis Sharp. C. 1153.

Cilibe Brême, 1842.

- opacula* Bates. C. 368.
nitidula Bates. C. 369.
otagensis Bates. C. 370.
elongata Brême. C. 371.
pascoei Bates. C. 372.
humeralis Bates. C. 373.
thoracica Bates. C. 373.
brevipennis Bates. C. 374.
granulosa Brême. C. 375.
rugosa Bates. C. 375.
tibialis Bates. C. 376.
impressifrons Bates. C. 376.
buchanani Broun. C. 377.
huttoni Sharp. C. 378.
marginalis Broun. C. 1155.

- phosphugoides* White. *Voy. Ereb. and Terr., Insects*, 11.

- costella* Broun. A. 7, xv, 544.
lateralis Broun. A. 8, iii, 408.
smithiana Broun. A. 8, iii, 409.
saragooides Pascoe. T. xli, 148.
subcostata Sharp. T. xli, 149.

Zolodinus Blanchard, 1853.

- zealandicus* Blanchard. C. 379.

Lorelus Sharp, 1876.

- priscus* Sharp. C. 381.
pubescens Broun. C. 381.
crassicornis Broun. C. 382.
quadricollis Broun. C. 786.
latulus Broun. B. 1, i, 43.
tarsalis Broun. B. 1, i, 43.
marginalis Broun. B. 1, i, 43.
sternalis Broun. B. 1, i, 44.
nigrescens Broun. B. 1, i, 45.

Demtrius Broun, 1895.

- carinulatus* Broun. A. 6, xv, 243.

Arthropus Sharp, 1876.

- browni* Sharp. C. 383.

Edalus Broun, 1893.

- alienus* Broun. C. 391.

- opuscus* Broun. C. 1160.

- pleuralis* Broun. A. 6, xii, 289.

Group HETEROMERA—continued.

Subfamily CNODALONINAE.

- Artystona* Bates, 1873.
erichsoni Bates. C. 384.
wakefieldi Bates. C. 385.
rugiceps Bates. C. 385.
obscura Sharp. C. 1154.
collaris Sharp. C. 1154.
obsoleta Sharp. C. 1154.
interrupta Redtenbacher. *Reise Novara, Coleop.*, 128.
philpoti Broun. B. 1, i, 45.
tinctella Broun. B. 1, i, 46.
vicina Broun. B. 1, i, 46.

Subfamily HELOPINAE.

- Pheloneis* Pascoe, 1866.
harpalooides White. *Voy. Ereb. and Terror, Insects*, 11.
bullatum Pascoe. C. 386.
intricatum Bates. C. 387.
aucklandicum Broun. C. 387.
amaroides White. C. 388.
aeratum Broun. C. 388.
lentum Broun. C. 389.
zealandicum Bates. C. 390.
thoracicum Bates. C. 390.
cheesemani Broun. C. 787.
chalmieri Broun. C. 787.
indagator Broun. C. 839.
rufilabrum Broun. C. 840.
hansenii Broun. C. 929.
nigritulum Broun. C. 930.
multistriatum Sharp. C. 1155.
simplex Sharp. C. 1156.
sericatum Sharp. C. 1156.
intermedium Sharp. C. 1156.
dunedensis Sharp. C. 1157.
urquharti Broun. C. 1157.
miniatum Broun. C. 1157.
gratiosum Broun. C. 1158.
turgidulum Broun. C. 1158.
tuberculatum Guerin. *Voy. Pole Sud*, iv, 175.
simulans Redtenbacher. *Reise Novara*, 32.
femorale Broun. B. 1, i, 48.
titahense Broun. B. 1, i, 49.
tinctum Broun. B. 1, ii, 111.
calcaratum Broun. B. 1, ii, 110.
hudsoni Broun. A. 8, iii, 411.
appositus Broun. B. 1, iv, 322.
halli Broun. B. 1, v, 392.
angulatus Broun. B. 1, v, 393.
dubitans Broun. B. 1, v, 394.
curtulus Broun. B. 1, v, 394.
complicatum Broun. T. xliv, 437.

Pseudhelops Guérin.

- tuberculatus* Guérin. S. 106.
quadricollis Broun. S. 107.
posticalis Broun. S. 107.
interruptus Broun. S. 108.
substriatus Broun. B. 1, i, 47.
nodosus Broun. B. 1, i, 47.

Cerodolus Sharp, 1886.

- chrysomelooides* Sharp. C. 1161.
genialis Broun. C. 1162.
aeneus Broun. C. 1162.
tuberculatus Broun. B. 1, v, 395.
capitalis Broun. B. 1, v, 395.
sulcisternus Broun. B. 1, v, 396.
curvellus Broun. T. xliv, 437.

Leioperpus Broun, 1893.

- expolitus* Broun. C. 392 and 1160.

Subfamily AMARYGMINAE.

Amarygmus Dalman, 1823.

- zealandicus* Bates. C. 394.

Gnathocerus.

- **cornutus* Fabricius.

Tribolium.

- **ferrugineum* Fabricius.

Family CISTELIDAE.

Amarosoma Redtenbacher, 1868.

- metallicus* White. C. 395.

- sophorae* Broun. C. 396.

- violacea* Broun. B. 1, i, 49.

Xylochus Broun, 1880.

- substriatus* Broun. C. 397.

- tibialis* Broun. C. 397.

- dentipes* Broun. C. 788.

- spinifer* Broun. C. 1168.

Omedes Broun, 1893.

- nitidus* Broun. C. 1169.

- fuscatus* Broun. C. 1170.

- apterus* Broun. A. 6, xv, 244.

Family LAGRIIDAE.

Lagrioda Fairmaire et Germain, 1860
brownii Pascoe. C. 408.

Family MELANDRYIDAE.

Ctenoplectron Redtenbacher, 1868.

- fasciatum* Redtenbacher. C. 401.

- fuliginosum* Broun. C. 402.

- maculatum* Broun. C. 691.

- costatum* Broun. C. 691.

- vittatum* Broun. C. 844.

- coloratum* Broun. C. 959.

- dignum* Broun. C. 960.

Allopterus Broun, 1886.

- reticulatus* Broun. C. 790.

- ornatus* Broun. C. 401.

- instabalis* Broun. C. 844.

- cavelli* Broun. C. 1356.

- simulans* Broun. B. 1, i, 52.

Hylobia Broun, 1880.

- velox* Broun. C. 404.

- undulata* Broun. C. 404.

- nebeculosa* Broun. C. 404.

- nulla* Broun. C. 405.

- nigricornis* Broun. C. 405.

- calida* Broun. C. 405.

- bifasciata* Broun. C. 406.

Group HETEROMERA—continued.

Hylobia—continued.

- minor* Broun. C. 406.
cylindrata Broun. C. 406.
usitata Broun. C. 407.
nigella Broun. C. 407.
sexnotata Broun. B. I, iv, 324.
acuminata Broun. B. I, iv, 325.
arboricola Broun. B. I, iv, 325.
plagiata Broun. T. xliv, 439.
Philpottia Broun, 1915.
maculatus Broun. B. I, iv, 323.
mollis Broun. C. 960.
Chalcodrya Redtenbacher, 1868.
variegata Redtenbacher. C. 403.
Onysius Broun, 1886.
anomalus Broun. C. 843.
pulcher Broun. C. 1357.
Doxozilora Broun, 1909.
punctata Broun. A. 8, iii, 412.
Azylita Broun, 1914.
sericophora Broun. B. I, ii, 112.
Allorchesia Broun, 1914.
valdipes Broun. B. I, ii, 113.
guinnessi Broun. T. xliv, 439.
Neorchesia Broun, 1914.
divergens Broun. B. I, ii, 114.
terricola Broun. B. I, iii, 197.
Mecorchesia Broun, 1914.
spectabilis Broun. B. I, ii, 116.
brevicornis Broun. B. I, ii, 116.
Lyperocharis Broun, 1914.
agilis Broun. B. I, ii, 118.
Nothotelus Broun, 1914.
ocularius Broun. B. I, ii, 119.

Family PYTHIDAE.

- Salpingus* Gyllenhal, 1810.
perpunctatus Broun. C. 398.
bilunatus Pascoe. C. 398.
angusticollis Broun. C. 399.
unguiculus Broun. C. 399.
lautus Broun. C. 400.
hirtus Broun. C. 789.
simplex Broun. C. 789.
quisquilius Broun. C. 789.
fossulatus Broun. C. 1459.
ornatus Broun. A. 6, xv, 245.
cognatus Broun. B. I, i, 49.
rugulosus Broun. B. I, i, 50.
tarsalis Broun. B. I, i, 51.
simplex Broun. B. I, i, 51.
semilaevis Broun. B. I, iii, 196.
atrellus Broun. B. I, iii, 196.
denticollis Broun. B. I, iii, 196.
aterrimus Broun. B. I, vi, 542.
nigricans Broun. B. I, vii, 618.
Agapytho Broun, 1919.
foveicollis Broun. B. I, vi, 543.

Family ANTHICIDAE.

Subfamily ANTHICINAE.

Cotes Sharp, 1877.*vestita* Sharp. C. 411.

13—Trans.

Cotes Sharp—continued.

- proba* Broun. C. 691.
crispi Broun. C. 412.
optima Broun. C. 1165.
punctata Broun. C. 1165.
dorsalis Broun. C. 1166.
distincta Broun. C. 1166.
proxima Broun. C. 1167.
rufa Broun. C. 1167.
insignis Broun. T. xliv, 438.
halliana Broun. B. I, viii, 690.
bullata Broun. B. I, viii, 690.
Anthicus Paykull, 1798.
obscuricornis Broun. C. 411.
pellucidipes Broun. C. 412.
minor Broun. C. 930.
anthracinus Broun. C. 1168.
fallax Broun. C. 1168.
flavitarvis Broun. B. I, ii, 119.
**floralis* Linnaeus.

Subfamily PEDILINAE.

- Macratria* Newman, 1838.
exilis Pascoe. C. 409.
verticalis Sharp. C. 409.
Scaptogetus Broun, 1893.
anthracinus Broun. C. 1358.

Family ODEMERIDAE.

- Selenopalpus* White, 1846.
cyaneus Fabricius. C. 420.
subviridis White. *Voy. Ereb. and Terr., Insects*, 13.
aciphyllae Broun. C. 845.
rectipes Broun. A. 8, iii, 413.
Sessinia Pascoe, 1863.
strigipennis White. C. 420.
latiuscula Broun. C. 421.
lineata Fabricius. C. 421.
pauperata Pascoe. C. 422.
nemoralis Broun. C. 845.
longicornis Broun. C. 1459.
thoracica Broun. A. 6, xi, 290.
reversa Broun. A. 6, xi, 291.
stictica Broun. B. I, iii, 199.
brookesi Broun. B. I, v, 397.
Thelyphassa Pascoe, 1876.
diaphana Pascoe. C. 422.
obscura Broun. C. 422.
conspicua Broun. C. 790.
limbata Broun. C. 961.
fuscate Broun. A. 7, xii, 70.
longicollis Broun. A. 7, xii, 70.
Dammarobius Broun, 1886.
mollis Broun. C. 846.
Baculipalpus Broun, 1880.
rarus Broun. C. 423.
maritimus Broun. A. 8, iii, 414.
Exocalopus Broun, 1893.
pectinatus Broun. C. 1170.
antennalis Broun. A. 7, xii, 71.
nitidiceps Broun. B. I, i, 53.
Phytilea Broun, 1893.
propera Broun. C. 1172.

Group HETEROMERA—continued.

- Techmessa* Bates, 1874.
concolor Bates. C. 424.
telephoroides Bates. C. 424.
attenuata Broun. C. 1172.
varians Broun. C. 1173.
rugicollis Broun. B. 1, i, 52.
longicollis Broun. A. 7, xii, 70.
Techmessodes Broun, 1893.
picticornis Broun. C. 424.
versicolor Broun. C. 1173.
cephalotes Broun. B. 1, i, 53.
Metasclera Broun, 1914.
nigricans Broun. B. 1, iii, 198.
arborea Broun. B. 1, iii, 198.
Nacerdes.
**melanura* Schmidt.

Family MORDELLIDAE.

Subfamily MORDELLINAE.

- Mordella* Linnaeus, 1758.
antarctica White. *Voy. Ereb. and Terror, Insects*, 12.
funerea Pascoe. C. 414.
tibialis Broun. C. 414.

- Mordella*—continued.
detracta Pascoe. C. 414.
tairuensis Broun. C. 414.
Mordellistena Costa, 1855.
jucunda Broun. C. 415.
neglecta Broun. C. 415.
Zeamordella Broun, 1880.
monacha Broun. C. 847.
Apesina Broun, 1881.
stewarti Broun. C. 693.
tener Broun. C. 693.
Cryptomera Broun, 1893.
nigra Broun. C. 1359.
Subfamily RHIPIPHORINAE.
Rhipistena Sharp, 1878.
lugubris Sharp. C. 417.
cryptartha Broun. A. 7, xiv, 58.
sulciceps Broun. A. 7, xiv, 59.
Allocinops Broun, 1919.
brookesi Broun. B. 1, vii, 619.
Sharpia Broun, 1880 (not of Tournier, 1873).
hirtella Broun. C. 418.
Xylophilus.
xenarthrus Broun. B. 1, i, 54.

Group RHYNCHOPHORA.

Family BRENTHIDAE.

- Teramocerus*.
barbicornis Fabricius. C. 544.
cylindricornis Fabricius. C. 543.

Family ANTHRIBIDAE.

- Anthribus* Geoffroy, 1762.
spinifer Sharp. C. 545.
sharpi Broun. C. 546.
brouni Sharp. C. 547.
halli Broun. B. 1, vi, 586.
sandageri Broun. C. 1261.
obsoletus Broun. C. 1262.
rudis Sharp. C. 547.
bulletus Sharp. C. 548.
vates Sharp. C. 548.
inornatus Sharp. C. 1255.
concolor Sharp. C. 1256.
brunneus Broun. C. 1262.
flavipilus Broun. A. 6, xv, 417.
tessellatus Broun. C. 1263.
curvatus Broun. C. 1263.
lanuginosus Broun. C. 549.
phyamatodes Redtenbacher. C. 550.
fuscopictus Broun. C. 564.
hetaera Sharp. C. 550.
discedens Sharp. C. 551.
tuberous Sharp. C. 1254.
altus Sharp. C. 551.
ornatus Sharp. C. 552.
huttoni Sharp. C. 553.
decens Broun. C. 1264.

Anthribus—continued.

- finimus* Broun. C. 1264.
anxius Broun. C. 1265.
laetabilis Broun. C. 1265.
deterius Broun. C. 1266.
impar Broun. C. 1266.
picipictus Broun. C. 742.
nigrescens Broun. C. 743.
torulosus Broun. C. 743.
crassus Sharp. C. 560.
nanus Sharp. C. 561.
atomus Sharp. C. 561.
purpureus Broun. C. 559.
minor Broun. C. 1260.
fungicola Broun. C. 1261.
thoracicus Broun. C. 1261.
cucullatus Sharp. C. 1255.
anguliceps Broun. B. 1, i, 72.
rugifer Broun. B. 1, i, 72.
maurus Broun. B. 1, i, 72.
suspectus Broun. B. 1, i, 73.
albiceps Broun. B. 1, ii, 138.
imitarius Broun. B. 1, ii, 139.
lewisi Broun. A. 8, iv, 159.
philpotti Broun. A. 8, iv, 160.
venustus Broun. B. 1, iii, 254.
cornutellus Broun. T. xlvi, 150.
levinensis Broun. T. xlvi, 150.
obscurus Broun. T. xlvi, 151.
warriensis Broun. T. xlvi, 152.
cristatellus Broun. T. xliii, 111.
propinquus Broun. T. xliii, 112.
pilicornis Broun. T. xliii, 112.

Group RHYNCHOPHORA—continued.

- Exilis* Pascoe, 1873 (= *Lawsonia* Sharp).
lawsoni Sharp. C. 556.
variabilis Sharp. C. 557.
Arecopais Broun, 1893.
spectabilis Broun. C. 558.
Proscoporhinus Montrouzier, 1860.
meinertzhageni Broun. C. 563.
signatus Broun. C. 1268.
viridescens Broun. C. 1269.
albifrons Sharp. C. 1269.
Etnalis Sharp, 1873.
spinicollis Sharp. C. 554.
Eugonissus Broun. C. 1893.
prictipes Broun. C. 1257.
convexus Broun. C. 555.
proximus Broun. C. 555.
obtusus Sharp. C. 1256.
turneri Broun. T. xlv, 148.
sylvanus Broun. T. xlv, 149.
Dysnocryptus Broun, 1893.
plagiatus Broun. C. 1258.
testaceus Broun. C. 1259.
pallidus Broun. C. 1259.
maculifer Broun. C. 1260.
nigricans Broun. C. 1260.
inflatus Sharp. C. 562.
dignus Broun. C. 563.
rugosus Sharp. C. 563.
setigerus Broun. B. 1, vii, 658.
Araeocerus Pascoe, 1876.
pardalis Pascoe. C. 559.
Xenanthribus Broun, 1893.
hirsutus Broun. C. 1270.
Doticus.
**pestilens* Olliff.

Family CURCULIONIDAE.

Subfamily OTIORHYNCHINAE.

- Nicæana* Pascoe, 1877.
modesta Pascoe. C. 428.
concinna Broun. C. 847.
cinerea Broun. C. 931.
cervina Broun. C. 1360.
tarsalis Broun. C. 1460.
infuscata Broun. A. 8, iv, 52.
catoptoides Broun. B. 1, iii, 200.
placida Broun. B. 1, iii, 201.
gracilicornis Broun. B. 1, iii, 201.
crassifrons Broun. B. 1, v, 398.
nesophila Broun. T. xlvi, 97.
cordipennis Broun. B. 1, vii, 619.
Aphela.
pictipes Broun. A. 7, xii, 75.
Stygeopetes Broun, 1893.
tittoralis Broun. C. 1176.
Thomitus Broun, 1910.
halli Broun. T. xlvi, 103.
Cecyropa Pascoe, 1875.
tychioides Pascoe. C. 437.
maritime Broun. C. 438.
brevipennis Broun. C. 438.
alba Broun. C. 698.
varia Broun. C. 698.
discors Broun. C. 699, and A. 7, xiv, 106.

- Cecyropa*—continued.
setigera Broun. C. 905.
macularia Broun. C. 961.
albicans Sharp. C. 1175.
fumosa Broun. C. 1175.
alternata Broun. A. 7, xiv, 105.
lineifera Broun. A. 7, xii, 72.
striata Broun. A. 7, xii, 73.
sulcifrons Broun. B. 1, v, 398.
striatella Broun. B. 1, v, 399.
jucunda Broun. B. 1, v, 400.
laticollis Broun. B. 1, v, 400.
litorea Broun. B. 1, vi, 343.
Nonnotus Sharp, 1886.
griseolus Sharp. C. 1177.
electus Broun. C. 1177.
pallescens Broun. C. 1178.
albicans Broun. C. 443.
albatus Broun. C. 694.
nigricans Broun. T. xlvi, 98.
Tigones Broun, 1886.
caudata Broun. C. 855.
oscularis Broun. C. 856.
certa Broun. C. 856.
grisea Broun. C. 856.
cervina Broun. C. 857.
arctica Broun. C. 1180.
variegata Broun. C. 1181.
scutellaris Broun. C. 1181.
cruda Broun. C. 1182.
antennalis Broun. C. 1182.
diversa Broun. C. 1183.
rufula Broun. C. 1183.
cavelli Broun. C. 1361.
obscura Broun. C. 1361.
cuspidata Broun. C. 694.
plana Broun. C. 700.
dispar Broun. A. 7, xiv, 107.
philpotti Broun. B. 1, i, 57.
humeralis Broun. B. 1, iii, 202.
flectiscapus Broun. B. 1, iii, 202.
nasalis Broun. B. 1, v, 401.
citimus Broun. B. 1, v, 401.
thoracica Broun. B. 1, v, 402.
variata Broun. B. 1, v, 403.
murina Broun. B. 1, v, 403.
assimilis Broun. B. 1, v, 404.
setosa Broun. B. 1, v, 404.
longiceps Broun. B. 1, v, 405.
bicostellus Broun. B. 1, vi, 544.
allopicta Broun. T. xlvi, 100.
gracilis Sharp. C. 1179.
binodulus Sharp. C. 1179.
robustus Sharp. C. 1180.
rugosa Broun. T. xlvi, 99.
longipes Broun. B. 1, vii, 620.
Epitimates Pascoe, 1877.
lutescens Pascoe. C. 435.
wakefieldi Sharp. C. 1183.
cupreus Broun. B. 1, v, 406.
foveiger Broun. B. 1, v, 406.
grisealis Broun. T. xlvi, 97.
bicolor Broun. B. 1, vii, 621.
densus Broun. B. 1, vii, 622.

Group RHYNCHOPHORA—continued.

- Exonastus* Broun, 1919.
amplus Broun. B. 1, vii, 623.
fumidus Broun. B. 1, vii, 623.
- Platyomida* White, 1846.
bimodes White. C. 442.
censoria Pascoe. C. 442.
aculeata Broun. C. 442.
perniciosa Broun. C. 442.
apicalis Broun. C. 701.
amota Broun. C. 793.
enysi Broun. C. 852.
aequa Broun. C. 932.
hochstetteri Redtenbacher. C. 476.
simulatrix Sharp. C. 1185.
depressa Broun. C. 1185.
brevicornis Broun. A. 7, xiv, 107.
caudata Broun. B. 1, i, 54.
humeralis Broun. B. 1, i, 55.
tibialis Broun. B. 1, i, 56.
verrucosa Broun. B. 1, ii, 120.
philpotii Broun. B. 1, vi, 545.
hamiltoni Broun. T. xlv, 100.
morosa Broun. T. xlv, 101.
dorsalis Broun. B. 1, v, 407.
cuprealis Broun. B. 1, v, 408.
fuscella Broun. B. 1, v, 409.
rectirostris Broun. B. 1, v, 409.
hystricula Broun. B. 1, v, 410.
sulcicollis Broun. B. 1, v, 410.
latipennis Broun. B. 1, v, 411.
versicolor Broun. T. xlvi, 103.
Thesius Broun, 1909.
inophlaeoides. A. 8, iv, 60.
- Rhynchogonus*.
germanus Broun. A. 6, xii, 293.
- Lyperobates* Broun, 1893.
asper Broun. C. 1462.
guinnessi Broun. T. xlvi, 101.
elegantulus Broun. T. xlvi, 102.
rostralis Broun. T. xlvi, 103.
punctatus Broun. T. xlvi, 103.
waterworthi Broun. B. 1, i, 57.
carinifer Broun. B. 1, i, 58.
ardens Broun. B. 1, i, 59.
virilis Broun. A. 8, iv, 58.
- Zenographus* Broun, 1915.
metallescens Broun. B. 1, iv, 327.
allinotatus Broun. B. 1, vii, 625.
- Hygrochus* Broun, 1881.
oscitans Broun. C. 703.
verrucosus Broun. C. 1220.
illepidus Broun. C. 1220.
oculatus Broun. A. 6, xii, 292.
cordipennis Broun. B. 1, i, 59.
scutellaris Broun. B. 1, ii, 121.
granifer Broun. A. 8, iv, 58.
monilifer Broun. B. 1, vii, 625.
- Homodus* Broun, 1881.
fumeus Broun. C. 703.
posticalis Broun. B. 1, viii, 697.
longicornis Broun. B. 1, viii, 698.
cuprealis Broun. B. 1, viii, 698.
- Aporolobus* Sharp, 1886.
irritus Pascoe. C. 431 and C. 1186.
hariolus Broun. C. 848.
- Aporolobus*—continued.
cecyropioides Broun. C. 905.
rugosus Broun. C. 930.
pallidus Broun. C. 1365.
albosparsus Broun. B. 1, ii, 120.
scapalis Broun. B. 1, vi, 545.
Proboscocelus Broun, 1909.
sculpturatus Broun. A. 8, iv, 56.
Notiopatae Broun, 1893.
setifer Broun. C. 1187.
sternalis Broun. C. 1462.
clarus Broun. C. 431.
terricola Broun. T. xlvi, 105.
Phaeocharis Broun, 1913.
cuprealis Broun. T. xlvi, 105.
punctatus Broun. T. xlvi, 105.
Paelocharis Broun, 1893.
inflata Broun. C. 1188.
vestita Broun. C. 1463.
corpulenta Broun. C. 431.
Getopsephus Broun, 1913.
acuminatus Broun. T. xlvi, 107.
costifer Broun. B. 1, vi, 558.
funestus Broun. B. 1, vi, 559.
Protolobus Sharp, 1886.
obscurus Sharp. C. 1188.
porculus Pascoe. C. 430.
granicollis Broun. B. 1, iii, 203.
nodosus Broun. B. 1, v, 412.
Drymaria Broun, 1909.
ciliipes Broun. A. 8, iv, 57.
Heterexis Broun, 1909.
scutipennis Broun. S. 114.
laevinsculus Broun. S. 114.
Catodryobius Broun, 1909.
vestitus Broun. S. 109.
benhami Broun. S. 110.
tetricus Broun. S. 110.
erubescens Broun. S. 111.
grandis Broun. S. 112.
Inocatoptes Broun, 1909.
incertus Broun. S. 113.
Catoptes Schonherr, 1842.
obliquis Schonherr. C. 428.
parilis Pascoe. C. 443.
compressus Broun. C. 429.
obliquisignatus Bohem. C. 693.
attenuatus Broun. C. 791.
pilosellus Broun. C. 853.
bicostatus Broun. C. 853.
vexator Broun. A. 7, xiv, 108.
egeus Broun. A. 7, xiv, 109.
duplex Broun. A. 7, xiv, 110.
spermophilus Broun. A.
aequalis Broun. A.
constrictus Broun. B. 1, i, 60.
tibialis Broun. C. 854.
stolidus Broun. C. 854.
caliginosus Broun. C. 1189.
posticalis Broun. C. 1189.
humeralis Broun. C. 1190.
chalmeri Broun. C. 1190.
asperellus Broun. C. 1191.
cheesemani Broun. C. 1191.
decorus Broun. C. 1192.

Group RHYNCHOPHORA—continued.

Catoptes—continued.

- brevicornis* Sharp. C. 1193.
aemulator Broun. C. 1193.
tenebricus Broun. C. 1194.
latipennis Broun. C. 1362.
furvus Broun. C. 1362.
vastator Broun. C. 1463.
scutellaris Sharp. C. 1503.
longulus Sharp. C. 1504.
limbatus Broun. A. 8, iv, 60.
spectabilis Broun. B. 1, iii, 205.
subnitidus Broun. B. 1, iii, 206.
curvatus Broun. B. 1, iii, 206.
carinalis Broun. B. 1, iii, 207.
argentalis Broun. B. 1, iii, 208.
fumosus Broun. B. 1, iii, 208.
subplicatus Broun. B. 1, v, 412.
dehiscens Broun. B. 1, v, 413.
robustus Broun. B. 1, v, 414.
albosparsus Broun. B. 1, v, 415.
pallidipes Broun. B. 1, v, 415.
flaviventris Broun. B. 1, v, 416.
nigrans Broun. B. 1, v, 416.
lobatus Broun. B. 1, vii, 626.
apicalis Broun. B. 1, viii, 691.
Haplolobus Broun, 1893.
saevus Broun. C. 1199.
aethiops Broun. C. 1199.
gregalis Broun. C. 1199.
frontalis Broun. B. 1, iii, 204.
granulatus Broun. B. 1, iii, 204.
Neoevas Broun, 1919.
celmisiae Broun. B. 1, vii, 624.
Heterodiscus Sharp, 1886.
insolitus Sharp. C. 1197.
Echinopeptus Broun, 1886.
dilatatus Broun. C. 852.
verrucatus Broun. B. 1, iii, 211.
dorsalis Broun. B. 1, in, 212.
Brachyolus White, 1846.
punctatus White. C. 432.
brevisculus Broun. C. 440.
elegans Broun. C. 1194.
inaequalis Sharp. C. 1195.
huttoni Sharp. C. 1195.
bagoooides Sharp. C. 1195.
punctipennis Sharp. C. 1196.
longicollis Sharp. C. 1196.
viridescens Broun. C. 1363.
posticalis Broun. C. 1364.
albescens Broun. A. 7, xii, 73.
cervalis Broun. A. 7, xii, 74.
sylvaticus Broun. B. 1, i, 61.
asperatus Broun. B. 1, iii, 209.
fuscipictus Broun. B. 1, iii, 210.
bicosatus Broun. B. 1, v, 416.
terricola Broun. B. 1, v, 417.
labeculatus Broun. T. xlv, 107.
varius Broun. T. xlv, 108.
nodirostris Broun. B. 1, vii, 627.
obscurus Broun. B. 1, vii, 628.
Bryodrassus Broun, 1917.
miricollis Broun. B. 1, v, 418.
Inocatoptes Broun, 1902.
incertus Broun. T. xxxiii, 178.

- Agatholobus* Broun, 1913.
waterhousei Broun. T, xlv, 109.
Inophloeus Pascoe, 1875.
traversii Pascoe. C. 439.
inurus Pascoe. C. 439.
rhesus Pascoe. C. 439.
vittiosus Pascoe. C. 440.
rubidus Broun. C. 699.
nigellus Broun. C. 700.
praelatus Broun. C. 851.
sulcifer Broun. C. 905.
costifer Broun. C. 932.
egregius Broun. C. 962.
nodifer Broun. C. 1200.
alacer Broun. C. 1200.
albonotatus Broun. C. 1201.
vestitus Broun. C. 1202.
suturalis Broun. C. 1464.
villaris Pascoe. A. 4, xvi, 220.
medius Broun. A. 6, xii, 294.
sternalis Broun. A. 7, xiv, 111.
discrepans Broun. A. 7, xiv, 112.
longicornis Broun. A. 7, xiv, 113.
quadricollis Broun. A. 8, iv, 53.
laeticus Broun. A. 8, iv, 54.
peusus Broun. B. 1, iii, 212.
sulcicollis Broun. B. 1, iii, 213.
aplorhinus Broun. B. 1, iv, 327.
tricostatus Broun. B. 1, iv, 328.
fuscatus Broun. B. 1, v, 419.
collinus Broun. B. 1, v, 419.
cuprellus Broun. B. 1, vii, 628.
obsoletus Broun. B. 1, vii, 629.
sexnotatus Broun. B. 1, vii, 630.
festucae Broun. B. 1, vii, 631.
Oiorhynchus Germar.
**sulcatus* Fabricius.

Subfamily CYLINDRORHINAE.

- Phaeophanus* Broun, 1886.
rugosus Broun. C. 793.
similis Broun. C. 1216.
lituratus Broun. C. 1216.
o'connori Broun. B. 1, vii, 631.
ornatus Broun. B. 1, vii, 632.
Heteraomus Broun, 1893.
longipes Broun. C. 1218.
Anogotus Sharp, 1882.
helmsi Sharp. C. 1215.
palleucus Broun. A. 7, xii, 76.
Mecosargon Broun, 1915.
costipennis Broun. B. 1, iv, 329.
Sargon Broun, 1903.
carinatus Broun. A. 7, xii, 77.
hudsoni Broun. A. 8, iv, 70.
Pparcus Broun, 1904.
lewisi Broun. A. 7, xiv, 114.
halli Broun. B. 1, iv, 330.
Phaedropholus Broun, 1910.
o'connori Broun. B. 1, i, 67.
Phoroteles Broun, 1893.
graniger Broun. C. 850.
Heterotyles Broun, 1886.
argentatus Broun. C. 795.

Group RHYNCHOPHORA—continued.

- Rhadinosomus* Schönherr, 1840.
acuminatus Schönherr. C. 430.
Heteromias Broun, 1913.
foveirostris Broun. T. xlv, 121.
Hadrampus Broun, 1910.
spinipennis Broun. T. xlvi, 105.

- Subfamily MOLYTINAE.
- Lyperobius* Pascoe, 1876.
huttoni Pascoe. C. 445.
tuberculatus Pascoe. C. 445.
carinatus Broun. C. 702.
cupiendus Broun. C. 962.
laeviusculus Broun. T. xxiii, 179.
hudsoni Broun. B. 1, ii, 127.
aciphyllae Broun. B. 1, v, 442.
fallax Broun. B. 1, v, 443.
spedenii Broun. B. 1, v, 443.
Liparogetus Broun, 1915.
sulcatus Broun. B. 1, iv, 331.
Lyperopais Broun, 1893.
mirus Broun. C. 1222.
alternans Broun. B. 1, ii, 128.

- Subfamily HYLOBITINAE.
- Eiratus* Pascoe, 1877.
parvulus Pascoe. C. 447.
tetricus Broun. C. 447.
costatus Broun. C. 794.
ornatus Broun. C. 906.
suavis Broun. C. 934.
pyriformis Broun. C. 1223.
rugosus Broun. C. 1223.
nitirostris Broun. B. 1, i, 65.
Bryocatus Broun, 1914.
alternans Broun. B. 1, iii, 219.
jugosus Broun. B. 1, iii, 219.
nodicollis Broun. B. 1, iii, 220.
nigrirostris Broun. B. 1, iii, 220.
ampalus Broun. B. 1, iii, 221.
burrowsi Broun. B. 1, iv, 332.
quadrucollis Broun. B. 1, vi, 559.
rubidus Broun. B. 1, vi, 559.
elegans Broun. B. 1, vi, 560.
humeratus Broun. B. 1, vi, 560.
lugubris Broun. B. 1, vi, 561.
crassirostris Broun. B. 1, vi, 561.
niticollis Broun. B. 1, vii, 633.
iridescent Broun. B. 1, vii, 633.
rugosus Broun. B. 1, vii, 634.
plicatus Broun. B. 1, viii, 693.
crassipes Broun. B. 1, viii, 694.
angustus Broun. B. 1, viii, 694.
fordi Broun. B. 1, viii, 694.
diversus Broun. B. 1, viii, 695.
thoracicus Broun. B. 1, viii, 695.
ovipennis Broun. B. 1, viii, 696.
Stiboderma Broun, 1909.
impressipennis Broun. A. 8, iv, 68.
Exeiratus Broun, 1914.
setarius Broun. B. 1, ii, 129.
Athor Broun, 1909.
arcifera Broun. A. 8, iv, 69.

- Dryopais* Broun, 1886.
variabilis Broun. C. 933.
Paedaretus Pascoe, 1876.
hispidus Pascoe. C. 444.
rufulus Broun. B. 1, i, 65.

Subfamily RHYPAROSOMINAE.

- Phrynxus* Pascoe, 1875.
terreus Pascoe. C. 433.
astutus Pascoe. C. 433.
celatus Broun. C. 433.
modicus Broun. C. 434.
facetus Broun. C. 695.
intricatus Broun. C. 848.
tuberculatus Broun. C. 849.
differens Broun. C. 849.
rufipes Broun. C. 849.
simplex Broun. C. 1202.
costirostris Broun. C. 1203.
humeralis Broun. C. 1467.
brevipennis Broun. C. 1468.
cepsis Broun. C. 1468.
longulus Broun. B. 1, i, 61.
rufiventris Broun. B. 1, ii, 121.
bicarinellus Broun. A. 8, iv, 61.
ventralis Broun. A. 8, iv, 62.
squamalis Broun. B. 1, vi, 546.
humilis Broun. B. 1, vi, 547.
amoenus Broun. B. 1, vi, 547.
setipes Broun. T. xlv, 110.
bindosus Broun. T. xlv, 111.
asper Broun. T. xlvi, 104.
conspicuus Broun. B. 1, vii, 634.
blandus Broun. B. 1, vii, 635.
Bradyptatae Broun, 1893.
capitatis Broun. C. 1206 and 850.
armiger Broun. A. 6, xii, 296.
dilaticollis Broun. A. 8, iv, 66.
interstitialis Broun. A. 8, iv, 67.
minor Broun. T. xlv, 112.
subnodifer Broun. B. 1, vii, 636.
impressum Broun. B. 1, vii, 637.
Erymneus Pascoe, 1877.
sharpia Pascoe. C. 436.
scabiosus Broun. C. 436.
castaneus Broun. C. 436.
granulatus Broun. C. 437.
longulus Broun. C. 931.
coenosus Broun. C. 932.
ferrugatus Broun. C. 1203.
irregularis Broun. C. 1367.
crassipes Broun. C. 1469.
firmus Broun. C. 1470.
probus Broun. A. 6, xii, 299.
terrestris Broun. B. 1, vi, 548.
Lithocia Broun, 1893.
fimbriata Broun. C. 1471.
angustula Broun. B. 1, iii, 214.
ciliata Broun. B. 1, v, 420.
setirostris Broun. B. 1, v, 421.
basalis Broun. B. 1, v, 421.
rectisetosa Broun. B. 1, v, 422.
nigrirista Broun. B. 1, v, 423.
acuminata Broun. T. xlv, 111.
stictica Broun. B. 1, viii, 692.

Group RHYNCHOPHORA—continued.

- Abrotheus* Broun, 1917.
placitus Broun. B. 1, v, 424.
Pachyprypnus Broun, 1886.
pyriformis Broun. C. 792.
longiusculus Broun. C. 434.
modicus Broun. A. 7, xiv, 117.
Dohroethus Broun, 1893.
dumetosus Broun. C. 1208.
vestitus Broun. C. 1209.
Astyplus Broun, 1893.
conicus Broun. C. 1210.
brevicornis Broun. B. 1, vii, 640.
Styphlotelus Broun, 1893.
foveatus Broun. C. 1368.
fascicularis Broun. C. 1369.
Halliea Broun, 1917.
squamipes Broun. B. 1, v, 425.
antennalis Broun. B. 1, v, 425.
longicollis Broun. B. 1, v, 426.
cuneata Broun. B. 1, vii, 640.
Allotyphlus Broun, 1919.
yugosus Broun. B. 1, vi, 549.
Phrynidodes Broun, 1919.
scruposus Broun. B. 1, vi, 551.
Cuneopterus Sharp, 1886.
conicus Sharp. C. 1204.
tenuicornis Broun. A. 6, xii, 297.
Chamaepsephus Broun, 1893.
aurisetifer Broun. C. 1207.
Amphisktira Broun, 1909.
umbricola Broun. A. 8, iv, 64.
Araeoscapus Broun, 1893.
ovipennis Broun. A. 6, xii, 300.
flavipes Broun. C. 1473.
thoracicus Broun. C. 1474.
obscurus Broun. C. 1475.
fasciculatus Broun. B. 1, i, 62.
mucronatus Broun. B. 1, i, 63.
punctipennis Broun. B. 1, i, 64.
brevicollis Broun. B. 1, ii, 125.
ardens Broun. A. 8, iv, 64.
estriatus Broun. A. 8, iv, 65.
ocularius Broun. B. 1, iii, 215.
subcostatus Broun. B. 1, vii, 637.
Memes Broun, 1903.
rufirostris Broun. A. 7, xii, 79.
Baeosomus Broun, 1904.
tacitus Broun. A. 7, xiv, 119.
Bantiades Broun, 1893.
fuscatus Broun. C. 1371.
valgus Broun. C. 1472.
suratalis Broun. B. 1, ii, 123.
nodosus Broun. B. 1, ii, 125.
cupiendus Broun. B. 1, iii, 215.
morosus Broun. B. 1, v, 426.
cylindricus Broun. B. 1, v, 427.
notatus Broun. B. 1, v, 428.
rectalis Broun. B. 1, vi, 551.
trifoveatus Broun. B. 1, vi, 552.
Abantiades Broun, 1914.
nodipennis Broun. B. 1, iii, 217.
pusillus Broun. B. 1, iii, 217.
gratulus Broun. B. 1, v, 428.
Rachidiscus Broun, 1893.
granicollis Broun. C. 1478.
multinodosus Broun. T. xlvi, J16.
- Rachidiscodes* Broun, 1917.
altipennis Broun. B. 1, v, 430.
glabrus Broun. B. 1, vi, 552.
Dermotrichus Sharp, 1886.
mundulus Broun. C. 1205.
multicostatus Broun. B. 1, v, 434.
elegantis Broun. B. 1, v, 435.
vicinus Broun. B. 1, vii, 638.
curvirostris Broun. B. 1, vii, 639.
Allalops Broun, 1917.
carinatus Broun. B. 1, v, 430.
Hycanus Broun, 1905.
cockaynei Broun. A. 7, xv, 546, and S. 116.
frontalis Broun. S. 116.
Stilbodiscus Broun, 1909.
setarius Broun. S. 118.
Clypeorhynchus Sharp, 1883.
gracilipes Sharp. C. 1211.
thoracicus Broun. C. 1211.
brevicornis Broun. C. 1212.
impressus Broun. C. 1212.
ovipennis Broun. C. 1213.
cordipennis Broun. C. 1370.
cristatus Broun. C. 1370.
inophloeoides Broun. A. 7, xiv, 117.
nodiceps Broun. B. 1, iii, 218.
striatus Broun. B. 1, iv, 333.
clarulus Broun. B. 1, v, 431.
merus Broun. B. 1, v, 432.
setosus Broun. B. 1, v, 433.
nitidellus Broun. B. 1, v, 433.
halli Broun. B. 1, v, 433.
dorsalis Broun. B. 1, vi, 553.
bicarinatus Broun. B. 1, vi, 554.
calvulus Broun. T. xlvi, 113.
caudatus Broun. T. xlvi, 114.
tenueculus Broun. B. 1, vii, 641.
furvus Broun. B. 1, vii, 642.
bifoveatus Broun. B. 1, viii, 692.
Phygoothalpus Broun, 1915.
sulcicollis Broun. T. xlvi, 118.
nitidulus Broun. B. 1, iv, 334.
majusculus Broun. B. 1, iv, 334.
philotti Broun. B. 1, v, 435.
granissimus Broun. B. 1, v, 436.
subcipennis Broun. B. 1, v, 437.
striatus Broun. B. 1, vi, 556.
anthracinus Broun. B. 1, vi, 557.
Daenophylla Broun, 1893.
setosa Broun. C. 1472.
variegata Broun. A. 6, xii, 301.
sparsa Broun. B. 1, vi, 554.
Nestrius Broun, 1893.
serripes Broun. C. 1480, and A. 6, xii, 302.
crassicornis Broun. B. 1, iv, 335.
sulcirostris Broun. B. 1, v, 437.
prolixus Broun. B. 1, v, 438.
zenoscelis Broun. B. 1, vi, 555.
simmondsi Broun. B. 1, vi, 556.
Plotinus Broun, 1893.
ovithorax Broun. C. 1481.
Phemus Broun, 1893.
scabralis Broun. C. 1214.
rufipes Broun. C. 1482.

Group RHYNCHOPHORA—continued.

Phemus—continued.

- curvipes* Broun. T. xlv, 114.
constrictus Broun. T. xlv, 115.

Rystheus Broun, 1893.

- ocularius* Broun. C. 1219.
notabilis Broun. B. 1, v, 441.

Mosogenes Broun, 1917.

- longiventris* Broun. B. 1, v, 439.
acerbus Broun. B. 1, vii, 643.

Sosgenes Broun, 1893.

- carinatus* Broun. C. 1477.
longicollis Broun. B. 1, ii, 122.
discalis Broun. B. 1, v, 440.
planirostris Broun. T. xlv, 116.

Phyllodytes Broun, 1893.

- foveatus* Broun. C. 1479.
irregularis Broun. B. 1, i, 64.

Phronira Broun, 1893.

- sulcirostre* Broun. C. 696.
asper Broun. C. 696.
costosum Broun. C. 696.
osculans Broun. C. 697.
simplex Broun. C. 697.
striatum Broun. C. 697.
nodosum Broun. C. 1476.

Subfamily AMYCTERINAE.

Tocris Broun, 1904.

- latirostris* Broun. A. 7, xiv, 160.
laevicostata Broun. B. 1, ii, 126.
hamiltoni Broun. T. xlv, 119.
pascoi Broun. B. 1, v, 442.
aterrima Broun. T. xlv, 118.

Subfamily ERIRHININAE.

Philacta Broun, 1880.

- testacea* Broun. C. 449.
maculifera Broun. A. 7, xiv, 119.
Erirhinus Schönherr, 1826.
acalyptoides Pascoe. C. 450.
limbatus Pascoe. C. 450.
glottis Pascoe. C. 450.
fusconotatus Broun. C. 451.
discoideus Broun. C. 451.
flavitarsis Broun. C. 451.
fasciatus Broun. C. 452.
rubricalis Broun. C. 452.
viridipennis Broun. C. 452.
sexfasciatus Broun. C. 706.
dolosus Broun. C. 707.
fascialis Broun. C. 707.
crucigerus Broun. C. 708.
anchoralis Broun. C. 708.
gracilirostris Broun. C. 708.
stramineus Broun. C. 709.
subconicollis Broun. B. 1, viii, 699.
nocens Broun. C. 709.
acceptus Broun. C. 710.
titahensis Broun. T. xlv, 122.
poecilus Broun. B. 1, viii, 700.
femoralis Broun. C. 710.
concolor Broun. C. 710.
eustictus Broun. C. 857.
veronicae Broun. C. 858.

Erirhinus—continued.

- thomsoni* Broun. C. 858.
bicolorus Broun. C. 859.
melastomus Broun. C. 859.
confusus Broun. C. 860.
xenorhinus Broun. C. 860.
durus Broun. C. 861.
fuscoventris Broun. C. 861.
fulvus Broun. C. 862.
cheesemani Broun. C. 963.
disformipes Broun. C. 964.
nesobius Broun. C. 965.
anxius Broun. C. 1372.
fuscipes Broun. C. 1373.
pectoralis Broun. B. 1, ii, 131.
insignis Broun. A. 8, iv, 130.
insolitus Broun. A. 8, iv, 131.
spadiceus Broun. A. 8, iv, 132.
castigatus Broun. A. 8, iv, 132
sylvaticus Broun. B. 1, iii, 225.
cordipennis Broun. B. 1, iv, 336.
celmisiae Broun. B. 1, v, 444.
dilucidus Broun. B. 1, vi, 562.
altivagans Broun. B. 1, vi, 562.
leucocomus Broun. B. 1, vi, 563.
oleariae Broun. T. xlv, 122.
exilis Broun. T. xlv, 123.
dracophyliae Broun. S. 118.
obscurus Broun. B. 1, viii, 699.
Anthonomus, Germar.
**pomorum*, Linnaeus.
Dorytomus Stephens, 1831.
trilobus Pascoe. C. 453.
rufirostris Broun. C. 453.
lateralis Broun. C. 711.
sudus Broun. C. 711.
ochraceus Broun. C. 712.
aericomus Broun. C. 795.
aciphyllae Broun. C. 965.
grossus Broun. C. 1224.
fulvescens Broun. B. 1, iii, 225.
melastictus Broun. B. 1, iii, 226.
albisetosus Broun. B. 1, iii, 227.
terrestris Broun. B. 1, iii, 227.
floricola Broun. B. 1, iii, 228.
methvenensis Broun. B. 1, iv, 336.
vittatus Broun. B. 1, vi, 563.
maorinus Broun. T. xlv, 123.
consonus Broun. T. xlv, 124.
australis Broun. B. 1, vii, 644.
Neomycta Pascoe, 1877.
pubicaris Pascoe. C. 457.
rubida Broun. C. 457.
seticeps Broun. B. 1, iii, 229.
Euprocas Broun, 1886.
scutulus Broun. C. 1375.
Alloprocas Broun, 1886.
rufus Broun. C. 1374.
niger Broun. C. 1374.
muticus Broun. B. 1, iii, 229.
Xerostygnum Broun, 1903.
binodulus Broun. A. 7, vii, 80.
pullus Broun. B. 1, i, 68.
Celestotelus Broun, 1886.
fulvus Broun. C. 1375.

Group RHYNCHOPHORA—continued.

- Phorostichus* Broun, 1882.
linearis Broun. C. 705.
Simachus Broun, 1886.
montanus Broun. C. 963.
cuneipennis Broun. B. I, iii, 230.
placens Broun. B. I, vii, 644.
Aganeuma Broun, 1886.
rufula Broun. C. 1484.
Stilobopsis Broun, 1886.
politus Broun. C. 1483.
Etheophanus Broun, 1886.
pinguis Broun. C. 1233.
striatus Broun. B. I, i, 69.
punctiventris. B. I, iii, 231.
optandus Broun. B. I, viii, 701.
nitidellus Broun. B. I, viii, 701.
obscurus Broun. B. I, viii, 702.
Praolepra Pascoe, 1880.
squamosa Pascoe. C. 454.
infusca Broun. C. 452.
albopicta Broun. C. 712.
rufescens Broun. C. 713.
varia Broun. C. 713.
asperirostris Broun. C. 713.
pallida Broun. C. 714.
fultoni Broun. C. 908.
Megacolabus Broun, 1886.
sculpturatus Broun. C. 1378.
Aneuma Pascoe, 1876.
fulvipes Pascoe. C. 455.
rubricalis Broun. C. 452.
stramineipes Broun. C. 862.
ferruginea Broun. C. 862.
compta Broun. C. 934.
erubescens Broun. B. I, i, 68.
rufa Broun. B. I, vi, 564.
oblonga Broun. B. I, vi, 565.
rostralis Broun. B. I, vi, 565.
conspersa Broun. B. I, vi, 566.
spinifera Broun. T. xlvi, 124.
Tysius Pascoe, 1875.
amplipennis Pascoe. C. 458.
purus Broun. C. 1224.
Cyrtalia Pascoe, 1873.
dispar Broun. C. 907.
Caenophanus Broun, 1886.
flavipilus Broun. C. 796.
carbonarius Broun. B. I, vii, 645.
Colabotelus Broun, 1914.
dealbatus Broun. B. I, iii, 232.
Hypotagea Pascoe, 1876.
rubido Pascoe. C. 456.
testaceipennis Broun. C. 456.
variegata Broun. C. 456.
creperus Broun. C. 706.
simulans Broun. C. 706.
castanea Broun. C. 714.
vestita Broun. C. 715.
dissoma Broun. C. 863.
tibialis Broun. C. 1376, and B. I, vi,
 567.
lewi Broun. T. xlvi, 131.
Eugnomus Schönherr, 1847.
elegans Pascoe. C. 458.
picipennis Pascoe. C. 459.

- Eugnomus*—continued.
fervidus Pascoe. C. 459.
wakefieldii Pascoe. C. 459.
fucosus Pascoe. C. 459.
interstitialis Broun. C. 460.
discolor Broun. C. 460.
cinnamoneus White. C. 461.
maculosus Broun. C. 715.
nubilans Broun. C. 715.
fasciatus Broun. C. 716.
luctuosus Broun. C. 863.
monachus Broun. C. 966.
argutus Sharp. C. 1225.
nobilis Broun. C. 1225.
aenescens Broun. C. 1226.
cyaneus Broun. C. 1227.
tarsalis Broun. C. 1227.
lituratus Broun. C. 1227.
aspersus Broun. C. 1376.
dennanensis Broun. T. xlvi, 125.
spuamifer Broun. C. 1377.
maurus Broun. C. 1377.
antennalis Broun. A. 8, iv, 133.
femoralis Broun. A. 8, iv, 134.
albisetosus Broun. B. I, vii, 645.
bryobius Broun. B. I, v, 445.
alternans Broun. B. I, v, 445.
tristis Broun. B. I, v, 446.
atratus Broun. B. I, vi, 566.
calvulus Broun. T. xlvi, 125.
Oreocaris Broun, 1886.
nigriceps Broun. C. 864.
pictularis Broun. C. 865.
nigrescens Broun. C. 865.
cyanea Broun. C. 716.
vittata Broun. C. 1228.
dealbata Broun. C. 1228.
ferruginea Broun. A. 7, xiv, 120.
pullata Broun. A. 7, xiv, 121.
carinulata Broun. B. I, iii, 233.
latipennis Broun. B. I, iii, 234.
lineirostris Broun. B. I, iii, 234.
pleuralis Broun. B. I, iv, 337.
fasciata Broun. B. I, v, 447.
congruens Broun. B. I, v, 447.
albosparsa Broun. T. xlvi, 126.
veronicae Broun. T. xlvi, 127.
picipennis var. Broun. T. xlvi, 127.
uniformis Broun. T. xlvi, 127.
dives Broun. T. xlvi, 128.
castanea Broun. T. xlvi, 128.
Hoplocneme White, 1846.
hookeri White. C. 461.
punctatissima Pascoe. C. 461.
spuamosa Broun. C. 462.
inaequalis Broun. C. 1229.
cyanea Broun. C. 1230.
propinquua Broun. B. I, iii, 233.
vicina Broun. T. xlvi, 129.
Stephanorhynchus White, 1846.
curvipes White. C. 462.
purus Pascoe. C. 463.
brevipennis Pascoe. C. 463.
lawsoni Sharp. C. 463.
crassus Broun. C. 464.

Group RHYNCHOPHORA—continued.

- Stephanorhynchus*—continued.
tuberosus Broun. C. 717.
fatus Broun. C. 718.
griseipictus Broun. C. 866.
nigrosparsus Broun. C. 1230.
insolitus Broun. C. 1231.
costifer Broun. C. 1231.
attelaboides Fabricius. *Ent. Syst.*, 156.
pygmaeus Broun. A. 7, xii, 80.
osculator Broun. A. 8, iv, 134.
morosus Broun. A. 8, iv, 135.
halli Broun. B. 1, iii, 235.
- Stenopactola* Broun, 1914.
proliza Broun. B. 1, iii, 237.
- Pactola* Pascoe, 1876.
variabilis Pascoe. C. 465.
demissa Pascoe. C. 465.
humeralis Broun. A. 6, xv, 407.
nitidula Broun. T. xlvi, 129.
fusicornis Broun. T. xlvi, 130.
binodiceps Broun. T. xlvi, 130.
- Pactolotyphus* Broun, 1909.
striatus Broun. S. 119.
- Brexius* Pascoe, 1870.
ascitus Pascoe. C. 465.
- Subfamily APIONINAE.
- Apion* Herbst, 1797.
metrosideros Broun. C. 466.
terricola Broun. B. 1, viii, 702.
- Subfamily RHINOMACERINAE.
- Rhinorhynchus* Sharp, 1882.
rufulus Broun. C. 467.
- Subfamily BELINAE.
- Pachyura* Hope, 1833.
metallica Pascoe. C. 469.
rubicunda Broun. C. 469.
sumptuosa Broun. C. 470.
albocoma Broun. C. 1233.
stictica Broun. C. 1379.
venusta Broun. A. 8, iv, 136.
violacea Broun. A. 8, iv, 137.
aeneescens Broun. B. 1, iv, 338.
brookesi Broun. B. 1, vi, 567.
Agathinus Broun, 1880.
tridens Fabricius. C. 471.
- Subfamily SCOLOPTERINAE.
- Scolopterus* White, 1846.
tetracanthus White. C. 472.
aequus Broun. C. 473.
penicillatus White. C. 473.
pectoralis Broun. C. 474.
aeneorufus Broun. C. 718.
- Icmalus* Broun, 1893.
abnormis Broun. C. 908.
helmsi Sharp. *Ent. Mon. Mag.*, xviii, 47.
- Nyxetes* Pascoe, 1870.
bidens Fabricius. C. 474.
rufipes Broun. C. 718.
- Ancistropterus* White, 1846.
quadrispinosus White. C. 475.
brouni Sharp. C. 475.
pilosus Broun. C. 476.
- Gonopterus* Broun, 1904.
spinicollis Broun. A. 7, xiv, 122.
- Amlyopterus* Broun, 1886.
prasinus Broun. C. 797.
- Oropterus* White, 1846.
coniger White. C. 477.
- Subfamily TYCHINAE.
- Sibinia* Germar, 1824.
tychoides Pascoe. C. 478.
- Subfamily HAPLONYCHINAE.
- Geochus* Broun, 1882.
inaequalis Broun. C. 446.
politus Broun. C. 704.
rugulosus Broun. C. 931.
similis Broun. C. 1221.
nodosus Broun. C. 1221.
plagiatus Broun. C. 1365.
puncticollis Broun. C. 1367.
squamosus Broun. C. 1465.
tibialis Broun. C. 1465.
marginatus Broun. C. 1466.
frontalis Broun. C. 1466.
nigripes Broun. C. 1467.
setiger Broun. A. 6, xii, 295.
sulcatus Broun. B. 1, ii, 129.
morosus Broun. B. 1, ii, 130.
pyriformis Broun. B. 1, iii, 221.
suffusus Broun. B. 1, iii, 222.
variegatus Broun. B. 1, iii, 223.
distinguens Broun. B. 1, iii, 223.
lateralis Broun. B. 1, iii, 224.
pictulus Broun. B. 1, vi, 568.
apicalis Broun. B. 1, vi, 569.
certus Broun. B. 1, vi, 570.
convexus Broun. B. 1, vi, 570.
posticalis Broun. T. xlvi, 121.
rufipictus Broun. B. 1, viii, 696.
- Subfamily CRYPTORHYNCHINAE.
- Psepholax* White, 1846.
sulcatus White. C. 479.
granulatus Broun. C. 479.
cornutus Broun. C. 479.
barbifrons White. C. 480.
punctulatus Broun. C. 480.
rostralis Broun. C. 480.
coronatus White. C. 481.
femoratus Broun. C. 481.
simplex Pascoe. C. 481.
mystacinus Broun. C. 866.
mediocris Broun. C. 909.
brevicornis Broun. C. 1382.
crassicornis Broun. A. 6, xv, 408.
denticostatus Broun. A. 8, iv, 139.
acanthomerus Broun. T. xlvi, 132.

Group RHYNCHOPHORA—continued.

- Aphocoelus* Broun, 1909.
versicolor Broun. A. 8, iv, 138.
Pseudoreda Broun, 1893.
tibialis Broun. C. 1383 and 482.
Oreda White, 1846.
notata White. C. 487.
brevis Pascoe. C. 487.
Kentraulax Broun, 1909.
murina Broun. C. 487.
flavisetosus Broun. T. xlili, 106.
Mesoreda Broun, 1893.
setigera Broun. C. 1235 and 488.
orthorhina Broun. C. 909.
sulcifrons Broun. A. 8, iv, 155.
longula Broun. T. xlvi, 133.
laminata Broun. B. 1, vii, 646.
Homoreda Broun, 1893.
punctata Broun. C. 1383.
Aldonus White, 1846.
hylobioides White. C. 483.
celator Pascoe. C. 483.
rostratus Broun. C. 483.
chathamensis Sharp. T. xli, 149.
misturatus Broun. T. xlili, 309.
lineifer Broun. T. xlili, 310.
Nothaldonus Broun, 1893.
peacei Broun. C. 484.
Indacentia Broun, 1880.
nubila Broun. C. 485.
straminea Broun. C. 486.
Ectopsis Broun, 1881.
ferrugalis Broun. C. 719.
simplex Broun. A. 6, xii, 383.
foveigerus Broun. B. 1, v, 448.
Tychanus Pascoe, 1876.
gibbus Pascoe. C. 498.
ferrugatus Pascoe. C. 498.
densus Broun. C. 499.
verrucosus Pascoe. C. 499.
lachrymosus Broun. C. 729.
scabrosus Broun. C. 801.
quadratus Broun. C. 867.
ventralis Broun. C. 935.
bufo Sharp. C. 1239.
dux Broun. C. 1240.
squamulosus Broun. B. 1, iii, 239.
lunulus Broun. B. 1, v, 448.
obscurus Broun. B. 1, vi, 573.
costatus Broun. T. xlvi, 137.
complexus Broun. B. 1, vii, 648.
simulans Broun. B. 1, vii, 648.
terricola Broun. B. 1, vii, 649.
Clypeolus Broun, 1909.
cineraceus Broun. A. 8, iv, 143.
Euacalles Broun, 1919.
cristatus Broun. B. 1, vii, 647.
Sympedius Pascoe, 1876.
testudo Pascoe. C. 500.
verzatus Pascoe. C. 500.
curtus Broun. C. 728.
lepidus Broun. C. 935.
rectirostris Broun. A. 8, iv, 149.
minor Broun. B. 1, vii, 651.
- Crisius* Pascoe, 1876.
binotatus Pascoe. C. 500.
variegatus Broun. C. 501.
scutellaris Broun. C. 501.
obesus Sharp. C. 1241.
picicollis Broun. C. 1241.
signatus Broun. C. 1242.
ornatus Broun. C. 1242.
fasciculatus Broun. C. 1492.
dorsalis Broun. A. 7, xiv, 123.
latirostris Broun. B. 1, iii, 240.
posticalis Broun. B. 1, v, 450.
contiguus Broun. B. 1, vi, 574.
eximius Broun. B. 1, vi, 574.
humeralis Broun. T. xlvi, 138.
semifuscus Broun. T. xlvi, 139.
decorus Broun. T. xlvi, 139.
dives Broun. B. 1, vii, 650.
longulus Broun. B. 1, vii, 650.
Agacalles Broun, 1886.
formosus Broun. C. 967.
Getacalles Broun, 1893.
rostralis Broun. C. 1380.
minor Broun. C. 1381.
humeralis Broun. C. 1381.
inaequalis Broun. C. 1493.
sparsus Broun. B. 1, ii, 135.
fasciatus Broun. B. 1, ii, 136.
favosus Broun. A. 8, iv, 157.
posticalis Broun. B. 1, iii, 241.
fulvicornis Broun. B. 1, iii, 242.
parvulus Broun. B. 1, iii, 242.
variellus Broun. B. 1, iii, 243.
oblongus Broun. B. 1, iii, 243.
baccatellus Broun. B. 1, v, 449.
foveiceps Broun. B. 1, v, 450.
substriatus Broun. T. xlvi, 147.
eucoelius Broun. B. 1, vii, 652.
fulvispareus Broun. B. 1, vii, 653.
grisealis Broun. B. 1, vii, 653.
Omeacalles Broun, 1909.
perspicuus Broun. A. 8, iv, 151.
Tychanopais Broun, 1893.
pictulus Broun. C. 1380.
flavisparsus Broun. T. xlvi, 140.
dealbatus Broun. B. 1, vii, 654.
Atyloides Broun, 1914.
foveiger Broun. B. 1, ii, 137.
asaphus Broun. B. 1, vi, 577.
Dendrostygynus Broun, 1895.
calcaratus Broun. A. 6, xv, 410.
Hadracalles Broun, 1893.
fuliginosus Broun. C. 1384.
Pachyderris Broun, 1909.
punctiventris Broun. S. 121.
Acalles Schönherr, 1826.
intutus Pascoe. C. 489.
erroneus Pascoe. C. 489.
hystriculus Pascoe. C. 489.
impexus Pascoe. C. 490.
perpusillus Pascoe. C. 490.
trinotatus Broun. C. 490.
pascoei Broun. C. 491.
signatus Broun. C. 491.

Group RHYNCHOPHORA—continued.

Acalles—continued.

- tortipes* Broun. C. 492.
crisioides Broun. C. 492.
vividus Broun. C. 493.
certus Broun. C. 493.
scitus Broun. C. 494.
ovatellus Broun. C. 720.
laeviculus Broun. C. 721.
cristatus Broun. C. 721.
rudis Broun. C. 721.
dorsalis Broun. C. 722.
volens Broun. C. 722.
horridus Broun. C. 723.
rubricus Broun. C. 723.
spurcus Broun. C. 724.
mundus Broun. C. 724.
canescens Broun. C. 725.
artetus Broun. C. 725.
vafrus Broun. C. 725.
griseus Broun. C. 797.
diversus Broun. C. 798.
cingulatus Broun. C. 798.
latirostris Broun. C. 799.
decemcristatus Broun. C. 800.
huttoni Broun. C. 801.
setifer Broun. C. 867.
posticalis Broun. C. 868.
terrictola Broun. C. 934.
floricola Broun. C. 966.
veratrus Broun. C. 1235.
maritimus Broun. C. 1236.
cryptobius Broun. C. 1236.
ingens Broun. C. 1236.
xanthostictus Broun. C. 1237.
adamsi Broun. C. 1237.
concininus Broun. C. 1238.
australis Broun. C. 1239.
mimus Broun. C. 1385.
lepirhinus Broun. C. 1385.
sympesioides Broun. C. 1386.
dolosus Broun. C. 1484.
comptus Broun. C. 1486.
facilis Broun. C. 1486.
quietus Broun. C. 1487.
integer Broun. A. 6, xii, 374.
ruficollis Broun. A. 6, xii, 376.
porcatus Broun. A. 6, xii, 377.
puncticollis Broun. A. 6, xii, 378.
farinosus Broun. A. 6, xii, 378.
fougeri Hutton. T. xxx, 157.
fuscatus Broun. A. 7, xix, 60.
multisetosus Broun. A. 7, xix, 61.
flynni Broun. B. 1, ii, 131.
notoporhinus Broun. B. 1, ii, 132.
fuscidorsis Broun. A. 8, iv, 143.
igneus Broun. A. 8, iv, 144.
altus Broun. A. 8, iv, 145.
albistrigalis Broun. A. 8, iv, 146.
aterrimus Broun. A. 8, iv, 146.
praesetosus Broun. A. 8, iv, 147.
robustus Broun. A. 8, iv, 148.
flavisetosus Broun. A. 8, iv, 149.
confusus Broun. B. 1, iii, 237.
bicristiceps Broun. B. 1, iii, 238.
ignotus Broun. B. 1, iii, 239.
furvus Broun. B. 1, iv, 339.

Acalles—continued.

- bicinctus* Broun. B. 1, iv, 339.
sternalis Broun. B. 1, v, 452.
nodigerus Broun. B. 1, v, 452.
zenomorphus Broun. B. 1, v, 454.
dentigerus Broun. B. 1, v, 455.
humeralis Broun. B. 1, vi, 571.
anceps Broun. B. 1, vi, 572.
cilicollis Broun. B. 1, vi, 573.
conicollis Broun. T. xlvi, 134.
eruensis Broun. T. xlvi, 134.
peleensis Broun. T. xlvi, 135.
consors Broun. T. xlvi, 136.
gracilis Broun. T. xlvi, 136.
contractus Broun. T. xlvi, 137.
lineirostris Broun. T. xlvi, 107.
subcarinatus Broun. T. xlvi, 107.
piciventris Broun. S. 120.
brevipennis Broun. B. 1, vii, 655.
hopensis Broun. B. 1, vii, 655.
bicostatus Broun. B. 1, vii, 656.
binodes Broun. B. 1, vii, 651.
sticticus Broun. B. 1, vii, 658.
brookesi Broun. B. 1, viii, 703.
Torilus Broun, 1909.
griseocollis Broun. A. 8, iv, 152.
Hatasu Broun, 1909.
dorsale Broun. A. 8, iv, 141.
tuberosa Broun. B. 1, viii, 704.
Onias Broun, 1909.
latisulcatus Broun. A. 8, iv, 153.
sentus Broun. C. 799.
ornatus Broun. A. 8, iv, 154.
albicristus Broun. B. 1, iii, 244.
irregularis Broun. T. xlvi, 145.
Allancalcis Broun, 1912.
aulacus Broun. C. 1489.
allostethus Broun. C. 1490.
incultus Broun. C. 1490.
formosus Broun. A. 6, xii, 375.
ignealis Broun. T. xlvi, 142.
oculatus Broun. T. xlvi, 142.
dilatatus Broun. T. xlvi, 143.
laticollis Broun. B. 1, ni, 245.
variatus Broun. B. 1, vi, 575.
seticollis Broun. B. 1, vi, 575.
albipictus Broun. B. 1, vi, 576.
melastictus Broun. B. 1, vi, 576.
blanditus Broun. B. 1, vi, 577.
Acallopa Pascoe, 1877.
rudis Pascoe. C. 495.
sculpturatus Broun. C. 495.
Scelodolichus Broun, 1880.
celsus Broun. C. 496.
lineithorax Broun. C. 496.
setosus Broun. C. 726.
villosus Broun. C. 726.
denotans Broun. C. 726.
attulus Broun. C. 968.
juncobius Broun. C. 1239.
halaris Broun. C. 1491.
politus Broun. A. 6, xv, 412.
quamulosus Broun. A. 6, xv, 413.
flexicollis Broun. B. 1, ii, 134.
decorus Broun. B. 1, viii, 703.
pyriformis Broun. B. 1, viii, 704.

Group RHYNCHOPHORA—continued.

- Zeacalles* Broun, 1893.
flavescens Broun. A. 6, xii, 380.
varius Broun. A. 6, xii, 381.
alpestris Broun. C. 1488.
picatus Broun. C. 1488.
binodosus Broun. B. 1, i, 69.
lepidulus Broun. A. 8, iv, 140.
carinellus Broun. B. 1, iii, 245.
estriatus Broun. B. 1, iii, 246.
scaber Broun. B. 1, iv, 340.
sparsus Broun. B. 1, iv, 341.
speciosus Broun. B. 1, v, 452.
aeratus Broun. B. 1, vi, 578.
coarctalis Broun. B. 1, vi, 578.
finitimus Broun. B. 1, vi, 579.
bisulcatus Broun. B. 1, vi, 579.
scruposus Broun. B. 1, vi, 580.
latulus Broun. B. 1, vi, 581.
inornatus Broun. B. 1, vi, 581.
parvus Broun. B. 1, vi, 582.
cordipennis Broun. B. 1, vi, 582.
pictus Broun. T. xliv, 144.
femoralis Broun. T. xliv, 145.
- Cyclacalles* Broun, 1880.
setiger Broun. C. 497.
vestitus Broun. C. 497.
- Xenacalles* Broun, 1912.
triangulatus Broun. C. 800.
squamiventris Broun. T. xliii, 108.
simplex Broun. B. 1, ii, 133.
nodifer Broun. B. 1, ii, 134.
nigricans Broun. B. 1, v, 451.
- Schylus* Broun, 1895.
nigricollis Broun. A. 6, xv, 412.
- Metacalles* Broun, 1893.
aspersus Broun. A. 6, xii, 382.
rugicollis Broun. A. 6, vii, 383.
cordipennis Broun. C. 720.
latus Broun. C. 727.
crinitus Broun. C. 727.
exiguus Broun. C. 728.
picatus Broun. B. 1, ii, 135.
crinitus Broun. T. xlvi, 143.
lanosus Broun. T. xlvi, 144.
- Dermothrius* Broun, 1882.
sanguineus Broun. C. 730.
- Euthyrrhinus* Schonherr, 1837.
squamiger White. C. 503.
- Baeorhynchodes* Broun, 1909.
cristatus Broun. A. 8, iv, 159.
- Rhyncodes* White, 1846.
ursus White. C. 502.
saundersi White. C. 503.
squamifer Broun. C. 730.
atrus Broun. C. 802.
weberi Colenso. T. xiv, 281.
rubipunctatus Colenso. T. xiv, 282.
- Mistrastethus* Redtenbacher, 1868.
bituberculatus Fabricius. C. 504.
- Paranomocerus* Redtenbacher, 1868.
spiculus Redtenbacher. C. 505.
maurus Broun. C. 505.
- Idotasia* Pascoe, 1871.
egena Pascoe. C. 506.
- Mecistostylus* Lacordaire, 1866.
douei Lacordaire. Gen. Col. vii.
- Sympiezoscelus* Waterhouse, 1853.
spencei Waterhouse. E. 2, ii, 204.
- Gonipterus.*
**reticulatus* Boisduval.
- Calandra* Clairville.
**granaria* Linnaeus.
**oryzae* Linnaeus.
- Oxyops.*
**concreta* Pascoe.
- Sitones.*
**lineatus* Linnaeus.

Subfamily COSSONINAE.

PENTARTHRIDES.

- Pentarthrum* Wollaston, 1854.
zealandicum Wollaston. T. xli, 156.
amicum Broun. T. xli, 156.
fultoni Broun. T. xli, 156.
proximum Broun. T. xli, 156.
brevirostre Sharp. T. xli, 157.
ruficorne Broun. T. xli, 157.
philotti Broun. T. xli, 157.
melanosternum Broun. T. xli, 157.
punctirostre Broun. T. xli, 157.
planicolle Broun. T. xli, 158.
subsericatum Wollaston. T. xli, 157.
assimilatum Broun. T. xli, 158.
reductum Broun. T. xli, 158.
brunneum Broun. T. xli, 158.
nubilum Broun. T. xli, 158.
breviscorne Broun. B. 1, iii, 246.
impressum Broun. T. xlvi, 147.
tenebrosum Broun. T. xlvi, 148.
dissimilum Broun. T. xliii, 109.
auripilum Broun. T. xliii, 109.
- Touropsis* Broun, 1908.
punctatus Broun. T. xli, 159.
- Euophryum* Broun, 1908.
rufum Broun. T. xli, 160.
asperillum Broun. T. xli, 160.
sculpturatum Broun. T. xli, 160.
punctatissimum Broun. T. xli, 160.
dubitans Broun. T. xli, 161.
confinum Broun. T. xli, 161.
antennale Broun. T. xli, 161.
crassellum Broun. T. xli, 161.
servulum Broun. T. xli, 161.
porcatum Sharp. T. xli, 161.
- Zenoteratus* Broun, 1908.
macrocephalus Broun. T. xli, 162.
diversus Broun. T. xli, 162.
cephalotes Sharp. T. xli, 162.
- Torostomo* Broun, 1908.
apicale Broun. T. xli, 163.
- Toura* Broun, 1908.
longirostre Wollaston. T. xli, 164.

Group RHYNCHOPHORA—continued.

Toura—continued.

- fulva* Broun. T. xli, 164.
morosa Broun. T. xli, 165.
Merisma Broun, 1908.
sharpiana Wollaston. T. xli, 165.
aurantiaca Broun. T. xli, 166.
Tanysoma Broun, 1908.
angusta Broun. T. xli, 167.
impresella Broun. B. 1, iii, 247.
aciphyllae Broun. B. 1, iii, 248.

Stenotoura Broun, 1908.

- exilis* Broun. T. xli, 167.
lateritia Broun. T. xli, 168.
prolifica Broun. T. xli, 168.

Eutassa Broun, 1908.

- comatum* Broun. T. xli, 169.
fuscollis Broun. T. xli, 169.

Adel Broun, 1908.

- crenatus* Broun. T. xli, 170.

Rhinanisus Broun, 1886.

- fulvicornis* Broun. T. xli, 171.
parvicornis Sharp. T. xli, 171.
contiguus Broun. T. xli, 172.
sagax Broun. T. xli, 172.
cheesemani Broun. T. xli, 172.
lewisi Broun. T. xli, 172.
elongatus Broun. T. xli, 172.
subconvexus Broun. T. xli, 172.
suturalis Broun. T. xli, 172.
confertus Sharp. T. xli, 173.
constrictus Sharp. T. xli, 173.
sulcirostris Broun. T. xli, 173.
halli Broun. B. 1, iii, 248.
fusiformis Broun. B. 1, iii, 249.

Campotoscapus Broun, 1893.

- sanguineus* Broun. T. xli, 174.
planiusculus Broun. T. xli, 174.
conicollis Broun. T. xli, 174.

Macrosyntalus Broun, 1886.

- laticollis* Broun. T. xli, 175.
depressus Broun. T. xli, 175.
russulus Broun. T. xli, 176.
remotus Sharp. T. xli, 176.
frontalis Broun. T. xli, 176.
badius Broun. T. xli, 176.
crenatus Broun. T. xli, 176.

Baeorhopalus Broun, 1886.

- glabrus* Broun. T. xli, 177.

Selocomis Broun, 1908.

- aeneopiceus* Broun. T. xli, 178.

Glyphoramphus Broun, 1886.

- rarus* Broun. T. xli, 179.

Belka Broun, 1908.

- spadicea* Broun. T. xli, 180.

Attarus Broun, 1908.

- tristis* Broun. T. xli, 181.
castus Broun. T. xli, 181.
vestitus Broun. T. xli, 181.

Agastegnus Broun, 1886.

- ruficollis* Broun. T. xli, 182.
gratus Broun. T. xli, 183.
longipes Broun. T. xli, 183.
simulans Sharp. T. xli, 183.

Agastegnus—continued.

- coloratus* Broun. T. xli, 183.
femoralis Broun. T. xli, 183.
nitidurostris Broun. T. xli, 183.
distinctus Broun. T. xli, 183.
sericatus Broun. T. xli, 184.
rufescens Broun. T. xli, 184.
biimpressus Broun. T. xli, 184.
concinnus Broun. B. 1, iii, 251.
thoracicus Broun. B. 1, iii, 251.
rugipennis Broun. B. 1, iii, 252.
ornatus Broun. T. xlii, 110.

Sericotrogus Wollaston, 1873.

- suboenezensis* Wollaston. T. xli, 185.
ovicollis Broun. T. xli, 185.
stramineus Broun. T. xli, 185.
plexus Broun. B. 1, iii, 250.
Gauromydas Broun, 1908.

auricomus Broun. T. xli, 186.*mirandus* Broun. B. 1, iii, 253.

- Eucossonus* Broun, 1886.
comptus Broun. T. xli, 187.
elegans Broun. T. xli, 187.
gracilis Broun. T. xli, 187.
setiger Sharp. T. xli, 188.
rostralis Broun. T. xli, 188.
antennalis Broun. B. 1, i, 70.
discalis Broun. B. 1, i, 71.
constrictus Broun. B. 1, vi, 583.
sulcicollis Broun. B. 1, vi, 583.
nasalis Broun. B. 1, vi, 584.
disparilis Broun. B. 1, vi, 584.
oreoculus Broun. B. 1, vi, 585.

Agrilochilus Broun, 1880.

- prolixus* Broun. T. xli, 189.

Dioedimorpha Broun, 1886.

- wollastonianum* Sharp. T. xli, 189.
debile Sharp. T. xli, 190.

Arecocryptus Broun, 1903.

- bellus* Broun. T. xli, 190.

Entium Sharp, 1878.

- aberrans* Sharp. T. xli, 191.

Mesoxenophasis Wollaston, 1874.

- brouni* Wollaston. T. xli, 192.

Proconus Broun, 1886.

- asperirostris* Broun. T. xli, 192.

- crassipes* Broun. T. xli, 192.

Heteropsis Wollaston, 1873.

- lawsoni* Wollaston. T. xli, 193.

Novitas Broun, 1880.

- rufum* Broun. T. xli, 194.

- nigrans* Broun. T. xli, 194.

- dispar* Broun. T. xli, 194.

Unas Broun, 1908.

- piceus* Broun. T. xli, 195.

Trachyglyphus Broun, 1908.

- rugirostris* Broun. T. xli, 196.

Microtribus Wollaston, 1873.

- huttoni* Wollaston. T. xli, 197.

- pictonensis* Sharp. T. xli, 197.

Idus Broun, 1893.

- caecus* Broun. T. xli, 198.

Progonum Broun, 1908.

- helmsianum* Sharp. T. xli, 199.

Group RHYNCHOPHORA—continued.

COSSONIDAE.

- Phloeophagosoma* Wollaston, 1873.
corvinum Wollaston. T. xli, 200.
thoracicum Wollaston. T. xli, 200.
dilutum Wollaston. T. xli, 200.
pedatum Wollaston. T. xli, 200.
rugipenne Broun. T. xli, 200.
abdominale Broun. T. xli, 201.
Eutornus Wollaston, 1873.
dubius Wollaston. T. xli, 202.
vicinus Broun. T. xli, 202.
breviceps Broun. T. xli, 202.
amplus Broun. T. xli, 203.
littoralis Broun. T. xli, 203.
cylindricus Broun. T. xli, 203.
parvulus Broun. T. xli, 203.
Stilbocara Broun, 1893.
nitida Broun. T. xli, 204.
constricticollis Broun. T. xli, 204.
serena Broun. T. xli, 204.
Arecophaga Broun, 1880.
varia Broun. T. xli, 205.
Pogonorhinus Broun, 1903.
oparus Broun. T. xli, 207.
Exomesites Broun, 1886.
optimus Broun. T. xli, 208.
Allaorus Broun, 1893.
urquharti Broun. T. xli, 209.
pedatus Broun. T. xli, 209.
sternalis Broun. T. xli, 209.
ovatus Broun. T. xli, 209.
versutus Broun. T. xli, 210.
rugosus Broun. T. xli, 210.
pyriformis Broun. T. xli, 210.
piciclavus Broun. T. xli, 210.
scutellaris Broun. B. 1, ii, 138.
impressus Broun. B. 1, v, 455.
carinifer Broun. B. 1, vi, 586.
Pselactus Broun, 1886.
punctatus Broun. T. xli, 212
ferrugineus Broun. T. xli, 212.

Inosomus Broun, 1903.

- rufopiceus* Broun. T. xli, 213.
Xenonema Wollaston, 1873.
spinipes Wollaston. T. xli, 214.
Hectoetus Broun, 1904.
rubidus Broun. T. xli, 215.

Family SCOLYTIDAE.

Subfamily SCOLYTINAE.

- Pachycotes* Sharp, 1877.
ventralis Sharp. C. 538.
Hylastes Erichson, 1836.
peregrinus Chapuis. C. 539.
Acrantus Broun, 1903 (substituted for
Homarus, 1881).
mundulus Broun. C. 740.
opus Broun. A. 6, xv, 417.
Dendrotrupes Broun, 1881.
vestitus Broun. C. 741.
costiceps Broun. C. 741.
Tomicus Latreille, 1810.
asper Broun. C. 742.
Mesoscolytus Broun, 1903 (substituted for
Apate).
inurbanus Broun. C. 346, and A. 7,
 xiv, 126.

Subfamily PLATYPIDAE.

- Platypus* Herbst, 1793.
douei Reich. C. 540.
apicalis White. C. 541.
caviceps Broun. C. 541.
castaneus Broun. C. 542.
lobatus Broun. C. 1253.
gracilis Broun. C. 1254.
inimicus Broun. B. 1, i, 71.

Family AGLYCYDERIDAE.

- Aglycyderes* Westwood.
wollastoni Sharp. C. 426.
badius Broun. C. 427.