

**TURKISH RED LIST CATEGORIES OF
LONGICORN BEETLES (COLEOPTERA: CERAMBYCIDAE)
PART VI – SUBFAMILY CERAMBYCINAE: ACHRYSONINI,
HESPEROPHANINI, PHORACANTHINI, CERAMBYCINI,
ROSALIINI, TRACHYDERINI AND CALLICHRMATINI**

Hüseyin Özdikmen*

* Gazi University, Faculty of Science, Department of Biology, 06500 Ankara, TURKEY. E-mail: ozdikmen@gazi.edu.tr

[**Özdikmen, H.** 2014. Turkish Red List Categories of Longicorn Beetles (Coleoptera: Cerambycidae) Part VI – Subfamily Cerambycinae: Achrysonini, Hesperophanini, Phoracanthini, Cerambycini, Rosaliini, Trachyderini and Callichromatini. *Munis Entomology & Zoology*, 9 (2): 609-623]

ABSTRACT: The aim of this study is to create a Turkish Red List of the longicorn beetles. Moreover, presence such a Red List is necessary for Turkey. Even governmental evaluations could cause some erroneous decisions due to absence such a Red List. Since, governmental evaluations at the present time are based on the works that are realized with respect to the European Red List. Furthermore, Turkey appears a continental property changeable in very short distances in terms of climatical features and field structures. So, the status of European fauna and the status of Turkish fauna are not the same. Clearly, there is no any work that subjected to create a Turkish Red List except Parts I-V. Hence, a series work is planned with this purpose. This type of study is the sixth attempt for Turkey.

KEY WORDS: Red List, Conservation, Cerambycidae, Turkey

The purpose of the current study was to create a Turkish Red List of longicorn beetles similarly to “European Red List of Saproxyllic Beetles” that was compiled by Ana Nieto & Keith N. A. Alexander and published by IUCN (International Union for Conservation of Nature) in collaboration with the European Union in 2010. “European Red List of Saproxyllic Beetles” includes 153 species within the subfamilies Prioninae, Cerambycinae (including Stenopterinae) and Lamiinae of the European Cerambycidae. In the future, I hope that the present work will be lead to preparation a more comprehensive “Turkish Red List”.

Hence, a series work is planned with this purpose. The present study is attempted as the sixth step of this aim. The previous works are Özdikmen (2014a,b,c,d,e). It should be noted that the using information at the present work on Turkish longicorn beetles are on the base of my personal database. The data of distribution are given on base of Löbl & Smetana (2010, 2011), Danilevsky (2010a,b, 2012a,b,c,d, 2013), Özdikmen (2011) and Miroshnikov (2011). Identification of chorotypes is based on the chorotype classification of the Anatolian fauna, proposed by Vigna Taglianti et al. (1999).

The evaluations of Turkish longicorn beetles at the present work based on “The IUCN Red List Categories” that was presented in Part I (Özdikmen, 2014a).

TURKISH RED LIST FOR STENOPTERINAE

SUBFAMILY CERAMBYCINAE Latreille, 1802: 211

TRIBE ACHRYSONINI Lacordaire, 1868: 203

GENUS ICOSIUM Lucas, 1854: VIII


SPECIES *I. tomentosum* Lucas, 1854: IX

SUBSPECIES *I. t. atticum* Ganglbauer, 1882: 743

According to European Red List of Saproxyllic Beetles, the species was placed in **LC**. It is represented only by the subspecies *I. tomentosum atticum* in Turkey. It probably is known only from SW Anatolia for Turkey. So, Turkish Red List category of the subspecies is **NT**.

Range: SE Europe, Transcaucasia (Azerbaijan), Turkey, Cyprus, Israel, Jordan.

Chorotype: Turano-Mediterranean (Turano-Apenninian)



TRIBE HESPEROPHANINI Mulsant, 1839: 61


GENUS HESPEROPHANES Dejean, 1835: 328

SPECIES *H. sericeus* (Fabricius, 1787: 152)

According to European Red List of Saproxyllic Beetles, the species was placed in **LC**. It probably is rather widely distributed in Anatolia for Turkey. However, Turkish Red List category of the species is **NT** now.

Range: S Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Turkmenistan, Turkey, Iraq, Cyprus, Israel, Jordan, North Africa (Algeria, Egypt, Libya, Morocco, Tunisia).

Chorotype: Turano-Mediterranean



GENUS TRICHOFERUS Wollaston, 1854: 427


SPECIES *T. fasciculatus* (Faldermann, 1837: 266)

SUBSPECIES *T. f. fasciculatus* (Faldermann, 1837: 266)

According to European Red List of Saproxyllic Beetles, the species was placed in **LC**. It is represented only by the nominative subspecies in Turkey. It probably is rather widely distributed in Anatolia for Turkey. So, Turkish Red List category of the subspecies is **LC**.

Range: S Europe, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Turkey, Iraq, Cyprus, Syria, Lebanon, Israel, Jordan, North Africa (Algeria, Egypt, Libya, Morocco, Tunisia).

Chorotype: Turano-Mediterranean




SPECIES *T. fissitarsis* Sama, Fallahzadeh & Rapuzzi, 2005: 125

European Red List of Saproxyllic Beetles does not include the species rightly. It is known only from E Anatolia. So, Turkish Red List category of the species is **VU**.

Range: Turkey, Iran, Iraq.

Chorotype: SW-Asiatic




SPECIES *T. griseus* (Fabricius, 1792: 325)

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It probably is rather widely distributed in SC and SW Anatolia for Turkey. So, Turkish Red List category of the species is **LC**.

Range: S Europe, Turkey, Iraq, Cyprus, Syria, Lebanon, Israel, Jordan, North Africa (Algeria, Egypt, Libya, Morocco, Tunisia).

Chorotype: Mediterranean




SPECIES *T. holosericeus* (Rossi, 1790: 153)

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It probably is rather widely distributed in N and W Anatolia for Turkey. However, Turkish Red List category of the species is **NT** now.

Range: E Europe (Ukraine), Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey, North Africa (Algeria, Libya, Morocco, Tunisia).

Chorotype: Turano-Mediterranean




SPECIES *T. kotschy* (Ganglbauer, 1883: 300)

European Red List of Saproxylic Beetles does not include the species rightly. It is known only from E Anatolia. So, Turkish Red List category of the species is **NT**.

Range: E Europe (Greece: Samos), Turkey, Syria, Lebanon.

Chorotype: E-Mediterranean




SPECIES *T. lunatus* (Szallies, 1994: 261)

European Red List of Saproxylic Beetles does not include the species rightly. It is known only from SE Anatolia. So, Turkish Red List category of the species is **VU**.

Range: Turkey, Iran.

Chorotype: SW-Asiatic (Irano-Anatolian)




SPECIES *T. pallidus* (Olivier, 1790: 256)

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It is known only from SW Anatolia for Turkey. So, Turkish Red List category of the species is **EN** now.

Range: C and E Europe, Caucasus, Turkey.

Chorotype: C and E European




SPECIES *T. preissi* (Heyden, 1894: 85)

European Red List of Saproxylic Beetles does not include the endemic species rightly. It is known only from S and SW Anatolia. So, Turkish Red List category of the species is **NT**.

Range: Turkey.

Chorotype: Anatolian




SPECIES *T. samai* Kadlec & Rejzek, 2001: 296

European Red List of Saproxylic Beetles does not include the endemic species rightly. It is known only from the type locality in SCE Anatolia. So, Turkish Red List category of the species is **DD**.

Range: Turkey.

Chorotype: Anatolian




SPECIES *T. sbordonii* Sama, 1982: 217

European Red List of Saproxylic Beetles does not include the endemic species rightly. It is known only from the type locality in SW Anatolia. So, Turkish Red List category of the species is **DD**.

Range: Turkey.


Chorotype: Anatolian

**SPECIES *T. spartii* (Müller, 1948: 67)**

European Red List of Saproxylic Beetles does not include the endemic species rightly. It is known only from WC Anatolia. So, Turkish Red List category of the species is **VU**.

Range: Turkey.


Chorotype: Anatolian

**GENUS *STROMATIUM* Audinet-Serville, 1834: 80****SPECIES *S. unicolor* (Olivier, 1795: 58)**

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It is widely distributed in Turkey. So, Turkish Red List category of the species is **LC**.

Range: S Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkmenistan, Iran, Turkey, Cyprus, Iraq, Syria, Lebanon, Israel, Jordan, North Africa (Algeria, Libya, Morocco, Tunisia).


Chorotype: Turano-Mediterranean

**TRIBE PHORACANTHINI Newman, 1840: 2****GENUS *PHORACANTHA* Newman, 1840: 19****SPECIES *P. recurva* Newman, 1840: 4**

European Red List of Saproxylic Beetles does not include the species. It is known only from SCW Anatolia. So, Turkish Red List category of the species is **VU**.

Range: S Europe, Turkey, Lebanon, Israel, North Africa (Libya, Morocco, Tunisia), and Afrotropical, Australian, Nearctic and Neotropical Regions.

Chorotype: Subcosmopolitan




SPECIES *P. semipunctata* (Fabricius, 1775: 180)

European Red List of Saproxylic Beetles does not include the species. It is known only from SCW Anatolia. So, Turkish Red List category of the species is **VU**.

Range: S Europe, Turkey, Cyprus, Syria, Lebanon, Israel, Jordan, North Africa (Algeria, Canary Islands, Egypt, Libya, Morocco, Madeira Archipelago, Tunisia), and Afrotropical, Australian, Nearctic and Neotropical Regions.

Chorotype: Subcosmopolitan



TRIBE CERAMBYCINI Latreille, 1802: 211

GENUS *CERAMBYX* Linnaeus, 1758: 388


SUBGENUS *CERAMBYX* Linnaeus, 1758: 388

SPECIES *C. carinatus* (Küster, 1845: 46)

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It is rather widely distributed in W half of Anatolia for Turkey. So, Turkish Red List category of the species is **NT** now.

Range: SE Europe (Italy, Malta, Balkans), Turkey.

Chorotype: Turano-Mediterranean (Turano-Apenninian)




SPECIES *C. cerdo* Linnaeus, 1758: 392

SUBSPECIES *C. c. cerdo* Linnaeus, 1758: 392

According to European Red List of Saproxylic Beetles, the species was placed in **NT**. It is represented only by the nominative subspecies in Turkey. It is widely distributed in Turkey. So, Turkish Red List category of the subspecies is **LC**.

Range: Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Turkey, Iraq, Syria, Lebanon, Israel, Jordan, North Africa (Morocco).

Chorotype: Europeo-Mediterranean




SPECIES *C. dux* (Faldermann, 1837: 264)

According to European Red List of Saproxylic Beetles, the species was placed in NT. It is widely distributed in Turkey. So, Turkish Red List category of the species is LC.

Range: E Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Turkey, Syria, Lebanon, Israel, Jordan.

Chorotype: Turano-Mediterranean (Turano-E-Mediterranean)




SPECIES *C. heinzianus* Demelt, 1976: 65

European Red List of Saproxylic Beetles does not include the endemic species rightly. It is known only from E Anatolia. So, Turkish Red List category of the species is VU.

Range: Turkey.

Chorotype: Anatolian




SPECIES *C. miles* Bonelli, 1812: 178

According to European Red List of Saproxylic Beetles, the species was placed in NT. It probably is rather widely distributed in S and W Turkey especially. So, Turkish Red List category of the species is LC.

Range: C and E Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey.

Chorotype: C and E European




SPECIES *C. nodulosus* Germar, 1817: 220

According to European Red List of Saproxylic Beetles, the species was placed in NT. It probably is rather widely distributed in Turkey. So, Turkish Red List category of the species is LC.

Range: SC and SE Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey, Cyprus, Syria, Lebanon.

Chorotype: Turano-Mediterranean (Turano-Apeninian)



SPECIES *C. welensii* (Küster, 1845: 44)

According to European Red List of Saproxylic Beetles, the species was placed in NT. It probably is rather widely distributed in S and W Turkey especially. So, Turkish Red List category of the species is LC.

Range: S Europe, Transcaucasia (Azerbaijan, Georgia), Iran, Turkey, Cyprus, Syria, Lebanon, Israel, Jordan.

Chorotype: Turano-Mediterranean




SUBGENUS *MICROCERAMBYX* Mikšić & Georgijević, 1973: 22

SPECIES *C. scopolii* Fuessly, 1775: 12

According to European Red List of Saproxylic Beetles, the species was placed in LC. It is widely distributed in Turkey. So, Turkish Red List category of the species is LC.

Range: Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey, Syria.

Chorotype: European




Moreover, the species is represented by two subspecies in Turkey as the nominative subspecies and *C. scopolii nitidus* Pic, 1892.

SUBSPECIES *C. s. scopolii* Fuessly, 1775: 12

According to European Red List of Saproxylic Beetles, the subspecies was placed in LC. It probably is rather widely distributed in N Turkey especially. So, Turkish Red List category of the subspecies is LC.

Range: Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey.

Chorotype: European




SUBSPECIES *C. s. nitidus* Pic, 1892: CXI

European Red List of Saproxylic Beetles does not include the subspecies. It is known only from SC and SW Anatolia. So, Turkish Red List category of the subspecies is **NT**.

Range: Turkey, Syria.

Chorotype: SW-Asiatic



TRIBE ROSALIINI Fairmaire, 1864: 137

GENUS *ROSALIA* Audinet-Serville, 1834: 561


SUBGENUS *ROSALIA* Audinet-Serville, 1834: 561

SPECIES *R. alpina* (Linnaeus, 1758: 392)

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It probably is rather widely distributed in N Turkey especially. However, Turkish Red List category of the species is **NT**.

Range: Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey.

Chorotype: European




Moreover, the species is represented by two subspecies in Turkey as the nominative subspecies and *R. alpina syriaca* Pic, 1895.

SUBSPECIES *R. a. alpina* (Linnaeus, 1758: 392)

According to European Red List of Saproxylic Beetles, the subspecies was placed in **LC**. It probably is rather widely distributed in N Turkey especially. However, Turkish Red List category of the subspecies is **NT**.

Range: Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey.

Chorotype: European




SUBSPECIES *R. a. syriaca* Pic, 1895: CCLXXXV

European Red List of Saproxylic Beetles does not include the endemic subspecies rightly. It is known only from SC Anatolia. So, Turkish Red List category of the subspecies is **EN**.

Range: Turkey.


Chorotype: Anatolian

**TRIBE TRACHYDERINI Dupont, 1836: 1****GENUS *PURPURICENUS* Dejean, 1821: 105****SPECIES *P. apicalis* Pic, 1905: 163**

European Red List of Saproxylic Beetles does not include the species. It is known only from SE Anatolia for Turkey. So, Turkish Red List category of the species is **VU**.

Range: Turkey, Iran, Iraq.


Chorotype: SW-Asiatic

**SPECIES *P. bitlisiensis* Pic, 1902: 27**

European Red List of Saproxylic Beetles does not include the endemic species rightly. It is known only from E Anatolia. So, Turkish Red List category of the species is **EN**.

Range: Turkey.


Chorotype: Anatolian

**SPECIES *P. budensis* (Götz, 1783: 70)**

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It is widely distributed in Turkey. So, Turkish Red List category of the species is **LC**.

Range: S and E Europe, W Siberia, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey, Cyprus, Syria, Lebanon, Israel.

Chorotype: Sibero-European




SPECIES *P. caucasicus* T. Pic, 1902: 27**SUBSPECIES *P. c. caucasicus* T. Pic, 1902: 27**

European Red List of Saproxylic Beetles does not include the species. It is represented only by the nominative subspecies in Turkey. It is known only sporadically from Anatolia. So, Turkish Red List category of the subspecies is **NT** now.

Range: Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Turkey.


Chorotype: SW-Asiatic

**SPECIES *P. cornifrons* Sabbadini & Pesarini, 1992: 58**

European Red List of Saproxylic Beetles does not include the endemic species rightly. It is known only from E Anatolia. So, Turkish Red List category of the species is **VU**.

Range: Turkey.


Chorotype: Anatolian

**SPECIES *P. dalmatinus* Sturm, 1843: 353**

According to European Red List of Saproxylic Beetles, the species was placed in **DD**. It probably is rather widely distributed in S half of Anatolia for Turkey. So, Turkish Red List category of the species is **LC**.

Range: SE Europe, Turkey, Syria, Lebanon, Israel, Jordan.


Chorotype: E-Mediterranean (NE-Mediterranean)

**SPECIES *P. desfontainii* (Fabricius, 1792: 258)****SUBSPECIES *P. d. inhumeralis* Pic, 1891: 24**

According to European Red List of Saproxylic Beetles, the species was placed in **DD**. It is represented only by the subspecies *P. desfontainii inhumeralis* in Turkey. It probably is rather widely distributed in Anatolia for Turkey. So, Turkish Red List category of the subspecies is **LC**.

Range: SE Europe (Greece), Turkey, Syria, Lebanon, Israel, Jordan.

Chorotype: E-Mediterranean (NE-Mediterranean)




SPECIES *P. interscapillatus* Plavilstshikov, 1937: 247 [RN]

European Red List of Saproxylic Beetles, does not include the species. The species is known only from SC and SCW Anatolia for Turkey. So, Turkish Red List category of the species is **NT**.

Range: Turkey, Cyprus, Syria, Lebanon, Israel.

Chorotype: E-Mediterranean (Palestino-Cyprioto-Taurian)




SPECIES *P. kaehleri* (Linnaeus, 1758: 393)

According to European Red List of Saproxylic Beetles, the species was placed in **LC**. It probably is rather widely distributed in N Turkey. However, Turkish Red List category of the species is **NT**.

Range: Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Turkey.

Chorotype: Turano-European or European




Moreover, the species is represented by two subspecies in Turkey as the nominative subspecies and *P. kaehleri menetriesi* Motschulsky, 1845.

SUBSPECIES *P. k. kaehleri* (Linnaeus, 1758: 393)

According to European Red List of Saproxylic Beetles, the subspecies was placed in **LC**. It probably is rather widely distributed in NW Turkey (European Turkey). So, Turkish Red List category of the subspecies is **NT**.

Range: Europe, Caucasus, Turkey.

Chorotype: European




SUBSPECIES *P. k. menetriesi* Motschulsky, 1845: 87

European Red List of Saproxylic Beetles, does not include the subspecies. It is known only from NE Anatolia for Turkey. So, Turkish Red List category of the subspecies is **NT**.

Range: Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Turkey.


Chorotype: SW-Asiatic

**SPECIES *P. nanus* Semenov, 1907: 254**

European Red List of Saproxylic Beetles, does not include the species. It is known only from W Anatolia for Turkey. So, Turkish Red List category of the species is **DD**.

Range: Iran, Turkey.


Chorotype: SW-Asiatic (Irano-Anatolian)

**SPECIES *P. nigrotatus* Pic, 1907: 169**

European Red List of Saproxylic Beetles, does not include the endemic species rightly. It is known only from SC Anatolia. So, Turkish Red List category of the species is **VU**.

Range: Turkey.


Chorotype: Anatolian

**SPECIES *P. nudicollis* Demelt, 1968: 65**

According to European Red List of Saproxylic Beetles, the species was placed in **EN**. It is known only from SC and SCW Anatolia for Turkey. So, Turkish Red List category of the species is **NT**.

Range: Turkey, Cyprus.

Chorotype: E-Mediterranean (Cyprioto-Taurian)




SPECIES *P. wachanrui* Levrat, 1858: 261

European Red List of Saproxyllic Beetles, does not include the species. It is known only from SE and E Anatolia. So, Turkish Red List category of the species is **NT**.

Range: Transcaucasia (Azerbaijan), Iran, Iraq, Turkey.


Chorotype: SW-Asiatic

**GENUS *CALCHAENESTHES* Kraatz, 1863: 97****SPECIES *C. diversicollis* Holzschuh, 1977: 129**

European Red List of Saproxyllic Beetles, does not include the species. It probably is distributed only in SE Anatolia for Turkey. So, Turkish Red List category of the species is **NE**.

Range: Iran, Iraq, Turkey.


Chorotype: SW-Asiatic

**SPECIES *C. oblongomaculata* (Guerin-Meneville, 1844: 234)**

According to European Red List of Saproxyllic Beetles, the species was placed in **DD**. It is known only from European Turkey for Turkey. So, Turkish Red List category of the species is **DD**.

Range: SE Europe (Bulgaria, Greece, Romania), Turkey, ? Middle East and ?Cyprus.


Chorotype: Turano-Mediterranean (Balkano-Anatolian)

**SPECIES *C. primis* Özdikmen, 2013: 150**

European Red List of Saproxyllic Beetles, does not include the endemic species rightly. It probably is rather widely distributed in Anatolia. So, Turkish Red List category of the species is **NT**.

Range: Turkey [? Middle East and ?Cyprus].

Chorotype: Anatolian or [E-Mediterranean (Palestino-Cyprioto-Taurian)]




TRIBE CALLICHRMATINI Swainson & Shuckard, 1840: 293**GENUS *AROMIA* Audinet-Serville, 1834: 559****SPECIES *A. moschata* (Linnaeus, 1758: 391)****SUBSPECIES *A. m. moschata* (Linnaeus, 1758: 391)**

According to European Red List of Saproxyllic Beetles, the species was placed in **LC**. It is represented only by the nominative subspecies in Turkey. It probably is widely distributed in N Turkey especially. So, Turkish Red List category of the subspecies is **LC**.

Range: Europe, Siberia, Kazakhstan, Mongolia, Caucasus, Transcaucasia (Georgia), Turkey.


Chorotype: Sibero-European

**SPECIES *A. ambrosiaca* (Steven, 1809: 40)****SUBSPECIES *A. a. ambrosiaca* (Steven, 1809: 40)**

European Red List of Saproxyllic Beetles, does not include the species. It is represented only by the nominative subspecies in Turkey. It probably is widely distributed in Anatolia. So, Turkish Red List category of the subspecies is **LC**.

Range: S Europe, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Turkey, Iraq, Syria, Lebanon, Jordan, North Africa (Algeria, Morocco, Tunisia).


Chorotype: Turano-Mediterranean

**GENUS *OSPHRANTERIA* Redtenbacher, 1850: 50****SPECIES *O. coerulescens* Redtenbacher, 1850: 50**

European Red List of Saproxyllic Beetles, does not include the species. It probably is rather widely distributed in E Anatolia. So, Turkish Red List category of the species is **NT**.

Range: Turkey, Iran, Iraq, Pakistan.

Chorotype: Turanian



Note: The conclusions and cited references for Turkish Cerambycinae will be presented at the end of evaluations in Part VIII.