

A ZOOGEOGRAPHICAL REVIEW OF SPONDYLIDINAE IN TURKEY (COLEOPTERA: CERAMBYCIDAE)

Hüseyin Özdikmen* and Semra Turgut*

* Gazi Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, 06500 Ankara, TÜRKİYE, e-mails: ozdikmen@gazi.edu.tr and semraturgut@mynet.com

[Özdikmen, H. & Turgut, S. 2006. A zoogeographical review of Spondylidinae in Turkey (Coleoptera: Cerambycidae). *Munis Entomology & Zoology* 1 (2): 279-288]

ABSTRACT: The taxa of Turkish Spondylidinae fauna that were recorded by various authors from different localities of Turkey are evaluated zoogeographically with the present paper. For this reason, the distribution patterns in Turkey and rest of the world for each taxon of related subfamily are determined. Also, chorotypes for each taxon are given in the parts of zoogeographical remarks.

Key Words: Coleoptera, Cerambycidae, Spondylidinae, Turkey, Zoogeography, Fauna

The longhorn beetles or Cerambycidae are classified together with Chrysomelidae and Bruchidae in the superfamily Chrysomeloidea. But, some authors recognized Cerambycidae as a separate superfamily Cerambycoidea (Svacha & Danilevsky, 1986). Cerambycidae sensu stricto is divided into several subfamilies. These are Parandrinae, Prioninae, Lepturinae, Necydalinae, Spondylidinae, Apatophyseinae, Cerambycinae and Lamiinae. Parandrinae are not represented in Turkey. Moreover, the subfamilies Necydalinae and Apatophyseinae are represented only a few species in Turkey.

The Turkish Spondylidinae fauna has not been evaluated as detailed by any authors yet. The works on the fauna of Spondylidinae in Turkey have been carried out especially towards late second half of the 20. century mostly faunistically. These are: Fairmaire (1884), Schimitschek (1944), Alkan (1946), Erdem (1947), Acatay (1948, 1961, 1963 and 1968), Defne (1954), Çanakçıoğlu (1956 and 1983), Demelt & Alkan (1962), Demelt (1963 and 1967), Villiers (1967), Erdem (1968), Beşçeli (1969), Gfeller (1972), Gül-Zümreoğlu (1972 and 1975), Tosun (1975), Erdem & Çanakçıoğlu (1977), Lobanov et al. (1981), Sekendiz (1981), Sama (1982 and 2002), Danilevsky & Miroshnikov (1985), Svacha & Danilevsky (1986), Öymen (1987), Önder et al. (1987), Adlbauer (1988 and 1992), Sama & Rapuzzi (1993), Yüksel (1996), Lodos (1998), Tuzin (2000), Alkan (2000), Tozlu (2001), Tozlu et al. (2002), Danilevsky (2005), Özdikmen & Çağlar (2004), Özdikmen & Şahin (2005), Malmusi & Saltini (2005), Özdikmen (2006). But, any work included the subfamily Spondylidinae completely is presently related to Turkish fauna. The known taxa of Spondylidinae in the studies above have been given together with other examined Cerambycidae taxa by various authors step by step. So nowadays, such data about the Turkish Spondylidinae fauna have reached a considerable level. The main aim of the present paper is to

evaluate the Turkish Spondylidinae fauna zoogeographically and faunistically by using known data.

In this paper classification and nomenclature of the longhorn beetles suggested by Danilevsky, 2005 and Althoff & Danilevsky, 1997 are followed. All genera are listed in the same order as in Danilevsky (2005), Althoff & Danilevsky (1997) and Sama (2002). Within the genera the species are listed alphabetically. Each name of a species or subspecies is accompanied by the author's name and description date. The data, "Current records from Turkey", "Range" and "Zoogeographical Remarks" under the title for each taxon is given in this present text. The present zoogeographical characterization is based on the chorotype classification of Anatolian fauna, recently proposed by Vigna Taglianti et al. (1999).

Subfamily Spondylidinae

Tribe Saphanini

Genus *Drymochares* Mulsant, 1847

Drymochares starcki Ganglbauer, 1888

The species has three subspecies in the World. These are nominate *Drymochares starcki starcki* Ganglbauer, 1888; *Drymochares starcki cavazzutii* Sama & Rapuzzi, 1993 and *Drymochares starcki ivani* Sama & Rapuzzi, 1993. *Drymochares starcki starcki* Ganglbauer, 1888 occurs only in Caucasus. *Drymochares starcki cavazzutii* is known from Abchazia, Armenia and north-eastern Turkey. *Drymochares starcki ivani* is only known from northern part of Turkey (Sama & Rapuzzi, 1993). These taxa may be distinguished by the different pubescence of elytra. Tauzin (2000) mentioned that Ilgaz Mountains in Kastamonu province was the hybridization area for the two subspecies occurring in Turkey.

Drymochares starcki cavazzutii Sama & Rapuzzi, 1993

Current records from Turkey: Artvin prov.: pass between from Borçka and Hopa (type locality); Turkey (Lobanov et al., 1981); Artvin prov.: from Borçka to Hopa (Sama, 1982); Northern Turkey (Danilevsky & Miroshnikov, 1985); Eastern Black Sea Region (Svacha & Danilevsky, 1986); Artvin prov.: Çifteköprüler and Murgul (Petek forest); Gümüşhane prov.: Zigana Mts. (Adlbauer, 1988; Sama & Rapuzzi, 1993); Giresun prov.: Dereci; Ordu prov.: Harçbeli; Hasancık; Gökçöy; Gürgentepe; Mesudiye; Tokat prov.: Niksar; Samsun prov.: Yenigün, Alaçam; Sinop prov.: Dranaz Mt., Yenikonak; Kastamonu prov.: Çatalzeytin, between Azdavay and Pınarbaşı, Ilgaz Mts. (Sama & Rapuzzi, 1993); Samsun prov.: Ladik and ?Layvikent (=Yakakent); Gümüşhane prov.: Şiran; Artvin prov.: Murgul; Ordu prov.: Akkuş; Kastamonu prov.: Ilgaz Mts. (Tauzin, 2000); Samsun prov.: Kavak, Hacılar pass; Artvin prov.: Cankurtaran pass (Malmusi & Saltini, 2005).

Range: Turkey, Caucasus (Sama & Rapuzzi, 1993).

Zoogeographical remarks: The subspecies is distributed in Turkey from Artvin to Kastamonu provinces in North Turkey. All old records before 1993 from Turkey should belong to this subspecies.

Drymochares starcki ivani Sama & Rapuzzi, 1993

Current records from Turkey: Bolu province: Abant (type locality) ; Bolu province (Tauzin, 2000); Bolu province: Abant; Zonguldak province: from Karadere to Eğerci (Malmusi & Saltini, 2005). **Range:** Turkey. **Zoogeographical remarks:** The subspecies is distributed only towards West of Kastamonu province in North Turkey. Up to now, it has been known from a rather local area in Turkey.

Genus *Saphanus* Serville, 1834

Saphanus piceus (Laicharting, 1784)

Current records from Turkey: There is no published record from Turkey. However, this species collected in Turkey is preserved in the collection of Stanislav Kadlec (Czechia) according to Danilevsky (2004). **Range:** Europe (Winkler, 1924-1932); Central and North of South Europe (Svacha & Danilevsky, 1986); Europe (Bense, 1985); Europe (Althoff & Danilevsky, 1997); Europe: from French Alps to Bulgaria and Greece (Sama, 2002); Europe

and Balkan Peninsula (Danilevsky, 2005). **Zoogeographical remarks:** The species is distributed only in Europe and probably European Turkey. It has the S-European chorotype.

Genus *Oxypleurus* Mulsant, 1839

***Oxypleurus nodieri* Mulsant, 1839**

Current records from Turkey: Antalya province: Bey Mts. (Demelt, 1963); Southern Turkey (Sama, 2002). **Range:** Mediterranean Region (Winkler, 1924-1932); Crimea, Europe, North Africa (Danilevsky & Miroshnikov, 1985); Mediterranean area (inc. Canary Islands), as far eastwards as the Black Sea, South Africa (Svacha & Danilevsky, 1986); Europe, North Africa (Bense, 1985); Europe (Althoff & Danilevsky, 1997); Southern Europe, Southern Turkey, Canary Islands, Madeira, North Africa, imported to South Africa (Sama, 2002); Crimea Peninsula, Caucasus, North Africa, South Europe, Balkan Peninsula (Danilevsky, 2005). **Zoogeographical remarks:** The species is distributed only in South part of Turkey. It has the Mediterranean chorotype.

Tribe Anisarthronini

Genus *Anisarthron* Redtenbacher, 1845

***Anisarthron barbipes* (Schrank, 1781)**

Current records from Turkey: The species has been reported only by Lodos (1998) from Turkey without exact locality. **Range:** Europe (Winkler, 1924-1932); Central Europe, up to Northern Italy and North of the Balkan Peninsula (Svacha & Danilevsky, 1986); Europe (Bense, 1985); Europe: from Italy to Ukraine (Althoff & Danilevsky, 1997); Western, Central and Eastern Europe (Sama, 2002); Europe and Balkan Peninsula (Danilevsky, 2005). **Zoogeographical remarks:** Unfortunately any record with exact locality in Turkey has not been given according to the literature yet. For this reason the real status in Turkey of this taxon needs to be clarified. It has the S-European chorotype.

Genus *Alocerus* Mulsant, 1862

***Alocerus moesiacus* (Frivaldsky, 1838)**

Current records from Turkey: Turkey (Svacha & Danilevsky, 1986); Isparta province: Eğirdir, Kovada (Adlbauer, 1992); Turkey (Lodos, 1998); Antalya province: Kumluca (Tozlu et al., 2002); Kırklareli province: İğneada, Mert lake (Özdikmen & Çağlar, 2004). **Range:** Mediterranean Region (Winkler, 1924-1932); Mediterranean Region (Svacha & Danilevsky, 1986); South Europe, North Africa (Bense, 1985); South Europe: from Portugal to Bulgaria and Greece (Althoff & Danilevsky, 1997). **Zoogeographical remarks:** The species is distributed in European Turkey and South Anatolia in Turkey. It has the Mediterranean chorotype.

Tribe Asemini

Genus *Nothorhina* Redtenbacher, 1845

***Nothorhina muricata* (Dalman, 1817)**

Current records from Turkey: The species has been reported as *N. punctata* by Lobanov et al., 1981 and Svacha & Danilevsky, 1986 from Turkey without exact locality. **Range:** Central and North Europe (Winkler, 1924-1932); Japan (Kusama & Takakuwa, 1984); Europe, South-West Siberia, Asia Minor, Japan (Svacha & Danilevsky, 1986); Europe, European Russia (Bense, 1985); Europe, European Russia, European Kazakhstan (Althoff & Danilevsky, 1997); Northern Palaearctic Region from Western Europe to Japan (Sama, 2002); Europe, Balkan Peninsula, Kazakhstan, West Siberia, Near East, Japan (Danilevsky, 2005). **Zoogeographical remarks:** Unfortunately any record with exact locality in Turkey has not been given according to the literatures yet. For this reason the real status in Turkey of this taxon needs to be clarified. It has the Sibero-European chorotype.

Genus *Aseum* Eschscholtz, 1830

***Aseum striatum* (Linnaeus, 1758)**

Current records from Turkey: Turkey (Acatay, 1948, 1961 and 1968); Sinop province: Hassan, 45 km SE Sinop (Villiers, 1967); Gümüşhane province: Torul Mts. (Gfeller, 1972); Gümüşhane province (Öymen, 1987); Tokat province: Topçam Mt. and Yıldız Mt., Almus, Tekneçik (Adlbauer, 1992); Turkey (Lodos, 1998); Kars province: Sarıkamış (Tozlu, 2001); Kars province: Sarıkamış (Tozlu et al., 2002); Northern Turkey (Sama, 2002). **Range:** Palaearctic Region (Winkler, 1924-1932); European Russia, Caucasus, Siberia, Europe, Mongolia, North-East China, Japan, Korea, Northern America (Danilevsky & Miroshnikov,

1985); Palaearctic Region, North America (Svacha & Danilevsky, 1986); Europe, European Russia (Bense, 1985); Europe, European Russia, European Kazakhstan (Althoff & Danilevsky, 1997); Europe, Northern Turkey, Caucasus, Siberia, Manchuria, Korea, Sakhalin, Japan, North America (Sama, 2002); Europe, Balkan Peninsula, Caucasus, Republics of Central Asia, Kazakhstan, West and East Siberia, Far East Russia, Sakhalin, Mongolia, Korea, Japan (Danilevsky, 2005). **Zoogeographical remarks:** Apparently the species is distributed only in North parts of Turkey. It has the Holarctic chorotype.

***Asemum tenuicorne* Kraatz, 1879**

Current records from Turkey: Ankara province: Kızılcahamam (Demelt, 1967); Turkey (Lobanov et al., 1981); Turkey (Danilevsky & Miroshnikov, 1985); Turkey (Svacha & Danilevsky, 1986); Turkey (Lodos, 1998); Asia Minor (Central and South-Eastern Turkey) (Sama, 2002). **Range:** Crimea, Caucasus, Syria (Winkler, 1924-1932); Crimea, Caucasus, Transcaucasia, Near East, Turkey (Danilevsky & Miroshnikov, 1985); Crimea, Caucasus, Transcaucasia, Turkey, Syria, Greece, ? Sweden (Svacha & Danilevsky, 1986); Italy, Greece, Rumania, Sweden, Crimea (Althoff & Danilevsky, 1997); Europe (incl. Sweden, Italy, Greece, Rumania, Crimea, Central and South-Eastern Turkey, West Caucasus, Northern Spain) (Sama, 2002); Crimea, Caucasus, Europe, Near East (Danilevsky, 2005). **Zoogeographical remarks:** Apparently the species is distributed in Central and South parts of Turkey. It has probably the Turano-European chorotype.

Genus *Arhopalus* Serville, 1834

***Arhopalus ferus* (Mulsant, 1839)**

Current records from Turkey: Hatay province: Akbez (Fairmaire, 1884); Bolu province: Abant as *C. polonicus* (Alkan, 1946); İzmir province: Central; Aydın province: Kuşadası (Demelt, 1963); Antalya province: Gebiz, Bozburan Mt. as *Arhopalus tristis* (Villiers, 1967); İzmir province: Bornova; Tokat province: Turhal (Gül-Zümrüoğlu, 1972 and 1975); Turkey as *Arhopalus tristis* Lobanov et al., 1981); Eskişehir province: Bokuyak (=?Bozüyük) as *Arhopalus tristis* (Sama, 1982); Çanakkale province: Gelibolu; Ankara province: Botanic Garden as *Arhopalus tristis* (Öymen, 1987); Çanakkale province: İntepe as *Arhopalus tristis* (Aldbauer, 1988); Tokat province: Turhal; İzmir province: Marmara Region, Aegean Region (Lodos, 1998); Adana province: Balcalı and Kozan; Antalya province: Central, Serik, Alanya, Topçam and Akseki; Bilecik province: Central; Kütahya province: Tavşanlı (Tozlu et al., 2002); Aydın province: Kuşadası; Kırklareli province: İslambeyli; Muğla province: Marmaris (Malmusi & Saltini, 2005); Turkey (Özdikmen & Şahin, 2006). **Range:** Europe, Siberia (Winkler, 1924-1932); European Russia, Caucasus, Transcaucasia, Siberia, Europe, Near East, Northern China, North Africa (Danilevsky & Miroshnikov, 1985); Most of Palaearctic Region (Svacha & Danilevsky, 1986); Europe, European Russia, North Africa (Bense, 1995); Europe, European Russia, European Kazakhstan (Althoff & Danilevsky, 1997); North Africa, Europe, Caucasus, Transcaucasia, Syria, Israel, Siberia, China (Sama, 2002); Europe, Balkan Peninsula, Crimea, Caucasus, Kazakhstan, West Siberia, Far East Russia, North Africa, Near East, China (Danilevsky, 2005). **Zoogeographical remarks:** Apparently the species is rather widely distributed in Turkey (especially West half of Turkey). It has the Sibero-European and the Turano-European-Mediterranean chorotypes.

***Arhopalus rusticus* (Linnaeus, 1758)**

Current records from Turkey: Between Ordu province and Bolu Mts. (Alkan, 1946); Sinop province: Ayancık (Schimitschek, 1944); Turkey (Acatay, 1948, 1961 and 1968); Tokat province: Arguslu (Villiers, 1967); Bolu province: Bolu Mts., Mengen (Demelt, 1967); Karabük province: Büyükdüz Research Forest (Beşçeli, 1969); Denizli province: Buldan (Gül-Zümrüoğlu, 1975); Antalya province: Manavgat (Karavca place), Korkuteli (Yazır forests), Serik (Belek forest) (Tosun, 1975); Bolu province, Sinop province: Ayancık, Gümüşhane province, Trabzon province: Akçaabat, Antalya province: Manavgat, Korkuteli, Serik, Denizli province (Erdem & Çanakçıoğlu, 1977); Gümüşhane province: Torul, Örümecek Serisi forests, Trabzon province: Akçaabat, around Düzköy (Sekendiz, 1981); Bolu province, Sinop province: Ayancık, Gümüşhane province, Trabzon province: Akçaabat, Antalya province: Manavgat, Korkuteli and Serik, Denizli province (Çanakçıoğlu, 1983); Northern Turkey (Danilevsky & Miroshnikov, 1985); Artvin province: Atilla Forest, Muğla province: Central, Balıkesir province: Dursunbey, Alaçam, İstanbul province: Büyükkada and Belgrad Forest (Öymen, 1987); Turkey (Önder et al., 1987); Kütahya province, Kastamonu province: Yarahgöz pass, Sinop province: Boyabat, Dıranaz pass, Rize province: İkizdere (Aldbauer,

1992); Artvin province: Şavşat (Yayla, Kocabey place) and (Veliköy, Karagöl Forests), Gümüşhane province: Karanlıkdere Forests (Yüksel, 1996); Gümüşhane province, Artvin province, Kars province, East Anatolian Region, Aegean Region, Western Black Sea Region, Mediterranean Region (Lodos, 1998); Artvin province: Ardanuç, Ovacık (Alkan, 2000); Kars province: Sarıkamış (Tozlu, 2001); Ankara province: Çamlıdere, Bayburt province: Central, Gümüşhane province: Central and Kürtün, Kars province: Sarıkamış (Tozlu et al., 2002); Turkey (Sama, 2002); Ordu province: Perşembe, Çaytepe Özdikmen, 2006); Karabük province: Büyükdüz Research Forest (Özdikmen & Şahin, 2006). **Range:** Europe (Winkler, 1924-1932); European Russia, Caucasus, Transcaucasia, Siberia, Europe, North Iran, North Turkey, Mongolia, Korea, Northern China, Japan, North Africa, North Amerika (Danilevsky & Miroshnikov, 1985); Most of Palaearctic Region, North America (Svacha & Danilevsky, 1986); Europe, European Russia, North Africa (Bense, 1995); Europe, European Russia, European Kazakhstan (Althoff & Danilevsky, 1997); North Africa, Europe, Turkey, Caucasus, Iran, Middle Asia, Siberia, Japan, North America, European Russia, European Kazakhstan (Sama, 2002); Europe, Balkan Peninsula, Crimea, Caucasus, Kazakhstan, West and East Siberia, Far East Russia, Sakhalin, Near East, Iran, Mongolia, Korea, Japan (Danilevsky, 2005). **Zoogeographical remarks:** Apparently the species is widely distributed in Turkey. It has the Holarctic chorotype.

***Arhopalus syriacus* (Reitter, 1895)**

Current records from Turkey: İzmir province: Kemalpaşa as *Criocephalus syriacus* (Demelt & Alkan, 1962); İzmir province: Kemalpaşa, Antalya province: Bey Mt. as *Criocephalus syriacus* (Demelt, 1963); Antalya province: Kepez forests (Tosun, 1975); Artvin province: Ardanuç Forests as *Criocephalus syriacus* (Sekendiz, 1981); Muğla province: Yeşilyurt (Adlbauer, 1992); Gaziantep province, Şanlıurfa province, İzmir province: Kemalpaşa as *Criocephalus syriacus* (Lodos, 1998); Adana province: Balcalı, Bağtepe and Kozan, Antalya province: Central (Tozlu et al., 2002); Nevşehir province: Cappadocia (Malmusi & Saltini, 2005); İçel province: Mut, Alahandüzü (Özdikmen & Şahin, 2006). **Range:** Syria (Winkler, 1924-1932); Mediterranean area, Canary Islands (Svacha & Danilevsky, 1986); South Europe (Bense, 1995); South Europe (Althoff & Danilevsky, 1997); Mediterranean Region (Sama, 2002). **Zoogeographical remarks:** Apparently it is very doubtful as a possible wrong identification. The status of this record must be revised. It has the Mediterranean chorotype.

Genus *Tetropium* Kirby, 1837

***Tetropium castaneum* (Linnaeus, 1758)**

Current records from Turkey: Turkey (Acatay, 1948, 1961 and 1968); Düzce province, Zonguldak province: Devrek, Western Black Sea Region (Erdem & Çanakçıoğlu, 1977, 1983); Düzce province: Çiçekli forest, Zonguldak province: Devrek, Beldibi, Davulgu forest, Zonguldak province: Devrek, Karadere as *Tetropium luridum* (Defne, 1954); North Anatolia as *Tetropium luridum* (Erdem, 1968); Bolu province: Seben (Öymen, 1987); Turkey (Önder et al., 1987); Artvin province: Atila Forests (Yüksel, 1996); Zonguldak province, Bolu province, Western Black Sea Region (Lodos, 1998); Artvin province: Central (Alkan, 2000). **Range:** Central and North Europe, Siberia (Winkler, 1924-1932); European Russia, Northern Caucasus, Siberia, Europe, Mongolia, Northern China, Korea, Northern Japan (Danilevsky & Miroshnikov, 1985); Most of Palaearctic Region, from Europe to Japan (Svacha & Danilevsky, 1986); Europe, European Russia (Bense, 1995); Europe, European Russia, European Kazakhstan (Althoff & Danilevsky, 1997); Northern Palearctic region from Europe to Japan (Sama, 2002); Europe, Balkan Peninsula, Crimea, Caucasus, Kazakhstan, West and East Siberia, Far East Russia, Sakhalin, Mongolia, China, Korea, Japan (Danilevsky, 2005). **Zoogeographical remarks:** Apparently it is distributed only in North parts of Turkey. It has the Sibero-European chorotype.

***Tetropium fuscum* (Fabricius, 1787)**

Current records from Turkey: Turkey (Acatay, 1948, 1961 and 1968); North Anatolia (Erdem, 1968); Artvin province: Şavşat (Veliköy, Karagöl Forests), (Karagöl, Meşeli, Mukali district), Ardanuç (Tepedüzü, Şahinkaya) (Yüksel, 1996); Turkey (Lodos, 1998); Artvin province: Şavşat, Karagöl (Alkan, 2000). **Range:** Central and North Europe, Siberia (Winkler, 1924-1932); Japan (Kusama & Takakuwa, 1984); European Russia, Georgia, West Siberia, Europe (Danilevsky & Miroshnikov, 1985); Europe, West Siberia (Svacha &

Danilevsky, 1986); Europe, European Russia (Bense, 1995); Europe, European Russia, European Kazakhstan (Althoff & Danilevsky, 1997); Europe, Western Siberia, European Kazakhstan (Sama, 2002); Europe, Balkan Peninsula, Caucasus, Kazakhstan, West Siberia, Japan (Danilevsky, 2005). **Zoogeographical remarks:** Apparently it is distributed only in North-East Turkey. It has the Sibero-European chorotype.

***Tetropium gabrieli* Weise, 1905**

Current records from Turkey: The species has only been reported by Acatay, 1948, 1961 and 1968 from Turkey without exact locality. **Range:** Central Europe (Winkler, 1924-1932); Central Europe and some adjacent regions, reaching to Denmark, Germany, France, Great Britain (Svacha & Danilevsky, 1986); Central and North Europe (Bense, 1995); Europe, ? North parts of European Russia (Althoff & Danilevsky, 1997); Europe from British Isles to Russia (Sama, 2002); Europe (Danilevsky, 2005). **Zoogeographical remarks:** Any record with exact locality in Turkey has not been given according to the literature yet. Unfortunately the records in Acatay (1948, 1961 and 1968) seem to be very doubtful. Because the species must not occur in Turkey according to the known distributional patterns in the world. Therefore Acatay's records are not valid. It has the Centralearctic chorotype.

Tribe Spondylidini

Genus *Spondylis* Fabricius, 1775

***Spondylis buprestoides* (Linnaeus, 1758)**

Current records from Turkey: Sinop province: Ayancık (Schimitschek, 1944); Kars province: Sarıkamış (Erdem, 1947); Turkey (Acatay, 1948, 1961, 1963 and 1968); Bursa province: Keles, Kocayayla place (Çanakçıoğlu, 1956); Trabzon province: Zigana Mts. (Villiers, 1967); Ankara province: Kızılcahamam (Demelt, 1967); Karabük province: Büyükdüz Research Forest (Beşçeli, 1969); Turkey (Lobanov et al., 1981); Turkey (Danilevsky & Miroshnikov, 1985); Trabzon province: Maçka (Öymen, 1987); Artvin province: Ardanuç (Yüksel, 1996); Turkey (Lodos, 1998); Artvin province: Ardanuç, Tosunlu (Alkan, 2000); Kars province: Sarıkamış (Tozlu, 2001); Kars province: Sarıkamış (Tozlu et al., 2002); Northern and South-Western Turkey (Sama, 2002); İstanbul province: Şile (Malmusi & Saltini, 2005); Ankara province: Kızılcahamam, Çamkoru (Özdikmen & Şahin, 2006). **Range:** Palaearctic Region (Winkler, 1924-1932); European Russia, Caucasus, Transcaucasia, Siberia, Europe, Turkey, Northern Iran, Mongolia, Northern China, Korea, Japan (Danilevsky & Miroshnikov, 1985); Most of Palaearctic Region, from Europe to Japan (Svacha & Danilevsky, 1986); Europe, European Russia (Bense, 1995); Europe, European Russia, European Kazakhstan (Althoff & Danilevsky, 1997); Europe, Northern and South-Western Turkey, Siberia, Sakhalin, Northern Mongolia, Northern China, Korea, Japan, ? North Africa (Sama, 2002); Europe, Balkan Peninsula, Crimea, Caucasus, Kazakhstan, West and East Siberia, Far East Russia, Sakhalin, Mongolia, China, Korea, Japan (Danilevsky, 2005). **Zoogeographical remarks:** Apparently it is distributed only in North parts of Turkey. It has the Sibero-European chorotype.

DISCUSSION

The Turkish Spondylidinae subfamily is represented by 4 tribes as Saphanini, Anisarthonini, Asemini and Spondylidini. The tribe Saphanini includes 3 genera as Drymochares, Saphanus and Oxyleurus. The tribe Anisarthonini includes 2 genera as Anisarthon and Alocerus. The tribe Asemini includes 4 genera as Nothorhina, Asemum, Arhopalus and Tetropium. The tribe Spondylidini includes only one genus as Spondylis normally.

In the tribe Saphanini, the genus Drymochares is represented by 2 subspecies as *D. s. cavazzutii* and *D. s. ivani* of the species *Drymochares starcki*. The other two genera, Saphanus and Oxyleurus are represented

by one each species as *Saphanus piceus* (Laicharting, 1784) and *Oxypleurus nodieri* Mulsant, 1839 respectively.

In the tribe Anisarthronini, both genera *Anisarthron* and *Alocerus* are represented by one species each as *Anisarthron barbipes* (Schrank, 1781) and *Alocerus moesiacus* (Fruvaldsky, 1838) respectively.

In the tribe Spondylidini, the genus *Spondylis* is represented by the species *Spondylis buprestoides* (Linnaeus, 1758).

In the tribe Asemini, the genus *Nothorhina* is represented by the species *Nothorhina muricata* (Dalman, 1817). The genus *Asemum* is represented by 2 species as *Asemum striatum* (Linnaeus, 1758) and *Asemum tenuicorne* Kraatz, 1879. The genus *Arhopalus* is represented by 3 species as *Arhopalus fesus* (Mulsant, 1839), *Arhopalus rusticus* (Linnaeus, 1758) and *Arhopalus syriacus* (Reitter, 1895). According to literatures, the genus *Tetropium* has been represented by 3 species as *Tetropium castaneum* (Linnaeus, 1758), *Tetropium fuscum* (Fabricius, 1787) and *Tetropium gabrieli* Weise, 1905. But, as seen in the text, *Tetropium gabrieli* has only been known by Acatay's records from Turkey. However, Acatay's records are wrong. We think the species must not occur in Turkey according to its known distribution patterns in the world.

Consequently, the Turkish Spondylidinae subfamily includes 4 tribes, 10 genera, 14 species and 2 subspecies until now according to the present paper. Taking into consideration the status of the species *Tetropium gabrieli*, the known real taxa number of Turkish Spondylidinae is 15 as species and subspecies. Moreover, Sama (pers. com.) mentioned that *Anisarthron barbipes*, *Saphanus piceus* and all species of the genus *Tetropium* are very likely absent in Turkey. Under these circumstances, Turkish Spondylidinae includes 4 tribes, 7 genera, 9 species and 2 subspecies.

Zoogeographically, *Asemum striatum* (Linnaeus, 1758) and *Arhopalus rusticus* (Linnaeus, 1758) have the Holarctic chorotype. *Oxypleurus nodieri* Mulsant, 1839, *Alocerus moesiacus* (Fruvaldsky, 1838) and *Arhopalus syriacus* (Reitter, 1895) have the Mediterranean chorotype. *Saphanus piceus* (Laicharting, 1784) and *Anisarthron barbipes* (Schrank, 1781) have the S-European chorotype. *Nothorhina muricata* (Dalman, 1817), *Tetropium castaneum* (Linnaeus, 1758), *Tetropium fuscum* (Fabricius, 1787) and *Spondylis buprestoides* (Linnaeus, 1758) have the Sibero-European chorotype. *Arhopalus fesus* (Mulsant, 1839) has the Sibero-European and the Turano-Europeo-Mediterranean chorotypes. *Asemum tenuicorne* Kraatz, 1879 has probably the Turano-European chorotype. *Tetropium gabrieli* Weise, 1905 has the Centraleuropean chorotype. The subspecies *Drymochares starcki cavazzutii* Sama & Rapuzzi, 1993 has been known from Turkey and Caucasus. The other subspecies *Drymochares starcki ivani* Sama & Rapuzzi, 1993 are known only from Turkey yet.

ACKNOWLEDGEMENTS

Thank to Dr. Gianfranco Sama (Italy) for his review and comments.

LITERATURE CITED

Acatay, A. 1948. Zararlı orman böcekleri teşhis anahtarı. T. C. Tarım Bakanlığı Orman Genel Müdürlüğü yayınları, Özel Sayı: 76, Ocak Matbaası, İstanbul, Türkiye, 113 pp.

Acatay, A. 1961. Zararlı orman böcekleri teşhis anahtarı. İstanbul Üniversitesi Orman Fakültesi yayınları No: 76, Fakülteler Matbaası, İstanbul, Türkiye, 152 pp.

Acatay, A. 1963. Tatbiki Orman Entomolojisi. İstanbul Üniversitesi Orman Fakültesi yayınları No: 94, Fakülteler Matbaası, İstanbul, Türkiye, 169 pp.

Acatay, A. 1968. Zararlı orman böcekleri teşhis anahtarı. İstanbul Üniversitesi Orman Fakültesi yayınları No: 132, Kutulmuş Matbaası, İstanbul, Türkiye, 154 pp.

Adlbauer, K. 1988. Neues zur Taxonomie und Faunistik der Bockkäferfauna der Türkei (Coleoptera, Cerambycidae). Entomofauna, 9 (12): 257-297.

Adlbauer, K. 1992. Zur Faunistik und Taxonomie der Bockkäferfauna der Türkei II (Coleoptera, Cerambycidae). Entomofauna, 13 (30): 485-509.

Alkan, B. 1946. Agricultural Entomology, T.C. Ministry of Agriculture, Institute of High Agriculture of Ankara, Book No: 31, Ankara, Turkey (Turkish with English summary).

Alkan, H. 2000. A study on the Cerambycidae species in Northern BlackSea Region and the introduction of Turkish forest Cerambycidae (Insecta, Coleoptera). K.T.Ü., Institute of Science, Engineering of Forestry, Trabzon, Turkey (Turkish).

Althoff, J. & Danilevsky, M. L. 1997. A Check-List of Longicorn Beetles (Coleoptera, Cerambycoidea) of Europe. Slovensko Entomološko Društvo Štefana Michielija. Ljubljana, 64 pp.

Bense, U. 1995. Illustrated key to the Cerambycidae (excl. Dorcadionini) and Vesperidae of Europe. Margraf Verlag, Germany, 512 pp.

Besçeli, Ö. 1969. Biologies des insectes nuisibles dans la forest de recherches de Büyükdüz et mesures de protection, Publications of Institute of Forestry Research, No: 33, Ankara, 94 pp.

Çanakçıoğlu, H. 1956. Entomological researches in the forests of Bursa. İ. Ü. Publication of Forest Faculty, No : 41, 41 pp., İstanbul, Turkey (Turkish with English summary).

Çanakçıoğlu, H. 1983. Forest Entomology: Special volume, İ.Ü. Publication of Forest Faculty, No: 349, İstanbul, Turkey (Turkish with English summary).

Danilevsky, M. L. 2005. Systematic list of Longicorn Beetles (Cerambycoidea) of the territory of the former USSR (on the base of "Systematic list of Longicorn Beetles of the USSR" by A.L. Lobanov, M.L. Danilevsky, S.V. Murzin, 1981, and computer databases by A.L. Lobanov, 1979-1990). Available from: <http://www.zin.ru/Animalia/Coleoptera/eng/dbase30.htm>

Danilevsky, M. L. & Miroshnikov, A. I. 1985. The longicorn beetles of the Caucasus (Coleoptera, Cerambycidae) A key book. Krasnodar, USSR, 419 pp.

- Defne, M. Ö.** 1954. Batı Karadeniz Bölgesindeki Göknarların Zararlı Böcekleri ve Mücadele Metodları. T. C. Tarım Bakanlığı, Orman Genel Müdürlüğü Yay. No. 105, 60-72.
- Demelt, C. V. & Alkan, B.** 1962. Short information of Cerambycidae Fauna of Turkey. Bitki Koruma Bülteni, 2 (10): 49-56.
- Demelt, C. V.** 1963. Beitrag zur Kenntnis der Cerambycidenfauna Kleinasien und 13. Beitrag zur Biologie palaearkt. Cerambyciden, sowie Beschreibung einer neuen Oberea-Art. Entomologische Blätter, 59 (3) : 132-151.
- Demelt, C. V.** 1967. Nachtrag zur Kenntnis der Cerambyciden-Fauna Kleinasien. Entomologische Blätter, 63 (2): 106-109.
- Erdem, R.** 1947. Sarıkamış Ormanlarında Entomolojik Gözlemler, T.C. Tarım Bakanlığı, Orman Genel Müdürlüğü Yayınları, Özel Sayı: 52, Ankara.
- Erdem, R.** 1968. Useful and Harmful Insects of Forest, İstanbul Univ. Publication of Forest Faculty, 43-58, İstanbul, Turkey (Turkish with English summary).
- Erdem, R. & Çanakçıoğlu, H.** 1977. Pests of Wood in Turkey, İstanbul Univ. Publication of Forest Faculty, 113-134, İstanbul, Turkey (Turkish with English summary).
- Fairmaire, M. L.** 1884. Liste des Coleopteres recueillis par M. l'abbé David a Akbes (Asie-Mineure) et Descriptions des Especies Nouvelles. Annales de la Societe entomologique de France, Octobre (1884): 165-180.
- Gfeller, W.** 1972. Cerambycidae (Coleoptera) der Türkei-Persienexpedition 1970 der Herren Dr. H. c. W. Wittmer und U. v. Botmer. Mitteilungen der Entomologischen Gesellschaft Basel, 22 (1): 1-8.
- Gül-Zümreoğlu, S.** 1972. Catalogue of Insect and common pests (1928-1969). T. C. Publications of Agriculture Ministry, Bornova, İzmir, 119 pp.
- Gül-Zümreoğlu, S.** 1975. Investigations on taxonomy, host plants and distribution of the Longhorned Beetles (Cerambycidae-Coleoptera) in Aegean Region. T. C. Ministry of Food, Agriculture and Stockbreeding, No : 28, , İstiklal Press, İzmir, 208 pp.
- Kusama, K. & Takakuwa, M.** 1984. Parandrinae. Prioninae. Spondyliinae. Aseminae. Lepturinae (part.). Cerambycinae. Lamiinae (part.). The Longicorn-beetles of Japan in Color. Kodansha, Tokyo: 131-172, 201-493, 511-549.
- Lobanov, A. L., Danilevsky, M. L. & Murzin, S. V.** 1981. [Systematic list of Longicorn beetles (Coleoptera. Cerambycidae) of the USSR. 1]. Rev. d'Entom. 60 (4): 784-803 [in Russian].
- Lodos, N.** 1998. Entomology of Turkey VI (General, Applied and Faunistic). Ege Ü. Ziraat Fak. Yayınları No: 529, E. Ü. Faculty of Agriculture Press, İzmir, 300 pp.
- Malmusi, M. & Saltini, L.** 2005. Cerambycidae raccolti dai componenti del Gruppo Modenese Scienze Naturali durante escursioni in Turchia tra il 1987-2003 (Contributo alla Fauna dei Cerambycidae di Turchia). Quaderno di studi e notizie di storia naturale della Romagna, n. 21, 28 pp. (unpublished).
- Önder, F., Karsavuran, Y., Tezcan, S. & Önder, P.** 1987. Scientific and Turkish names of some useful and harmful species of Agricultural, Forestic and Domestic Animals in Turkey, T.C. Ministry of Agriculture, Ankara, Turkey (Turkish with English summary).
- Öymen, T.** 1987. The Forest Cerambycidae of Turkey. İ. Ü. Forest Faculty, İstanbul, 146 pp.

- Özdikmen, H.** 2006. Contribution to the knowledge of Turkish longicorn beetles fauna (Coleoptera: Cerambycidae). *Munis Entomology & Zoology*, 1 (1): 71-90.
- Özdikmen, H. & Çağlar, Ü.** 2004. Contribution to the Knowledge of Longhorned Beetles (Coleoptera, Cerambycidae) from Turkey, Subfamilies Prioninae, Lepturinae, Spondylidinae and Cerambycinae. *Journal of the Entomological Research Society*, 6 (1): 39-69.
- Özdikmen, H. & Şahin, Ö.** 2006. Longhorned beetles collection of the entomology museum of Central Anatolia forestry research directorship, Ankara, Turkey (Coleoptera, Cerambycidae). *G. U. Journal of Science* 19 (1): 1-8.
- Sama, G.** 1982. Contributo allo studio dei coleotteri Cerambycidae di Grecia e Asia Minore. *Fragmenta Entomologica*, Roma, 16 (2): 205-227.
- Sama, G.** 2002. Atlas of the Cerambycidae of Europe and the Mediterranean Area, Volume I, Kabourek, Zlin, 173 pp.
- Sama, G. & Rapuzzi, P.** 1993. Revisione dei generi *Saphanus* Serville, 1834 e *Drymochares* Mulsant, 1847. *Lambillionea*, 93: 278-294.
- Schmitschek, E.** 1944. Forstinsekten der Türkei und Ihre Umwelt Grundlagen der türkischen Forstentomologie, Volk und Reich Verlag Prag, 125-141 pp.
- Sekendiz, O. A.** 1981. Researches on important damaging animal species of Eastern Black Sea Region. K. T. Ü. Publication of Forest Faculty, No : 12, Trabzon, 114 pp.
- Svacha, P. & Danilevsky, M. L.** 1986. Cerambycoid larvae of Europe and Soviet Union (Coleoptera, Cerambycoidea). Part I. *Acta Universitatis Carolinae – Biologica*, 30 : 1-176.
- Taglianti, A. V., Audisio, P. A., Biondi, M., Bologna, M. A., Carpaneto, G. M., De Biase, A., Fattorini, S., Piattella, E., Sindaco, R., Venchi, A. & Zapparoli, M.** 1999. A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palaearctic Region. *Biogeographia* 20: 31-59.
- Tauzin, P.** 2000. Complement a l'inventaire des Coleopteres Cerambycidae de Turquie. *L'Entomologiste*, 56 (4): 151-153.
- Tosun, İ.** 1975. The damaging insect species at conifer forests in Mediterranean Region and researches on their predators and parasites. *İ. Ü. Orman Fakültesi Dergisi*, 26 (2): 218-254.
- Tozlu, G.** 2001. Sarıkamış (Kars) Ormanlarında Sarıçam (*Pinus sylvestris* L.)'da zarar yapan Elateridae, Buprestidae, Cerambycidae, Curculionidae (Coleoptera) ve Diprionidae (Hymenoptera) familyalarına bağlı türler üzerinde çalışmalar. *Türkiye Entomoloji Dergisi* 25(2): 133-146.
- Tozlu, G., Rejzek, M. & Özbek, H.** 2002. A contribution to the knowledge of Cerambycidae (Coleoptera) fauna of Turkey. Part I: Subfamilies Prioninae to Cerambycinae. *Biocosme Mèsoèèn*, Nice, 19 (1-2): 55-94.
- Villiers, A.** 1967. Coléoptères Cérambycides de Turquie (1. Partie) - *L' Entomologiste*, 23 (1): 18-22.
- Winkler, A.** 1924-1932. *Catalogus Coleopterorum regionis palaearticae*. Verlag von Albert Winkler, 1135-1226.
- Yüksel, B.** 1996. The damaging insect species at oriental spruce forests and their predators and parasite species –I (Pest Insects), K.T.Ü. Institute of Science, Engineering of Forest, Trabzon, 222 pp. (Turkish with English summary).

[Received December 2005. Accepted March 2006]