

Faunae Insectorum Germanicae

Initia

Deutschlands Insecten

herausgegeben

von

Dr. G. W. F. Panzer

VIIItes Heft.

Nürnberg

in der Felseckerschen Buchhandlung.
Preis 12. gg. oder 54. Kr. R. Münz.

29

Bibliothek

Landhous Hogeschool
WAGENINGEN

Stabat
Walden

LEPTVRA arcuata.

Der bogrichte Schmalbock.

Leptura arcuata: coleoptris attenuatis testaceis fasciis quatuor nigris, prima triangulari, antennis pedibusque flavis.

Habitat in sylvaticis Harcyniae, Austriae.

Statura *Lept. subspinosae* F. at paullo minor. Caput nigrum, ore flavo. Antennae (maris) nigrae, articulis sex ultimis flavis. Thorax pube aurea nitet. Elytra attenuata, testacea, fasciis quatuor, prima triangulari, reliquis simplicibus punctoque anguli antici nigris. Corpus subtus aureo-villosum. Pedes in femina flavii; in mare supra nigri, subtus flavii.

♂ Mas. ♀ Femina, magnitudine naturali. * Coleopteron lente auctum.



Leptura arcuata Hellw.

LEPTVRA signata.

Der gezeichnete Schmalbock.

Leptura signata; nigra, elytris lividis, punctis, fasciisque duabus interruptis nigris, pedibus ruffis.

Habitat in Harcyniae, Göttingae sylvaticis.

Statura et magnitudo *L. melanur*. Corpus cum capite et thorace nigrum. Antennae flavae articulis apice nigricantibus. Elytra fastigiata laevia, livida, fasciis medio duabus interruptis, s. suturam haud attingentibus, punctis versus apicem plurimis, et maculis ad apicem oblongis nigris. Pedes ruffi genubus atris.

a. Coleoptron lente auctum.



Leptura signata Hellw.